Homework 1

- Questions from Lecture unit 1
- Please refer to the lecture slides for more information about the context in which the questions arise.
- Due date: September 4, 2019

Questions

- 1. Why do you think keyed cryptography is more practical nowadays?
- 2. Shift cipher can be generalized where the encryption function is of the form

$$E(x) = ax + b \mod 26$$

for $x \in \{0,...25\}$, where $a, b \in \mathbb{Z}$. The encryption function needs to be injective (one-to-one) so that every ciphertext can be uniquely decrypted. What conditions should be posed on a and b in order for E to be injective?

- 3. Is $\Gamma=\begin{pmatrix}7&14\\3&23\end{pmatrix}$ invertible? If so, what is Γ^{-1} (so that $\Gamma^{-1}\Gamma=\begin{pmatrix}1&0\\0&1\end{pmatrix}$?
- 4. Let's say it takes a second to try a key, that is an m by m matrix Γ , where we will need to construct its "inverse" Γ^{-1} and use it to try to decrypt a ciphertext. Supposedly we may need to try all possible m by m matrices. How small can m be so that it will take us at least 1000 years to try all keys in order to break the cipher?
- 5. Which do you think is a stronger evidence for security?
 - (a) Matrix-cipher: exhaustively trying all possible keys takes too much time (26^{m^2}) .
 - (b) RSA: Breaking the system may require factoring integers, a problem believed to be computationally hard.
- 6. Suppose

- (a) Pr[b=1] = 2/3 and Pr[b=0] = 1/3.
- (b) r is generated with the following probability, independently of what b is: Pr[r=1] = Pr[r=0] = 1/2.

Questions:

- (a) What is the probability that $c = b \wedge r$ is 1?
- (b) What is the probability that $c = b \wedge r$ is 0?
- (c) Is it a good idea to hide b in c using the \wedge operation?