GE23131-Programming Using C-2024



Status	Finished
Started	Monday, 13 January 2025, 1:57 PM
Completed	Monday, 13 January 2025, 2:13 PM
Duration	15 mins 57 secs



You are transporting some boxes through a tunnel, where each box is a parallelepiped, and is charac its length, width and height.

The height of the tunnel 41 feet and the width can be assumed to be infinite. A box can be carried the tunnel only if its height is strictly less than the tunnel's height. Find the volume of each box that can successfully transported to the other end of the tunnel. Note: Boxes cannot be rotated.

Input Format

The first line contains a single integer n, denoting the number of boxes.

n lines follow with three integers on each separated by single spaces - $length_i$, $width_i$ and $height_i$ w length, width and height in feet of the i-th box.

Constraints

 $1 \le n \le 100$

 $1 \le length_i$, width_i, height_i ≤ 100

Output Format

For every box from the input which has a height lesser than 41 feet, print its volume in a separate lin

Sample Input 0

4

5 5 5

1 2 40

10 5 41

7 2 42

Sample Output 0

125

80

Explanation 0

The first box is really low, only 5 feet tall, so it can pass through the tunnel and its volume is $5 \times 5 \times 10^{-5}$

The second box is sufficiently low, its volume is $1 \times 2 \times 4 = 80$.

The third box is exactly 41 feet tall, so it cannot pass. The same can be said about the fourth box.

Input	Expected	Got	
4 5 5 5 1 2 40 10 5 41 7 2 42	125 80	125 80	

Passed all tests!

Question **2**Correct

Flag

question

You are given n triangles, specifically, their sides a_i , b_i and c_i . Print them in the same style but sorted areas from the smallest one to the largest one. It is guaranteed that all the areas are different.

The best way to calculate a volume of the triangle with sides \boldsymbol{a} , \boldsymbol{b} and \boldsymbol{c} is Heron's formula:

$$S = \ddot{O} p * (p - a) * (p - b) * (p - c)$$
 where $p = (a + b + c) / 2$.

Input Format

First line of each test file contains a single integer n. n lines follow with a_i , b_i and c_i on each separate spaces.

Constraints

 $1 \le n \le 100$

 $1 \leq a_i, b_i, c_i \leq 70$

 $a_i + b_i > c_i$, $a_i + c_i > b_i$ and $b_i + c_i > a_i$

Output Format

Print exactly n lines. On each line print 3 integers separated by single spaces, which are a_i , b_i and c_i corresponding triangle.

Sample Input 0

3

7 24 25

5 12 13

3 4 5

Sample Output 0

REC-CIS


5 12 13

7 24 25

Explanation 0

The square of the first triangle is **84**. The square of the second triangle is **30**. The square of the third is **6**. So the sorted order is the reverse one.

Answer: (penalty regime: 0 %)



Passed all tests!

Save the state of the flags