

## GE23131-Programming Using C-2024

## Quiz navigation

[Show one page at a time](#)[Finish review](#)

Status	Finished
Started	Monday, 23 December 2024, 5:33 PM
Completed	Saturday, 14 December 2024, 6:20 PM
Duration	8 days 23 hours

## Question 1

Correct

Marked out of 5.00

☐ Flag question

Sunny and Johnny like to pool their money and go to the ice cream parlor. Johnny never buys the same flavor that Sunny does. The only other rule they have is that they spend all of their money.

Given a list of prices for the flavors of ice cream, select the two that will cost all of the money they have.

For example, they have  $m = 6$  to spend and there are flavors costing  $cost = [1, 2, 3, 4, 5, 6]$ . The two flavors costing 1 and 5 meet the criteria. Using 1-based indexing, they are at indices 1 and 4.

**Function Description**

Complete the code in the editor below. It should return an array containing the indices of the prices of the two flavors they buy.

It has the following:

- $m$ : an integer denoting the amount of money they have to spend
- $cost$ : an integer array denoting the cost of each flavor of ice cream

**Input Format**

The first line contains an integer,  $t$ , denoting the number of trips to the ice cream parlor. The next  $t$  lines each describe a visit. Each trip is described as follows:

1. The integer  $m$ , the amount of money they have pooled.
2. The integer  $n$ , the number of flavors offered at the time.
3.  $n$  space-separated integers denoting the cost of each flavor:  $cost[1], cost[2], \dots, cost[n]$

**Note:** The index within the cost array represents the flavor of the ice cream purchased.

**Constraints**

- $1 \leq t \leq 50$
- $2 \leq m \leq 10^4$
- $2 \leq n \leq 10^4$
- $1 \leq cost[i] \leq 10^4, \forall i \in [1, n]$
- There will always be a unique solution.

**Output Format**

For each test case, print two space-separated integers denoting the indices of the two flavors purchased in ascending order.

**Sample Input**

5  
1 4 5 3 2  
4  
4  
2 2 4 3

Sample Output

1 4  
1 2

Explanation

Sunny and Johnny make the following two trips to the parlor:

- 1. The first time, they pool together  $m = 4$  dollars. Of the five flavors available that day, flavors **1** and **3** have a total cost of  $1 + 3 = 4$ .
- 2. The second time, they pool together  $m = 4$  dollars. Of the four flavors available that day, flavors **1** and **2** have a total cost of  $2 + 2 = 4$ .

Answer: (penalty regime: 0 %)

	Input	Expected	Got	
	2	1 4	1 4	
	4	1 2	1 2	
	5			
	1 4 5 3 2			
	4			
	4			
	2 2 4 3			

Passed all tests!

Question **2**

Correct

Marked out of 5.00

☐ Flag question

Numeros the Artist had two lists that were permutations of one another. He was very proud. Unfortunately while transporting them from one exhibition to another, some numbers were lost out of the first list. find the missing numbers?

As an example, the array with some numbers missing,  $arr = [7, 2, 5, 3, 5, 3]$ . The original array of numbers was  $arr = [7, 2, 5, 4, 6, 3, 5, 3]$ . The numbers missing are  $[4, 6]$ .

Notes

lists is the same. If that is not the case, then it is also a missing number.

- You have to print all the missing numbers in ascending order.
- Print each missing number once, even if it is missing multiple times.
- The difference between maximum and minimum number in the second list is less than or equal to 101.

Complete the code in the editor below. It should return an array of missing numbers.

It has the following:

- `arr`: the array with missing numbers
- `brr`: the original array of numbers

### Input Format

There will be four lines of input:

***n*** - the size of the first list, ***arr***

The next line contains ***n*** space-separated integers ***arr[i]***

***m*** - the size of the second list, ***brr***

The next line contains ***m*** space-separated integers ***brr[i]***

### Constraints

- $1 \leq n, m \leq 2 \times 10^5$
- $n \leq m$
- $1 \leq brr[i] \leq 2 \times 10^4$
- $X_{max} - X_{min} < 101$

### Output Format

Output the missing numbers in ascending order.

### Sample Input

```
10
203 204 205 206 207 208 203 204 205 206
13
203 204 204 205 206 207 205 208 203 206 205 206 204
```

### Sample Output

```
204 205 206
```

### Explanation

**204** is present in both arrays. Its frequency in ***arr*** is **2**, while its frequency in ***brr*** is **3**. Similarly, **205** and **206** occur twice in ***arr***, but three times in ***brr***. The rest of the numbers have the same frequencies in both lists.

**Answer:** (penalty regime: 0 %)

Question **3**

Correct

Marked out of 5.00

☐ Flag question

	Input	Expected	Got	
	10 203 204 205 206 207 208 203 204 205 206 13 203 204 204 205 206 207 205 208 203 206 205 206 204	204 205 206	204 205 206	

Passed all tests!

Watson gives Sherlock an array of integers. His challenge is to find an element of the array such that all elements to the left is equal to the sum of all elements to the right. For instance, given the array `a = [8, 11]`, `8` is between two subarrays that sum to `11`. If your starting array is `[11]`, that element satisfies the left and right sum to `0`.

You will be given arrays of integers and must determine whether there is an element that meets the criterion.

Complete the code in the editor below. It should return a string, either YES if there is an element meeting the criterion or NO otherwise.

It has the following:

- `arr`: an array of integers

Input Format

- The first line contains `T`, the number of test cases.
- The next `T` pairs of lines each represent a test case.
- The first line contains `n`, the number of elements in the array `arr`.
  - The second line contains `n` space-separated integers `arr[i]` where  $0 \leq i < n$ .

Constraints

- $1 \leq T \leq 10$
- $1 \leq n \leq 10^5$
- $1 \leq arr[i] \leq 2 \times 10^4$
- $0 \leq i \leq n$

Output Format

For each test case print YES if there exists an element in the array, such that the sum of the elements on its left is equal to the sum of the elements on its right; otherwise print NO.

Sample Input 0

REC-CIS

3  
1 2 3  
4  
1 2 3 3

**Sample Output 0**

NO  
YES

**Explanation 0**

For the first test case, no such index exists.

For the second test case,  $arr[0] + arr[1] = arr[3]$ , therefore index **2** satisfies the given conditions.

**Sample Input 1**

3  
5  
1 1 4 1 1  
4  
2 0 0 0  
4  
0 0 2 0

**Sample Output 1**

YES  
YES  
YES

**Explanation 1**

In the first test case,  $arr[2] = 4$  is between two subarrays summing to **2**.

In the second case,  $arr[0] = 2$  is between two subarrays summing to **0**.

In the third case,  $arr[2] = 2$  is between two subarrays summing to **0**.

**Answer:** (penalty regime: 0 %)

REC-CIS

3	YES	YES	
5	YES	YES	
1 1 4 1 1	YES	YES	
4			
2 0 0 0			
4			
0 0 2 0			
2	NO	NO	
3	YES	YES	
1 2 3			
4			
1 2 3 3			

Passed all tests!

Save the state of the flags