

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

<u>Chemical Composition :</u>	Content % w/w
Rhodopol 23 as 5% Gel -	2.5 % w/w
Demineralized water -	QS to make 100% -
Sodium Bentonite -	0.25 % w/w
TOTAL -	100.0 % w/w
Kresoxim-methyl a.i. content -	44.3 % w/w
Polydimethyl siloxane (Defoamer) -	1.0 % w/w
Ethoxylated derivative of Styrenated Phenol -	1.5 % w/w
Ethoxylated polymethacrylate in Propylene glycol and water -	4.0 % w/w
Trihydroxy Propane (Glycerol) -	5.0 % w/w
Blend of Methyl isothiazolone and its chloro derivative -	0.5 % w/w
Total:	100.00%

GÙÊ

xiÀ 9 ' - \$

KỶ OÙ RỪNG KỶ Ỗ I NỖ K

K ũ Š ~~BO~~Ž æ Yl

there is no specific antidote, hence symptomatic and supportive therapy should be given.

Manufactured by

Rallis India Limited
Rallis India Limited, IIIRD Block, IInd Floor, KSCMF Ltd., Building, 8,
Cunningham Road, Bangalore - 560052 Rallis India Limited
156/157, Nariman Bhavan, 15th Floor, 228, Nariman Point, Mumbai
- 400021 Bangalore Bangalore Karnataka 560052

Premises Address

- (1). M/s Rallis India Limited, Plot No. 3301, GIDC Industrial Area,
Dist. Bharuch, Ankleshwar-393 002, GUJARAT
Gujarat
- (2). M/s Rallis India Limited, C-5/C-6, MIDC Industrial Area,Phase-
III, Shivani, Akola ĩ 444 104, MAHARASTHRA
Maharashtra
- (3). Rallis India Ltd Plot No. C44, Chemical Zone, Dahej Taluka-
Vagra, District- Bharuch - 392130, GUJARAT
- (4). M/s Rallis India Limited, Plot No. Z-110, Dahej SEZ Part II, P.O.
Lakhigam, Tal ĩ Vagra, Dist : Bharuch 392 130, GUJARAT

Customer Care Details

-

Safety Pictograms:



00YŃÜ¼ 9

¾ı| 0ÜeÜĤÈ 0ÜYß0 ©YĐ Ńôßußß] Üà ĩ ¾ı yŃPj0 ©0 «ÄŃ¾ēPj ¾ı
z Əj0 00 {ÜjĀ Īğ ĀjŃğ Ājôß]

ĴēÖĴİ¾Ź

0ŖğP {ĤĤÊ×j εÜeÖÊĔ
0ŖğP {ĤĤÊ×j εÜeÖÊĔ+ 3 ĞÜj¾+ 2 ēÜi0+ ¾ıŁŁP-εP-ŁÖ-ŁÓ- εÜ- ĞÜiÊēĀ+
8+ ¾ĤēŃēĀßÖ 0ĤÊ+ 0ēĀÜ0 ĩ 560052 0ŖğP {ĤĤÊ×j εÜeÖÊĔ+ 156/157+
Ńjô0ÖŃ 0jÜŃ+ 15 ēÜi0+ 227 Ńj0ğÖŃ 0j{ēÊ+ ÖĖ0| ĩ 400021 0ēĀÜ0
560052

0ò0P0 ¾ı 0 Ījğ

- (1). ????? ??????? ??????, ????? ?? 3301, ??,??,??,??.
????????????? ?????, ????? ?????, ?????????? 393002, ??????
- (2). ????? ????????? ???????, ??-5/??-6, ??,??,??,??, ???-3, ?????,
????? - 444104, ????????????
- (3). ????? ????????? ???????, ????? ??, ??,44, ?????? ???, ?????,
??????-?????, ????? ????? - 392130, ??????
- (4). ??? ?????? ??????? ???????, ????? ??, ??? - ???, ??? ??.?.???
????? ?, ??,?. ?????, ?? ĩ ???????, ????? ????? ĩ ??? ???, ??????

Öjß¾ ĪŁÖjÜ 0ÜÜ0İ

-

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

| ÎîŔĆƮÙÈÊ ÊƧŮÊÛ

(1). M/s Rallis India Limited, Plot No. 3301, GIDC Industrial Area,
Dist. Bharuch, Ankleshwar-393 002, GUJARAT
Gujarat

(2). M/s Rallis India Limited, C-5/C-6, MIDC Industrial Area,Phase-
III, Shivani, Akola i 444 104, MAHARASTHRA
Maharashtra

(3). Rallis India Ltd Plot No. C44, Chemical Zone, Dahej Taluka-
Vagra, District- Bharuch - 392130, GUJARAT

(4). M/s Rallis India Limited, Plot No. Z-110, Dahej SEZ Part II, P.O.
Lakhigam, Tal i Vagra, Dist : Bharuch 392 130, GUJARAT

(1). ????? ??????? ???????, ????? ?? 3301, ?? ?? ?? ??
????????????? ?????, ??? ???? , ????????? 393002, ?????

(2). ????? ??????? ???????, ??-5/??-6, ?? ?? ?? ?? ??-3, ?????,
???? - 444104, ?????????

(3). ????? ??????? ???????, ????? ?? ??44, ?????? ??, ???,
?????-????, ??? ???? - 392130, ?????

(4). ??? ????? ??????? ???????, ????? ?? ?? - ??, ??? ???.??
???? ? , ???. ?????, ?? i ?????, ??? ???? i ?? ??, ?????

Customer Care Details

Öİß¼ İkÖİÜ 0000 İ

-

-

Safety Pictograms:



- 1. not to be used for any purpose other than subscribed on label & leaflets
- 2. destroy container after use as directed on leaflet
- 3.Toxic to fish and other aquatic animals. Therefore, should not be used near aquaculture.
- 4. Toxic to bees. Therefore, should not be used during active foraging of bees.

Manufactured by

Rallis India Limited
Rallis India Limited, Illrd Block, IInd Floor, KSCMF Ltd., Building, 8, Cunningham Road, Bangalore - 560052 Rallis India Limited
156/157, Nariman Bhavan, 15th Floor, 228, Nariman Point, Mumbai - 400021 Bangalore Bangalore Karnataka 560052

Premises Address

- (1). M/s Rallis India Limited, Plot No. 3301, GIDC Industrial Area, Dist. Bharuch, Ankleshwar-393 002, GUJARAT Gujarat
- (2). M/s Rallis India Limited, C-5/C-6, MIDC Industrial Area,Phase-III, Shivani, Akola ĩ 444 104, MAHARASTHRA Maharashtra
- (3). Rallis India Ltd Plot No. C44, Chemical Zone, Dahej Taluka-Vagra, District- Bharuch - 392130, GUJARAT
- (4). M/s Rallis India Limited, Plot No. Z-110, Dahej SEZ Part II, P.O. Lakhigam, Tal ĩ Vagra, Dist : Bharuch 392 130, GUJARAT

Customer Care Details

-

- ÙÏÙ ©Ø ÒÃ    Āi yēŌĬā ßĴ }þþ[ēŌēÑ yē× Ŧ×iĀÑ Ō[}Ō×iĀ ¼[ēÙ  Ĩßġ]
a- ÒÃ    ĩÑôĬĥĒ }gŪ¼ ¼[yĨþĵØ Ŧĵ×iĀ ¼[ŌĵĬ ¼ĀĥÑØ 'iĒĜŌ[¼Āi ĨĥĒ ¼ĀØ ĬŦ]
3-ŌĀŪġ©Ø yē×ĀŪġ×ĀġŪĬ¼ĤŪ  ŪŸĴĈ Ĭ] { þēŪ +
 ĈŪĵ¼ġĀØ¼Ĥĵþ { Ĥ Ĭ ŤĵŪĨßġ  ×ĵĀĴĨĵĀĵ ß ]
4-Ō   ĈĈ   ×Ĭ¼ĤŪ  ŪŸĴĈ Ĭ] { þēŪ +
Ō    ĈĈ   ×Ĭ¼ ĤŪŌ×ŌĥĀĀ y Ū   ¼[Ĭ ŪĵĨ } Ō×iĀÑßġ  ×ĵĀĴĨĵĀĵ ß 
}  Ĭ Ĭ¼ 

   þ {ĥĒ×ĵ ēŪēŌĒĒ
   þ {ĥĒ×ĵ ēŪēŌĒĒ+ 3 ĈŪĵ¼+ 2     + ¼[  þ-ēþ- Ō- Ō- ēŪ- ĈŪĒēĀ+
8+ ¼ĥēĨēĀßŌ ŌiĒ+ Ō ĀŪ  Ĭ 560052    þ {ĥĒ×ĵ ēŪēŌĒĒ+ 156/157+
Ĩĵ  ŌÑ ŌĵŪĨ+ 15     + 227 Ĩĵ  ŌÑ Ōĵ{ēĒ+ Ō  Ĭ Ĭ 400021 Ō ĀŪ  560052

-       ¼[Ō Ĭĵ
- (1). ????? ??????? ???????, ????? ?? 3301, ??,??,??,??, ????????????? ?????, ????? ?????, ?????????? 393002, ??????
 - (2). ????? ????????? ???????, ??-5/??-6, ??,??,??,??, ???-3, ?????, ????? - 444104, ??????????
 - (3). ????? ????????? ???????, ????? ??, ??,44, ?????? ???, ???, ??????-????, ????? ??? - 392130, ??????
 - (4). ??? ????? ??????? ???????, ????? ??, ??? - ???, ??? ??.?.?? ????? ? , ??,?. ?????, ?? Ĭ ?????, ??? ????? Ĭ ??? ???, ??????

ŌĴ¼ ĬĤ ŌĵŪ      Ĭ

-

Safety Pictograms:



ÖĆ¼ȳ	ÈØεƁ¼Ö ò ì xyÀÖjØŷ ©Ø AȢ z	ad_	d__	d__	ÒÛxÔÑĦÒŕÍ Ɓ ȚĂÑÑ̃ yÔĤÎȳ í¼]	Ț×iÀ ¼ ac ÁĤ ¼ Ôȳİ ɔß ȚÛÛ ¼Øſ]	ad	øiÀ ¼ İÔȳÔ ¼ z ƏȳØ ÒØ a,b Țŕ í řĂÊ¼ȳÔ ¼Øſ]
ÖȳĂε	Ăũ xyεƁÍȳ+ ÔÛ İÔÛĂÑ 'αŞĆÑiƁ(+ Êȳ{ Ôſ¼ ©Ø œØÛÀ ĞÛȳ È	ad_	d__	d__	ÒÛxÔÑĦÒŕÍ Ɓ ȚĂÑÑ̃ yÔĤÎȳ í¼]	Ț×iÀ ¼ ac ÁĤ ¼ Ôȳİ ɔß ȚÛÛ ¼Øſ]	b	øiÀ ¼ İÔȳÔ ¼ z ƏȳØ ÒØ a,b Țŕ í řĂÊ¼ȳÔ ¼Øſ]
Ɓi×İÖÑ	ÀȢ z	ad_	d__	d__	ÒÛxÔÑĦÒŕÍ Ɓ ȚĂÑÑ̃ yÔĤÎȳ í¼]	Ț×iÀ ¼ ac ÁĤ ¼ Ôȳİ ɔß ȚÛÛ ¼Øſ]	cb	øiÀ ¼ İÔȳÔ ¼ z ƏȳØ ÒØ a,b Țŕ í řĂÊ¼ȳÔ ¼Øſ]

Note:-

Kùl·nùŘK ĦnŰŲk NĶ'nùKzB̃ſ̃b̃N Ħ ā ŸĦ ı ũśȳ

ȳÔ×iÀ ¼|εÛα řÑİŰ

Ôİȳİx¼ȳ řÑĐȳØÍ Öȳś ¼ȳ ȳÔ×iÀ ¼Øſ] Ôİȳİx¼i řÑĐȳØÍ ÔȳŊŷ ¼ȳ Öȳś ÖſȳćĂε Ț¼ȳØ Ɓ|εÔÛȳα]

ÖlƁÖ ¼ȳ ħĤİŕÍ

ÔØƁȳÍ+ ƁÛȳ ©Ø ÀÖxĐÛ yÔkĐ ¼|İœȳŊ řĂÊ¼ȳÔ Ŋ ¼Øſſ

εÖœÊŷ ©Ø ÔȳŊŷ

×ß ÔİȳİxÔăŷ řĂÊ¼ȳÔ ¼|εÛα ȳÔ×iÀ ȳ¼×ȳ ĂȳÍȳ ƁȢ {ƁεÛα εÔœɔÈ Ɓ| {Ɓ¼ȳ í¼ØŷÔÑ ¼i| ɔØĤÍȳ Ŋɔß ƁȢ} ĂεÛ× ¼œŰ Ö| ©Ø {Ɓ¼| z Ɓ ÔȳƁ
ȳÔ×iÀ Ŋ ¼Øſ]

ÓƁÛ yÔĤÎȳ

ÒÛxÔÑĦÒŕÍ Ɓ|ȚĂÑÑ̃ yÔĤÎȳ í¼]

z ÛİİŊ í¼Ŋŷȳ¼

ȳćĂ Ûȳġ×Đ ÛȳÛ|ĤȚ×Ɓx¼| ĂôØα Ôăŷ řĂÊ¼ȳÔ ¼Øſ]

ĬȳKđń TKũđȳ

Use compression Knapsack sprayer, Foot sprayer, High & low volume sprayer or Battery operated sprayers.

ŋźLœ TKũđȳ

ƁiÛi ¼iŊ ŊiĂÛ

kđŰKȳ

Please refer recommendation table.

ŰöYń KŸœĬ ŰŋŊăȳ

ÓÓũŊȳÛȳ¼ ¼ȳ řÑĐȳØÍ Öȳśȳ ¼i ÒßÛ| İİİ|ÔȳŊŷ Ö|ȳćĂε Ț¼ȳØ εÔÛȳ×| ȳƁ¼|Ôȳİ ÔȳŊŷ ¼ȳ z ÔĤ×ȳ¼ Öȳśȳ Ăi řÑĐȳØÍ ¼ȳ Ă×ŷ Ɓi εÔÛȳ×| ©Ø
Ăiŷ ȚİĂİcȚȳćĂε Ț¼ȳØ εÔÛȳ×İ
ŋŰŰ ȚİĂİcȚȳćĂε

Ț×iÀ ¼|ac ÁĤ| ¼|Ôȳİ ɔß ȚÛÛ ¼Øſ]

¼Êȳ| Ɓ|ÔßÛ| ¼ȳ yŰŵȳÛ

b_

ĥ rŊŋ Kũ śđŲ ĩȳ đ śđŲȳ

øiÀ ¼|ÛĐ z Íŷ Ûà ì ŊéØ z Ŋİ|ÔØ Ôăŷ řĂÊ¼ȳÔ ¼Øſ]

z ÛİİŊ ¼ȳ ƁÖ×

Caution:ñNot to Be Used on Crops Other Than Specified on This Label/Leafletò.

Kresoxim-methyl 44.3 % w/w (500 g/l) SC

(Systemic Fungicide)

Kresoxim-methyl 44.3% w/w (500 g/L) SC is a broad spectrum systemic fungicide used for the control of Blast & Sheath blight on Rice; Powdery mildew & Downy mildew on Grapes; Powdery mildew, Fruit rot, Die back & Twig blight on Chillies; Rust on Soybean; Late blight & Early blight on Potato; Leaf spot & Grey mildew on Cotton; Rust & Leaf blight on Wheat and Turcicum leaf blight & Rust on Maize.

Chemical Composition:	Ingredient	%W/W
Blend of Methyl isothiazolone and its chloro derivative	-	0.5 % w/w
Demineralized water	-	QS to make 100% -
Ethoxylated derivative of Styrenated Phenol	-	1.5 % w/w
Ethoxylated polymethacrylate in Propylene glycol and water	-	4.0 % w/w
Kresoxim-methyl a.i. content	-	44.3 % w/w
Polydimethyl siloxane (Defoamer)	-	1.0 % w/w
Rhodopol 23 as 5% Gel	-	2.5 % w/w
Sodium Bentonite	-	0.25 % w/w
TOTAL	-	100.0 % w/w
Trihydroxy Propane (Glycerol)	-	5.0 % w/w
Total:		100.00% w/w

Recommendation

<u>Crop(s)</u>	<u>Common Name of Pest</u>	<u>Dosage/HA</u>		<u>Dilution in Water Dilution in water (Litre)</u>	<u>Crop Stage</u>	<u>Re-entry Period</u>	<u>Pre Harvest</u>	<u>Frequency</u>
		<u>AI a.i. (g)</u>	<u>Formulation Formulation (mL)</u>					
Grapes	Powdery mildew Downy mildew	300-350	600-700	500	Early vegetative to reproductive stage	24 hours of spraying.	7	2-3 foliar spray depending on disease pressure
Potato	Late blight Early blight	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	23	2-3 foliar spray depending on disease pressure
Cotton	Leaf spot Grey mildew	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	26	2-3 foliar spray depending on disease pressure
Wheat	Rust Leaf blight	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	34	2-3 foliar spray depending on disease pressure
Rice	Blast Sheath blight	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	30	2-3 foliar spray depending on disease pressure
Maize	Turcicum leaf blight Rust	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	25	2-3 foliar spray depending on disease pressure
Chillies	Powdery mildew Fruit rot (Anthracnose) Die back Twig blight	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	3	2-3 foliar spray depending on disease pressure
Soybean	Rust	250	500	500	Early vegetative to reproductive stage	24 hours of spraying.	43	2-3 foliar spray depending on disease pressure

Note:-

Good Practices relating to pesticide application techniques

Caution:ñNot to Be Used on Crops Other Than Specified on This Label/Leaflet.

Direction of Use

Use recommended dose. Dilute the formulation in required quantity of water.

Weather Condition(in general):

Do not spray during rainy, windy and hot sunny period.

Soil & Water(when ever applicable):

This is foliar application combination product, so hardly have any relation to soil. Avoid using near aquaculture.

Crop Stage for application:

Early vegetative to reproductive stage

Application Technique

Foliar spray using high volume sprayers.

Equipment Type:

Use compression Knapsack sprayer, Foot sprayer, High & low volume sprayer or Battery operated sprayers.

NozzleType:

hollow cone nozzle

Dosage:

Please refer recommendation table.

Instruction for Mixing:

Take the required quantity of fungicide, dilute with a little water and stir using a stick. Then add the remaining quantity of water as per spray volume required and mix well again by stirring.

Re-entry Period

24 hours of spraying.

Pre-Harvesting Interval

30

Manner and Time of Application

Foliar spray. As soon as the disease appears.

Application Timing

Should be applied preferably in the morning or evening hours.

Frequency

2-3 foliar spray depending on disease pressure

Stages during which pesticides should not be applied on any crop :**Not to be applied during crop maturity and just before crop harvest.****Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :**

- 1.Avoid for fish and other aquatic animals. Therefore, should not go to experiment with aquaculture.
2. Toxic for honey bees. Therefore, should not be used during active foraging of bees.

Symptoms Of Poisoning

If ingested, the early symptoms may be a combination of nausea, vomiting, diarrhea, abdominal pain.

Caution:ñNot to Be Used on Crops Other Than Specified on This Label/Leaflet.

First Aid

1. If ingested : If person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious person. Call the doctor. Do not induce vomiting until and unless desired by attending physician. 2. If inhaled : Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, preferably with mouth to mouth breathing. Consult a physician. 3. In case of skin contact : Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All the contaminated clothes should be segregated and thoroughly washed with soap and water separately before use. 4. In case of eye contact : Rinse gently with plenty of water for at least 15 minutes. If pain, irritation, redness or photophobia persists, consult eye specialist.

Antidote

There is no specific antidote, hence symptomatic and supportive therapy should be given.

Phytotoxicity

Kresoxim-methyl 500 g/L SC is not phytotoxic when used as per recommendations.

Storage Conditions

1. The packages containing the fungicides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles particularly food articles or shall be kept in separate almirahs under lock and key depending on the quantity and the nature of the fungicides.
2. The store room should be well built, well lit, ventilated and sufficient in dimensions. The condition of the store should be dry and cool.

Safety Precaution:

Protective clothing should be worn while mixing and spraying. Wash contaminated clothes and body parts with water after application. Avoid inhalation, do not smoke, drink, eat & chew anything during application.

Cautionary Statement:

- 1. not to be used for any purpose other than subscribed on label & leaflets**
2. destroy container after use as directed on leaflet
3.Toxic to fish and other aquatic animals. Therefore, should not be used near aquaculture.
4. Toxic to bees. Therefore, should not be used during active foraging of bees.

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

M/s Rallis India Limited, C-5/C-6, MIDC Industrial Area,Phase-III,
Shivani, Akola ĩ 444 104, MAHARASTHRA
Karnataka

M/s Rallis India Limited, Plot No. 3301, GIDC Industrial Area, Dist.
Bharuch, Ankleshwar-393 002, GUJARAT
Karnataka

M/s Rallis India Limited, Plot No. Z-110, Dahej SEZ Part II, P.O.
Lakhigam, Tal ĩ Vagra, Dist : Bharuch 392 130, GUJARAT
Karnataka

Rallis India Ltd Plot No. C44, Chemical Zone, Dahej Taluka-
Vagra, District- Bharuch - 392130, GUJARAT
Karnataka

Manufactured by

Rallis India Limited
Rallis India Limited, IIIrd Block, IIInd Floor, KSCMF Ltd., Building, 8,
Cunningham Road, Bangalore - 560052 Rallis India Limited
156/157, Nariman Bhavan, 15th Floor, 228, Nariman Point,
Mumbai - 400021 Bangalore Karnataka 560052

Customer Care Details:

-

Safety Pictograms:



Caution:ñNot to Be Used on Crops Other Than Specified on This Label/Leafletò.