

Case of Initial Number of Susceptibles Slightly Higher than the Threshold

$$\boxed{\frac{dx}{dt} = x(Ay - B)} \text{ and } \boxed{x = (x_0 + y_0) - y + \frac{B}{A} \ln\left(\frac{y}{y_0}\right)}$$

When $\boxed{\frac{dx}{dt} = 0}$ Either $\boxed{y = B/A = p}$ or $\boxed{x = 0}$.

i) The former case is a turning point in which $\boxed{y = p}$ is a maximum $\Rightarrow \boxed{d^2x/dt^2 < 0}$.

ii) $\boxed{\frac{d^2x}{dt^2} = \frac{dx}{dt}(Ay - B) + x \frac{d}{dt}(Ay - B)}$. When $\boxed{x = 0}$

at $\boxed{\frac{dx}{dt} = 0} \Rightarrow \boxed{\frac{d^2x}{dt^2} = 0}$. This is an equilibrium point. (at $x = 0$).

When $\boxed{x = 0}$, we write $\boxed{y = y_\infty}$. With these,

We get $\boxed{0 = x_0 + y_0 - y_\infty + \frac{B}{A} \ln\left(\frac{y_\infty}{y_0}\right)}$. Usually

$\boxed{x_0 \ll y_0}$, and so we neglect x_0 above.

Now y_0 is the initial value of the Susceptibles, and y_∞ is the final value of the Susceptibles.

$\therefore \boxed{y_0 - y_\infty}$ is the number of people who have Contracted the infectious Disease. Further,

$\boxed{y_\infty < y_0}$, which implies that $\boxed{y_0 - y_\infty > 0}$.

Hence we can write $\boxed{(y_0 - y_\infty) + \frac{B}{A} \ln\left(\frac{y_0 - y_0 + y_\infty}{y_0}\right) \approx 0}$

$\Rightarrow \boxed{(y_0 - y_\infty) + p \ln\left(1 - \frac{y_0 - y_\infty}{y_0}\right) \approx 0}$ in which we have neglected x_0 .
(P.T.O.)

(Continued) - 23 -

We now consider the case where the initial number of susceptibles is slightly greater than the threshold, i.e. $y_0 = p + \epsilon$, in which $\epsilon \ll p$. $\therefore \frac{(y_0 - p)}{p} = \frac{\epsilon}{p} \ll 1$.

From the plot of $x-y$ we can see that in this case $(y_0 - y_\infty) \ll y_0$. Hence

using the formula $\ln(1+u) = u - \frac{u^2}{2} + \frac{u^3}{3} - \dots$,

we can write $\ln\left[1 - \left(\frac{y_0 - y_\infty}{y_0}\right)\right] = -\left(\frac{y_0 - y_\infty}{y_0}\right) - \frac{1}{2}\left(\frac{y_0 - y_\infty}{y_0}\right)^2$

going only up to the second order term.

Hence, $(y_0 - y_\infty) + p \left[-\left(\frac{y_0 - y_\infty}{y_0}\right) - \frac{1}{2}\left(\frac{y_0 - y_\infty}{y_0}\right)^2 \right] \approx 0$

$\Rightarrow (y_0 - y_\infty) \left[1 - \frac{p}{y_0} - \frac{p}{2} \frac{(y_0 - y_\infty)}{y_0^2} \right] \approx 0$ (Now, $y_0 - y_\infty \neq 0$)

$\Rightarrow y_0 - y_\infty \approx \frac{2y_0^2}{p} \left(1 - \frac{p}{y_0} \right) \approx 2y_0 \left(\frac{y_0}{p} - 1 \right)$

But $y_0 = p + \epsilon$ and $\left(\frac{y_0}{p} - 1 \right) = \frac{\epsilon}{p}$. Using these $\epsilon \ll p$

we get, $y_0 - y_\infty \approx 2y_0 \epsilon / p \approx 2(p + \epsilon) \epsilon / p$

Neglecting ϵ in $p + \epsilon \approx p$, we finally get.

$y_0 - y_\infty \approx 2\epsilon$ $\Rightarrow y_0 - y_\infty = 2\epsilon$.

$\therefore y_0 - y_\infty \approx 2(y_0 - p)$, an approximate result that is valid only when y_0 is slightly greater than p .

- 24 -

$Z(t)$ as a function of Time : $Z \equiv Z(t)$.

$$\boxed{\frac{dz}{dt} = Bx} , \quad \boxed{x = (x_0 + y_0) - y + \frac{B}{A} \ln\left(\frac{y}{y_0}\right)} \text{ and}$$

$$\boxed{y = y_0 \exp\left(-\frac{AZ}{B}\right)} \Rightarrow \boxed{Z = -\frac{B}{A} \ln\left(y/y_0\right)}.$$

Hence, $\boxed{\frac{dz}{dt} = B \left[x_0 + y_0 - y_0 \exp\left(-\frac{AZ}{B}\right) - Z \right]}$,

an equation in an autonomous form $\boxed{\frac{dz}{dt} = f(z)}$.

Noting $\boxed{\frac{B}{A} = p}$, we consider the case of $\boxed{Z \ll p}$.

$$\therefore \boxed{e^{-Z/p} \approx 1 - \frac{Z}{p} + \frac{1}{2} \left(\frac{Z}{p}\right)^2 + \dots}$$
, bring up to the second order.

$$\Rightarrow \frac{dz}{dt} \approx B \left[x_0 + \cancel{y_0} - \cancel{y_0} + y_0 \frac{Z}{p} - \frac{y_0}{2} \left(\frac{Z}{p}\right)^2 - Z \right]$$

$$\Rightarrow \boxed{\frac{dz}{dt} = B \left[x_0 + \left(\frac{y_0}{p} - 1\right) Z - \frac{y_0}{2} \left(\frac{Z}{p}\right)^2 \right]}$$

\downarrow
 $\frac{1}{B} \frac{dz}{dt} = x$
 $\Rightarrow x \equiv x(Z)$

We define $\boxed{\beta = Z/p}$, by which we set,

$$\boxed{\frac{p}{B} \frac{d\beta}{dt} = x_0 + (y_0 - p)\beta - \frac{y_0}{2} \beta^2}$$

Multiplying through out by $\frac{2}{y_0}$ we get.

$$\boxed{\frac{2}{y_0} \frac{p}{B} \frac{d\beta}{dt} = \frac{2x_0}{y_0} + 2\left(1 - \frac{p}{y_0}\right)\beta - \beta^2}$$

(P.T.O.)

We further define $\boxed{\alpha^2 = 2x_0/y_0}$ and $\boxed{\beta^* = 1 - \frac{p}{y_0}}$.

Hence
$$\frac{2}{y_0} \frac{l}{B} \frac{d\beta}{dt} = -(\beta^2 - 2\beta\beta + \beta^2 - \beta^2) + \alpha^2$$

$$\Rightarrow \frac{2}{y_0 A} \frac{d\beta}{dt} = (\beta^2 + \alpha^2) - (\beta - \beta)^2 \quad \because \left[\frac{l}{B} = \frac{R}{A} \cdot \frac{1}{2} \right]$$

Further define $k^2 = \alpha^2 + \beta^2$ and $\xi = \beta - \beta$,

which gives $\frac{d\beta}{dt} = \frac{d\xi}{dt}$. Using this we get,

$$\frac{2}{A y_0} \frac{d\xi}{dt} = k^2 - \xi^2 \Rightarrow \frac{2}{A y_0 k} \frac{d(\xi/k)}{dt} = 1 - \left(\frac{\xi}{k}\right)^2$$

Again, defining $\psi = \xi/k$ and $\tau = \frac{A y_0 k}{2} t$,

we obtain finally, $\frac{d\psi}{d\tau} = 1 - \psi^2$. This

Equation can be integrated to obtain

$$\ln \left(\frac{1+\psi}{1-\psi} \right) = 2\tau + 2C$$
, where C is an integration constant.

(Check for the solution of $\frac{dx}{dT} = 1 - x^2$ equation)

Writing $T = \tau + C$, we get, $\frac{1+\psi}{1-\psi} = e^{2T}$, from

which we get $1+\psi = (1-\psi)e^{2T} \Rightarrow \psi(1+e^{2T}) = e^{2T} - 1$

$$\Rightarrow \psi = \frac{e^{2T} - 1}{e^{2T} + 1} \Rightarrow \psi = \frac{e^T - e^{-T}}{e^T + e^{-T}} = \tanh(T)$$

$$\therefore \frac{\xi}{k} = \tanh(\tau + C) \Rightarrow \frac{\beta - \beta}{k} = \tanh(\tau + C)$$

(continued) - 26 -

$$\Rightarrow \boxed{S = \beta + k \tanh(\tau + c)}$$

$$\Rightarrow \boxed{\frac{Z}{p} = \beta + k \tanh\left(\frac{A y_0 k}{2} t + c\right)}$$

$$\Rightarrow \boxed{Z = \beta p + k p \tanh\left(\frac{A y_0 k}{2} t + c\right)} \quad \boxed{Z \equiv Z(t)}$$

Hence, $\boxed{\frac{dZ}{dt} = k p \cdot \left(\frac{A y_0 k}{2}\right) \operatorname{sech}^2\left(\frac{A y_0 k}{2} t + c\right)}$


$$\Rightarrow \boxed{\frac{dZ}{dt} = \frac{B y_0 k^2}{2} \operatorname{sech}^2\left(\frac{A y_0 k}{2} t + c\right)} \quad \left\{ \begin{array}{l} \frac{1}{\beta} \frac{dZ}{dt} = \kappa \\ \Rightarrow \kappa \equiv \kappa(t) \end{array} \right.$$

($\varphi = B/A$)

The integration constant c can be fixed from the initial condition, $\boxed{Z=0}$ at $\boxed{t=0}$.

$$\Rightarrow \boxed{0 = \beta p + k p \tanh(c)} \Rightarrow \boxed{\tanh(c) = -\frac{\beta}{k}}$$

Now $\boxed{\beta = 1 - \frac{p}{y_0}} \therefore \boxed{\tanh(c) = \frac{1}{k} \left(\frac{p}{y_0} - 1\right)}$.



i) The epidemic curve reaches a maximum when $\boxed{\frac{A y_0 k}{2} t + c = 0}$.

$$\Rightarrow \boxed{t = -2c/Aky_0} \quad (c < 0)$$

ii) The removal rate $\boxed{dZ/dt}$ climbs to a peak value and dies out.

iii) Data matched by the Bombay plague of 1905.