

# ENGG2440 Discrete mathematics for engineers

Ryan Chan

November 30, 2024

### **Abstract**

This is a note for **ENGG2440 - Discrete mathematics for engineers** for self-revision purpose ONLY.  
Some contents are taken from lecture notes and reference book.  
Mistakes might be found. So please feel free to point out any mistakes.

# Contents

<b>1</b>	<b>Mathematical Induction</b>	<b>2</b>
1.1	Introduction . . . . .	2
1.2	Mathematical Induction . . . . .	2
1.3	Strong Mathematical Induction . . . . .	3
<b>2</b>	<b>Summation Techniques</b>	<b>5</b>
2.1	Summation . . . . .	5
2.2	Close Form Formula . . . . .	6
2.3	Perturbation Method . . . . .	6
2.4	Guess-and-Verify method . . . . .	8
2.5	Multiple Summation . . . . .	9
<b>3</b>	<b>Recurrences</b>	<b>12</b>
3.1	Homogeneous Recurrences . . . . .	12
3.2	Non-homogeneous Recurrences . . . . .	13
<b>4</b>	<b>Asymptotics</b>	<b>16</b>
4.1	Big O . . . . .	16
4.2	Big Omega . . . . .	17
4.3	Theta . . . . .	18
4.4	Little O . . . . .	18
4.5	Little Omega . . . . .	19
4.6	Properties for Asymptotic Analysis . . . . .	19
<b>5</b>	<b>Set Theory and Counting Principle</b>	<b>20</b>
5.1	Set Theory . . . . .	20
5.2	Counting Principle . . . . .	20
5.3	Permutation and Combination . . . . .	23
<b>6</b>	<b>Binomial Coefficients</b>	<b>24</b>
6.1	Introduction . . . . .	24
<b>7</b>	<b>Elements of Discrete Probability</b>	<b>26</b>
7.1	. . . . .	26

# Chapter 1

## Mathematical Induction

### 1.1 Introduction

In mathematics, there are some basic proof techniques that we can apply, including direct proof, proof by induction, proof by contradiction, and proof by contraposition. For most of these proving methods, you won't be learning their reasons or applications, but you will still use them in some simple proving questions. In this chapter, we will mainly discuss mathematical induction.

**Definition 1.1.1 (Proposition).** A **proposition** is a statement that is either true or false.

**Definition 1.1.2 (Predicate).** A **predicate** is a proposition whose truth depends on one or more variables.

### 1.2 Mathematical Induction

An analogy of the principle of mathematical induction is the game of dominoes. Suppose the dominoes are lined up properly, so that when one falls, the successive one will also fall. Now by pushing the first domino, the second will fall; when the second falls, the third will fall; and so on. We can see that all dominoes will ultimately fall.

The key point is only two steps:

1. the first domino falls;
2. when a domino falls, the next domino falls.

We use the above principle of Mathematical Induction to prove.

Process:

1. Let  $P(n)$  be a predicate.
2. (Base Case) Show that  $P(1)$  is true.
3. (Inductive Steps) Show that for  $n = 1, 2, \dots$ , if  $P(n)$  is true, then  $P(n + 1)$  is true.

**Example.**

$$P(n) : 1 + 2 + \dots + n = \frac{n(n+1)}{2}$$

1. Base Case

We need to show that  $P(1)$  is true.

$$1 = \frac{(1)(1+1)}{2},$$

which is obviously true.

## 2. Inductive Step

For inductive hypothesis, we can assume

$$1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

Now, to show that  $P(n+1)$  is true,

$$\begin{aligned} L.H.S. &= 1 + 2 + \cdots + n + (n+1) \\ &= \frac{n(n+1)}{2} + (n+1) \\ &= \frac{n(n+1) + 2(n+1)}{2} \quad , \\ &= \frac{(n+1)(n+2)}{2} \\ &= R.H.S \end{aligned}$$

which shows that  $P(n+1)$  is also true.

Hence, by the principle of MI, we can conclude that  $P(n)$  is true for all integers  $n \geq 1$ .

**Exercise.** Show that for any integer  $n \geq 1$ ,  $n^3 - n$  is divisible by 3.

**Note.** In inductive step, consider putting a constant  $q$  as  $3q$  is divisible by 3.

**Exercise.** Prove that  $n^3 < 2^n$  for all integers  $n \geq 10$ .

**Note.** Consider bonding the lower order terms in terms of  $n^3$ .

## 1.3 Strong Mathematical Induction

As the name suggests, the method of induction used in this section is "stronger". This is because assuming only that  $P(n)$  is true may be too restrictive, i.e., insufficient to prove the predicate. Thus, in the inductive step, you may show that  $P(1), P(2), \dots, P(n)$  are true, and then prove that  $P(n+1)$  is true.

**Example.** The Fibonacci sequence is a sequence of number defined via the following recursion:

$$\begin{aligned} F_n &= F_{n-1} + F_{n-2}, \quad n \geq 2 \\ F_0 &= 0; F_1 = 1 \end{aligned}$$

Prove that

$$P(n) : F_n \leq \phi^{n-1}, \text{ where } \phi = \frac{1 + \sqrt{5}}{2}$$

### 1. Base Case

$$\begin{aligned} F_1 &= 1 \leq \phi^0 = 1 \\ F_2 &= 1 \leq \phi^1 \approx 1.618 \end{aligned}$$

Thus,  $P(1)$  and  $P(2)$  hold true, which means  $P(3)$  also holds true.

### 2. Inductive Step

---

For inductive hypothesis, we assume

$$F_k \leq \phi^{k-1} \text{ for } k = 1, 2, \dots, n$$

Given the Fibonacci sequence

$$F_{n+1} = F_n + F_{n-1}$$

By the strong inductive hypothesis, we have

$$F_n \leq \phi^{n-1}, \quad F_{n-1} \leq \phi^{n-2}$$

From the definition of  $\phi$  we have  $\phi^2 = 1 + \phi$

Hence, we obtain

$$F_{n+1} \leq \phi^{n-1} + \phi^{n-2} = \phi^{n-2}(1 + \phi) = \phi^n$$

## Chapter 2

# Summation Techniques

### 2.1 Summation

When summing numbers with certain patterns, we can use summation notation. For example,

$$a_1 + a_2 + \cdots + a_n = \sum_{k=1}^n a_k$$

#### 2.1.1 Distributive Law

Let  $c$  be a constant. Then, we can take  $c$  out of the summation:

$$\sum_{k \in \mathcal{K}} ca_k = c \sum_{k \in \mathcal{K}} a_k$$

**Example.**

$$\sum_{k=1}^n 2k = 2(1) + 2(2) + 2(3) + \cdots + 2(n) = 2(1 + 2 + 3 + \cdots + n) = 2 \sum_{k=1}^n k$$

#### 2.1.2 Associative Law

We can split the summand as follows:

$$\sum_{k \in \mathcal{K}} (a_k + b_k) = \sum_{k \in \mathcal{K}} a_k + \sum_{k \in \mathcal{K}} b_k$$

**Example.**

$$\begin{aligned} \sum_{k=1}^n (k + k^2) &= (1 + 1^2) + (2 + 2^2) + \cdots + (n + n^2) \\ &= (1 + 2 + \cdots + n) + (1^2 + 2^2 + \cdots + n^2) \\ &= \sum_{k=1}^n k + \sum_{k=1}^n k^2 \end{aligned}$$

---

## 2.2 Close Form Formula

Close form formula is the formula that does not have the summation index  $k$  for a sum by simply writing it out explicitly. For example,

$$\sum_{k=1}^n (a_k - a_{k-1})$$

By expanding the sum, we have

$$\sum_{k=1}^n (a_k - a_{k-1}) = (a_1 - a_0) + (a_2 - a_1) + \cdots + (a_n - a_{n-1}) = a_n - a_0$$

By cancelling the terms, we get  $a_n - a_0$ , which is the close form formula for the summation  $\sum_{k=1}^n (a_k - a_{k-1})$ .

## 2.3 Perturbation Method

It could be difficult to derive the close form formula for some summation. Therefore, we can use the perturbation method.

For summation

$$S_n = \sum_{k=1}^n a_k,$$

we can split off the first term and the last term, then rewrite it as

$$a_1 + \sum_{k=2}^{n+1} a_k = S_{n+1} = \sum_{k=1}^n a_k + a_{n+1}$$

**Example (Geometric Sum).** Let  $x$  be any number. Consider the sum

$$S_n = \sum_{k=1}^n x^k$$

$$x + \sum_{k=2}^{n+1} x^k = S_{n+1} = \sum_{k=1}^n x^k + x^{n+1}$$

Observe that

$$\sum_{k=2}^{n+1} x^k = x^2 + x^3 + \cdots + x^{n+1} = x(x + x^2 + \cdots + x^n) = xS_n$$

By substitution, we have

$$x + xS_n = S_n + x^{n+1}$$

If  $x \neq 1$ , then we can solve for  $S_n$  to get

$$S_n = \frac{x(1 - x^n)}{1 - x}$$

This summation is also called geometric sum.

By applying the perturbation method, we can find the close form formula for some common summation. Another example is Quadratic Series.



**Example (Quadratic Series).** By applying perturbation method to the sum

$$S_n = \sum_{k=1}^n k^2,$$

we have

$$1 + \sum_{k=2}^{n+1} k^2 = S_{n+1} = \sum_{k=1}^n k^2 + (n+1)^2$$

Let  $j = k - 1$ ,

$$\begin{aligned} \sum_{k=2}^{n+1} k^2 &= \sum_{j=1}^n (j+1)^2 \\ \sum_{j=1}^n (j+1)^2 &= \sum_{j=1}^n (j^2 + 2j + 1) \\ &= \sum_{j=1}^n j^2 + 2 \sum_{j=1}^n j + \sum_{j=1}^n 1 \\ &= S_n + 2 \sum_{j=1}^n j + n \end{aligned}$$

Then, we have

$$\begin{aligned} 1 + \sum_{k=2}^{n+1} k^2 &= \sum_{k=1}^n k^2 + (n+1)^2 \\ 1 + S_n + 2 \sum_{j=1}^n j + n &= S_n + (n+1)^2 \\ \sum_{j=1}^n j &= \frac{n(n+1)}{2} \end{aligned}$$

However, we obtain the Euler's trick here instead. Thus, we may apply the perturbation method to another sum.

$$\begin{aligned} C_n = \sum_{k=1}^n k^3 &\Rightarrow 1 + \sum_{k=2}^{n+1} k^3 = C_{n+1} = \sum_{k=1}^n k^3 + (n+1)^3 \\ \sum_{k=2}^{n+1} k^3 &= \sum_{j=1}^n (j+1)^3 \quad (\text{By applying } j = k - 1) \\ &= \sum_{j=1}^n j^3 + 3 \sum_{j=1}^n j^2 + 3 \sum_{j=1}^n j + \sum_{j=1}^n 1 \\ &= C_n + 3S_n + \frac{3n(n+1)}{2} + n \end{aligned}$$

By substitution, we get

$$\begin{aligned} 1 + C_n + 3S_n + \frac{3n(n+1)}{2} + n &= C_n + (n+1)^3 \\ 2 + 6S_n + 3n(n+1) + 2n &= 2(n+1)^3 \\ S_n &= \frac{2(n+1)^3 - 2n - 2 - 3n(n+1)}{6} \\ S_n &= \frac{(n+1)(2(n+1)^2 - 2 - 3n)}{6} \\ S_n &= \frac{n(n+1)(2n+1)}{6} \end{aligned}$$

There are some shortcut expression that might be used without finding the closed form formula on your own.

**Proposition 2.3.1** (Close form formula). (You can try the perturbation method to find the close form formula by yourself.)

- Geometric series

$$\sum_{k=0}^n ar^k = \frac{a(r^{n+1} - 1)}{r - 1}, r \neq 1$$

- Euler's trick

$$\sum_{k=1}^n k = \frac{n(n+1)}{2} \text{ or } \sum_{k=a}^b k = \frac{(b-a+1)(a+b)}{2}$$

- Quadratic series

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

- Cubic series

$$\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4}$$

## 2.4 Guess-and-Verify method

As the name suggests, we can often guess the closed-form formula. But how can we be certain it's correct? This is where mathematical induction comes in handy.

**Example.**

$$S_n = \sum_{k=1}^n k^2$$

Observe that  $S_n$  behaves like the sum of terms in a polynomial, allowing us to form an  $n$ -term polynomial. Since  $n^2$  is the largest term in  $S_n$ , we conclude that  $S_n \leq n^3$

$$S_n = a + bn + cn^2 + dn^3$$

$$1 = S_1 = a + b + c + d,$$

$$5 = S_2 = a + 2b + 4c + 8d,$$

$$14 = S_3 = a + 3b + 9c + 27d,$$

$$30 = S_4 = a + 4b + 16c + 64d$$

Solving for above, we have  $a = 0$ ,  $b = \frac{1}{6}$ ,  $c = \frac{1}{2}$ ,  $d = \frac{1}{3}$ .

$$S_n = \frac{1}{6}n + \frac{1}{2}n^2 + \frac{1}{3}n^3$$

After hypothesizing the closed-form formula, we need to use induction to verify its correctness. The steps are straightforward, and you can try proving it yourself. This process confirms that the formula is indeed the closed form for the summation.

## 2.5 Multiple Summation

All the summations above use only a single index. For example,

$$S_n = \sum_{k=1}^n k^2 = 1^2 + 2^2 + \cdots + n^2.$$

In this section, however, we introduce a summation with multiple indices. You can think of a single-index summation as summing over a 1D array. Extending this idea, a summation with two indices corresponds to summing over a 2D array. For example,

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

To sum up all the terms, we can use

$$r_j = a_{j1} + a_{j2} + \cdots + a_{jn} = \sum_{k=1}^n a_{jk}$$

Then, we can rewrite it as

$$S = \sum_{j=1}^m r_j = \sum_{j=1}^m \sum_{k=1}^n a_{jk}$$

We can also interchange the order of summation, which can be very useful for finding the closed-form formula. For the above summation, it is rather simple, we can simply do the interchange by

$$\sum_{j=1}^m \sum_{k=1}^n a_{jk} = \sum_{k=1}^n \sum_{j=1}^m a_{jk}$$

However, this does not work for all the summation.

**Example.** Considering

$$S = \sum_{j=1}^n \sum_{k=j}^n a_{jk}$$

To visualize that, we can again use matrix

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ & a_{22} & \cdots & a_{2n} \\ & & \ddots & \vdots \\ & & & a_{nn} \end{bmatrix}$$

Then, we have

$$\begin{aligned} S = & a_{11} + a_{12} + a_{13} + \cdots + a_{1n} \\ & + a_{22} + a_{23} + \cdots + a_{2n} \\ & \quad \quad \quad \ddots \\ & \quad \quad \quad + a_{nn} \end{aligned}$$

Let

$$c_k = a_{1k} + a_{2k} + \cdots + a_{nk} = \sum_{j=1}^n a_{jk},$$

we get

$$S = \sum_{k=1}^n c_k = \sum_{k=1}^n \sum_{j=1}^k a_{jk}$$

**Remark.** Informally, form

$$S = \sum_{j=1}^n \sum_{k=j}^n a_{jk}$$

we have  $1 \leq j \leq k \leq n$ , then we can simply interchange the order by

$$S = \sum_{k=1}^n \sum_{j=1}^k a_{jk}$$

Let's see another example,  $n$ -th harmonic number ( $H_n$ ).

**Example.**

$$H_n = \sum_{k=1}^n \frac{1}{k}$$

Again we can visualize it by using matrix

$$\begin{bmatrix} 1 & & & & \\ 1 & 1/2 & & & \\ 1 & 1/2 & 1/3 & & \\ & & & \ddots & \\ 1 & 1/2 & 1/3 & \cdots & 1/n \end{bmatrix}$$

$$S = \sum_{j=1}^n H_j = \sum_{j=1}^n \sum_{k=1}^j \frac{1}{k}$$

Since  $1 \leq k \leq j \leq n$

$$S = \sum_{j=1}^n H_j = \sum_{j=1}^n \sum_{k=1}^j \frac{1}{k} = \sum_{k=1}^n \sum_{j=k}^n \frac{1}{k}$$

Then we get

$$\begin{aligned} S &= \sum_{k=1}^n \sum_{j=k}^n \frac{1}{k} \\ &= \sum_{k=1}^n \frac{n - k + 1}{k} \\ &= \sum_{k=1}^n \frac{n}{k} - \sum_{k=1}^n 1 + \sum_{k=1}^n \frac{1}{k} \\ &= (n+1)H_n - n \end{aligned}$$

**Definition 2.5.1 (Floor function).** Given a real number  $x$ , define

$$\lfloor x \rfloor = \text{greatest integer} \leq x$$

The symbol  $\lfloor x \rfloor$  is usually read as "the floor of  $x$ ".

**Exercise.** For any real number  $x$ ,  $\lfloor x \rfloor \leq x \leq \lfloor x \rfloor + 1$ .

Find the closed form formula for

$$S_n = \sum_{k=1}^n \lfloor \sqrt{k} \rfloor$$

where, for simplicity, we assume that  $n$  is a perfect square, i.e.,  $n = a^2$  for some integer  $a \geq 1$ .

**Remark.** Consider  $\lfloor x \rfloor = \sum_{k=1}^x 1$ , then how to do the order interchange such that we can get the following summation

$$S_n = \sum_{j=1}^a \sum_{k=j^2}^{a^2} 1$$

Let's look at the last example.

**Example.** Let  $n \geq 1$  be an integer and  $x \neq 1$  be a real number.

$$S = \sum_{j=1}^n \sum_{k=1}^j kx^j$$

By interchanging the summation orders yields

$$S = \sum_{k=1}^n \sum_{j=k}^n kx^j$$

Notice that for any integer  $k$  satisfying  $1 \leq k \leq n$  it holds that

$$\sum_{j=k}^n x^j = \sum_{j=1}^n x^j - \sum_{j=1}^{k-1} x^j = \frac{x - x^{n+1}}{1 - x} - \frac{x - x^k}{1 - x} = \frac{x^k - x^{n+1}}{1 - x}$$

Equivalently,

$$\sum_{j=k}^n x^j = \frac{x^k - x^{n+1}}{1 - x} = x^{k-1} \cdot \frac{x - x^{n-k+2}}{1 - x} = x^{k-1} \cdot \sum_{j=1}^{n-k+1} x^j$$

Then we have

$$\begin{aligned} S &= \sum_{k=1}^n \sum_{j=k}^n kx^j \\ &= \sum_{k=1}^n k \sum_{j=k}^n x^j \\ &= \sum_{k=1}^n k \cdot \frac{x^k - x^{n+1}}{1 - x} \\ &= \frac{1}{1 - x} \cdot \sum_{k=1}^n kx^k - \frac{x^{n+1}}{1 - x} \cdot \sum_{k=1}^n k \\ &= \frac{1}{1 - x} \cdot \frac{x \cdot (nx^{n+1} - (n+1)x^n + 1)}{(x-1)^2} - \frac{x^{n+1}}{1 - x} \cdot \frac{n(n+1)}{2} \end{aligned}$$

# Chapter 3

## Recurrences

Sometimes recurrences are closely related to sums. Thus, if we are going to find the closed-form formula, we can express the sum as a recurrence, then the problem will be comparatively easier.

### 3.1 Homogeneous Recurrences

**Definition 3.1.1 (Linear homogeneous recurrence).** A linear homogeneous recurrence relation of degree  $d$  with constant coefficients is a recurrence relation of the form

$$T(n) = a_1T(n-1) + a_2T(n-2) + \cdots + a_dT(n-d),$$

where  $a_1, a_2, \dots, a_k \in \mathbb{R}$  are given constants and  $a_k \neq 0$

To solve for homogeneous recurrences with distinct root, we can use the following procedure:

1. Solve the characteristic equation to get root  $r_1, \dots, r_d$ .
2. If the roots are all distinct, form, the candidate solution

$$T_0(n) = \theta_1 r_1^n + \theta_2 r_2^n + \cdots + \theta_d r_d^n$$

3. Use the initial conditions on  $T(1), \dots, T(d)$  to determine  $\theta_1, \dots, \theta_d$

**Example.**

$$\begin{cases} T(n) = T(n-1) + 2T(n-2) & \text{for } n \geq 2 \\ T(0) = 2, T(1) = 7 \end{cases}$$

Let  $T(n) = x^n$ . Then, for characteristic equation, we have

$$\begin{aligned} x^n &= x^{n-1} + 2x^{n-2} \\ x^2 &= x + 2 \quad (\text{characteristic equation}) \end{aligned}$$

The roots are  $r_1 = 2, r_2 = -1$ . Since they are distinct, we can form the candidate solution

$$T_0(n) = \theta_1 2^n + \theta_2 (-1)^n.$$

By using the initial conditions, we have

$$\begin{aligned} 2 &= T_0(0) = \theta_1 + \theta_2 \\ 7 &= T_0(1) = 2\theta_1 - \theta_2 \end{aligned}$$

Solving above, we have  $\theta_1 = 3, \theta_2 = -1$ . Then, we have

$$T(n) = 3 \times 2^n - (-1)^n$$

However, if the root of the characteristic equation has a multiplicity  $m \geq 1$ , i.e., the root is repeated for  $m$  times, then we have  $T(n) = n^{m-1}x^n$ . Yet the procedures are the same as solving linear homogeneous recurrence with distinct roots.

**Example.**

$$\begin{cases} T(n) = 2T(n-1) - T(n-2) & \text{for } n \geq 2, \\ T(0) = 0, T(1) = 1 \end{cases}$$

Characteristic equation:  $x^2 = 2x - 1$ .

Solving above we have  $r_1 = 1$  with multiplicity  $m_1 = 2$ . Then we have

$$T_0(n) = \theta_1(1)^n + n\theta_2(1)^n = \theta_1 + n\theta_2$$

By using initial conditions, we have

$$\begin{aligned} 0 &= T_0(0) = \theta_1 \\ 1 &= T_0(1) = \theta_1 + \theta_2 \end{aligned}$$

Then, it follows that the solution to the recurrences is given by

$$T(n) = n$$

Let's see another example

**Example.**

$$\begin{cases} T(n) = 4T(n-1) - 5T(n-2) + 2T(n-3) & \text{for } n \geq 3, \\ T(0) = 0, T(1) = 1, T(2) = 3. \end{cases}$$

Characteristic equation:  $x^3 = 4x^2 - 5x + 2$

This equation has two distinct roots,  $r_1 = 1, r_2 = 2$ . The multiplicity of  $r_1$  is  $m_1 = 2$ . Hence,

$$T_0(n) = \theta_1(1)^n + n\theta_2(1)^n + \theta_32^n = \theta_1 + n\theta_2 + \theta_32^n$$

Using initial conditions, we have

$$\begin{aligned} 0 &= T_0(0) = \theta_1 + \theta_3 \\ 1 &= T_0(1) = \theta_1 + \theta_2 + 2\theta_3 \\ 3 &= T_0(2) = \theta_1 + 2\theta_2 + 4\theta_3 \end{aligned}$$

Solving for above, we have

$$T(n) = -1 + n \times 0 + 1 \times 2^n = -1 + 2^n$$

## 3.2 Non-homogeneous Recurrences

A recurrence relation of the form

$$T(n) = a_1T(n-1) + a_2T(n-2) + \cdots + a_dT(n-d) + f(n)$$

is called non-homogeneous recurrences.

To solve for non-homogeneous recurrences, we can use the following procedures:

1. Solve the associated linear homogeneous recurrence.
2. Find the particular solution  $T_p(n)$  to the linear non-homogeneous recurrence by examining the function class of  $f(n)$ .
3. Form the candidate solution  $T_0(n) = T_h(n) + T_p(n)$ , and use the initial conditions to find the parameters in  $T_h(n)$ .

In general, to solve non-homogeneous recurrence, we can consider the following particular solutions:

$f(n)$	$T_p(n)$
$s$	$x_0$
$n$	$x_1n + x_0$
$n^2$	$x_2n^2 + x_1n + x_0$
$s^n$	$x_0s^n$
$ns^n$	$(x_1n + x_0)s^n$

Let's see some examples

**Example.** Consider the recurrence

$$\begin{cases} T(n) = 2T(n-1) + 1 & \text{for } n \geq 1 \\ T(1) = 1 \end{cases}$$

Let  $T_0(n) = T_h(n) + T_p(n)$ , where  $T_h(n) = 2T(n-1)$ .

For  $T_h(n)$ , characteristic equation:  $x = 2$ , then we have  $T_h(n) = \theta 2^n$ .

Since  $f(n) = 1$ , let  $T_p(n) = x$ ,

$$x = 2x + 1$$

$$x = -1$$

Then, we have

$$T_0(n) = T_h(n) + T_p(n) = \theta 2^n - 1$$

Using the initial conditions, we have

$$1 = T_0(1) = 2\theta - 1$$

This gives  $\theta = 1$ . Hence, the solution to the recurrence is given by

$$T(n) = 2^n - 1$$

**Example.** Consider the recurrence

$$\begin{cases} T(n) = 5T(n-1) - 6T(n-2) + 7^n & \text{for } n \geq 2 \\ T(0) = 0, T(1) = 1. \end{cases}$$

Let  $T_0(n) = T_h(n) + T_p(n)$ , where  $T_h(n) = 5T(n-1) - 6T(n-2)$ .

For  $T_h(n)$ , characteristic equation:  $x^2 = 5x - 6$ , with  $r_1 = 3, r_2 = 2$ .

Hence, we have  $T_h(n) = \theta_1 3^n + \theta_2 2^n$ , which is the homogeneous solution.

Since  $f(n) = 7^n$  and  $s = 7$  is not a root of the characteristic equation. Let  $T_p(n) = x_0 7^n$  (particular solution),

$$x_0 7^n = 5x_0 7^{n-1} - 6x_0 7^{n-2} + 7^n$$

$$x_0 = 5x_0 7^{-1} - 6x_0 7^{-2} + 1$$

$$x_0 - \frac{5}{7}x_0 + \frac{6}{49}x_0 = 1$$

$$x_0 = \frac{49}{20}$$

Then, we have

$$T_0(n) = T_h(n) + T_p(n) = \theta_1 3^n + \theta_2 2^n + \frac{49}{20} 7^n$$



Using the initial conditions, we have

$$\begin{aligned} 0 &= T_0(0) = \theta_1 + \theta_2 + \frac{49}{20} \\ 1 &= T_0(1) = 3\theta_1 + 2\theta_2 + \frac{343}{20} \end{aligned}$$

This gives  $\theta_1 = -\frac{225}{20}, \theta_2 = \frac{176}{20}$ . Hence, the solution to the recurrence is given by

$$T(n) = -\frac{225}{20}3^n + \frac{176}{20}2^n + \frac{49}{20}7^n$$

**Remark.** Let  $f(n) = s^n$ ,  $r_1$  and  $r_2$  be the roots of the characteristic equation. Then

- If  $s \neq r_1, s \neq r_2$ , then  $T_p(n) = x_0 s^n$ ;
- If  $s = r_1, r_1 \neq r_2$ , then  $T_p(n) = x_0 n s^n$ ;
- If  $s = r_1 = r_2$ , then  $T_p(n) = x_0 n^2 s^n$ ;

where  $x_0$  is constant to be determined in all cases.

**Example.** Consider the recurrence

$$\begin{cases} T(n) = 6T(n-1) - 9T(n-2) + 3^n & \text{for } n \geq 2 \\ T(0) = 0, T(1) = \frac{1}{2}. \end{cases}$$

Let  $T_0(n) = T_h(n) + T_p(n)$ , where  $T_h(n) = 6T(n-1) - 9T(n-2)$ .

For  $T_h(n)$ , characteristic equation:  $x^2 = 6x - 9$ ,  $r_1 = 3$  with multiplicity  $m_1 = 2$ .

Hence, we have  $T_h(n) = \theta_1 3^n + n\theta_2 3^n$ , which is the homogeneous solution.

Since  $f(n) = 3^n$  and  $s = 3$  is also a root of the characteristic equation. Let  $T_p(n) = x_0 n^2 3^n$  (particular solution),

$$\begin{aligned} x_0 n^2 3^n &= 6x_0(n-1)^2 3^{n-1} - 9x_0(n-2)^2 3^{n-2} + 3^n \\ 9x_0 n^2 &= 18x_0(n-1)^2 - 9x_0(n-2)^2 + 9 \\ 9x_0 n^2 &= 18x_0(n^2 - 2n + 1) - 9x_0(n^2 - 4n + 4) + 9 \\ 9x_0 n^2 &= 9x_0 n^2 - 18x_0 + 9 \\ x_0 &= \frac{1}{2} \end{aligned}$$

Then, we have

$$T_0(n) = T_h(n) + T_p(n) = \theta_1 3^n + n\theta_2 3^n + \frac{1}{2} n^2 3^n$$

Using the initial conditions, we have

$$\begin{aligned} 0 &= T_0(0) = \theta_1 \\ \frac{1}{2} &= T_0(1) = 3\theta_1 + 3\theta_2 + \frac{3}{2} \end{aligned}$$

This gives  $\theta_1 = 0, \theta_2 = -\frac{1}{3}$ . Hence, the solution to the recurrence is given by

$$T(n) = -\frac{1}{3} n 3^n + \frac{1}{2} n^2 3^n$$

# Chapter 4

## Asymptotics

Asymptotic notation is a shorthand used to give a quick measure of the behavior of a function  $f(n)$  as  $n$  grows large.

### 4.1 Big O

Big O is the most frequently used asymptotic notation. It is used to give an upper bound on the growth of a function, such as the running time of an algorithm.

**Definition 4.1.1.** We say that  $f(x) = O(g(x))$  iff there exists a constant  $c > 0$  and an  $x_0 \geq 0$  such that

$$f(x) \leq cg(x) \quad \text{for all } x \geq x_0$$

Or we can use the following notation

$$\exists c > 0, x_0 \geq 0 \text{ such that } f(x) \leq cg(x) \forall x \geq x_0$$

**Example.** Let  $f(x) = x^2$  and  $g(x) = x^3$ .

By taking  $c = 1$  and  $x_0 = 1$ , we can simply conclude that  $x^2 = O(x^3)$ .

However, to prove that **there is no constant  $c$  such that  $x^2 \leq cx^3$  for all  $0 \leq x \leq 1$** , we need to use proof by contradiction.

For  $0 \leq x \leq 1$ , we have  $x^2 \geq x^3$ . Therefore, if such a constant  $c > 0$  exists, then we must have  $c \geq 1$  such that  $x^2 \leq cx^3$ . However, whenever  $0 \leq x \leq \frac{1}{c}$ , we have

$$x^3 \geq \frac{1}{c}x^2 > x^3.$$

Since we get  $x^3 > x^3$ , by contradiction, we know that there is no constant  $c$  such that  $x^2 \leq cx^3$  for all  $0 \leq x \leq 1$ .

We can also use differentiation to show the upper bound of a function.

**Example.** Let  $f(x) = \ln x$  and  $g(x) = x$ .

Observe that  $f(1) = \ln 1 = 0 < 1 = g(1)$ . Moreover,  $f'(x) = \frac{1}{x}$  and  $g'(x) = 1$  for all  $x > 0$ , which implies that  $g'(x) > f'(x)$  for all  $x > 1$ . It is the same as saying that  $g(x)$  grows faster than the function  $f(x)$  because the slope of the former is larger than that of the latter. It follows that

$$\ln x \leq x \text{ for all } x \geq 1.$$

Hence, by taking  $c = 1$  and  $x_0 = 1$  we have  $\ln x = O(x)$ .

**Exercise.** Let  $f(x) = x^2$  and  $g(x) = 2^x$ . Show that  $f(x) = O(g(x))$ .

**Example.** Let

$$f(n) = n \cdot (n+1) \cdot (n+2) + (-1)^n, \quad g(n) = n^\alpha$$

for any integer  $n \geq 1$ . What is the smallest  $\alpha \in \mathbb{R}$  such that  $f(n) = O(g(n))$ ? Note that

$$\begin{aligned} |f(n)| &= |n \cdot (n+1) \cdot (n+2) + (-1)^n| \\ &= |n^3 + 3n^2 + 2n + (-1)^n| \\ &= n^3 + 3n^2 + 2n + (-1)^n \quad (\text{for any } n \geq 1) \end{aligned}$$

We claim that  $\alpha = 3$  is the smallest  $\alpha \in \mathbb{R}$  such that  $f(n) = O(g(n))$ .

Let us first prove that  $\alpha$  satisfies  $\alpha \geq 3$  and then that  $\alpha = 3$ .

Indeed, we must have  $\alpha \geq 3$ . The proof is by contradiction. If  $\alpha < 3$ , then

$$\lim_{n \rightarrow \infty} \frac{|f(n)|}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^3 + 3n^2 + 2n + (-1)^n}{n^\alpha} = \infty,$$

which shows that  $f(n) = \omega(g(n))$ . However, this implies that  $f(n) = O(g(n))$  is wrong. We have reached the promised contradiction.

So let us now show that for  $\alpha = 3$  we have  $f(n) = O(g(n))$ . Indeed, choosing  $c = 7$  and  $n_0 = 1$ , we get

$$\begin{aligned} |f(n)| &= n^3 + 3n^2 + 2n + (-1)^n \\ &\leq n^3 + 3n^2 \cdot n + 2n \cdot n^2 + n^3 \\ &= 7n^3 \\ &= cn^3 \quad (\text{for all } n \geq n_0) \end{aligned}$$

## 4.2 Big Omega

As we use Big O notation to express upper bound, for lower bound, we have the "Big Omega" notation.

**Definition 4.2.1.** We say that  $f(x) = \Omega(g(x))$  iff there exists a constant  $c > 0$  and an  $x_0 \geq 0$  such that

$$f(x) \geq cg(x) \quad \text{for all } x \geq x_0$$

Since Big O and Big Omega are essentially "mirror image" of one another, we have

**Theorem 4.2.1** (Big O vs. Big Omega).

$$f(x) = O(g(x)) \iff g(x) = \Omega(f(x)).$$

**Proof.** By definition of big O,  $f(x) = O(g(x))$  means

$$\exists c_1 > 0, x_1 > 0 \text{ such that } f(x) \leq c_1 g(x) \quad \forall x \geq x_1,$$

which means the same as

$$\exists c_1 > 0, x_1 > 0 \text{ such that } g(x) \geq \frac{1}{c_1} f(x) \quad \forall x \geq x_1,$$

Hence, by taking  $c_0 = \frac{1}{c_1} > 0$  and  $x_0 = x_1 \geq 0$ , we have  $g(x) = \Omega(f(x))$  ■

## 4.3 Theta

Theta can be understood as the approximation of a function.

**Definition 4.3.1.** We say that  $f(x) = \Theta(g(x))$  iff  $f(x) = O(g(x))$  and  $g(x) = O(f(x))$ .

**Remark.** It is important to note that  $f(x) = \Theta(g(x))$  does not mean that  $f(x) = g(x)$ , it just means that

$$\exists c_{1,2} > 0 \quad \text{such that} \quad c_1 g(x) \leq f(x) \leq c_2 g(x) \quad \forall x \geq x_0$$

## 4.4 Little O

Little O notation can be understood as the strict upper bound on the growth of a function.

**Definition 4.4.1 (Little O notation).** For functions  $f, g : \mathbb{R} \rightarrow \mathbb{R}$ , with  $g$  nonnegative, we say  $f$  is asymptotically smaller than  $g$ , in symbols,

$$f(x) = o(g(x)) \quad \text{iff} \quad \lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0.$$

**Example.** For example, let  $f(x) = x, g(x) = e^x - 1$ . Because

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x}{e^x - 1} = 0.$$

Hence, we have  $f(x) = o(g(x))$

**Example.** Let

$$f(n) = \begin{cases} 0 & \text{if } n \text{ is odd,} \\ 2 & \text{if } n \text{ is even.} \end{cases}$$

Is  $f(n) = o(1)$ ?

Let  $g(n) = 1$ . Upon noting  $f(n) = 1 + (-1)^n$ , we see that

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} f(n).$$

However, this limit does not exist, as  $f(n)$  fluctuates between 0 and 2. Thus,  $f(n) \neq o(1)$ .

On the other hand, if  $f(n) = o(n)$ ? Here, let  $g(n) = n$ . Note that  $0 \leq f(n) \leq 2$ . It follows that

$$0 \leq \frac{f(n)}{g(n)} \leq \frac{2}{n}$$

for  $n \geq 1$ . By the sandwich theorem, we then obtain

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0.$$

Thus,  $f(n) = o(n)$ .

## 4.5 Little Omega

Little Omega notation can be understood as the strict lower bound on the growth of a function.

**Definition 4.5.1 (Little O notation).** For functions  $f, g : \mathbb{R} \rightarrow \mathbb{R}$ , with  $g$  nonnegative, we say  $f$  is asymptotically smaller than  $g$ , in symbols,

$$f(x) = \omega(g(x)) \quad \text{iff} \quad \lim_{x \rightarrow \infty} \frac{g(x)}{f(x)} = 0.$$

## 4.6 Properties for Asymptotic Analysis

### 4.6.1 Rules for Asymptotic Analysis

- Transitivity

If  $f(n) = \Pi(g(n))$  and  $g(n) = \Pi(h(n))$ , then  $f(n) = \Pi(h(n))$ , where  $\Pi = O, o, \Omega, \omega, \Theta$ .

- Rule of sums

$$f(n) + g(n) = \Pi(\max\{f(n), g(n)\}), \quad \text{where } \Pi = O, \Omega, \text{ or } \Theta.$$

- Rule of products

If  $f_1(n) = \Pi(g_1(n))$ ,  $f_2(n) = \Pi(g_2(n))$ , then  $f_1(n)f_2(n) = \Pi(g_1(n)g_2(n))$ , where  $\Pi = O, o, \Omega, \omega, \Theta$ .

- Transpose symmetry

$$f(n) = O(g(n)) \quad \text{iff} \quad g(n) = \Omega(f(n)).$$

- Transpose symmetry

$$f(n) = o(g(n)) \quad \text{iff} \quad g(n) = \omega(f(n)).$$

- Reflexivity


$$f(n) = \Pi(f(n)), \quad \text{where } \Pi = O, \Omega, \Theta.$$

- Symmetry

$$f(n) = \Theta(g(n)) \quad \text{iff} \quad g(n) = \Theta(f(n)).$$

### 4.6.2 Graph for Functions

One can understand the growth of functions by the following graph.



## Chapter 5

# Set Theory and Counting Principle

### 5.1 Set Theory

**Definition 5.1.1.** Set is a collection of elements, in which each element appears only once.

**Definition 5.1.2.** Given a set  $S$ , the number of elements in  $S$  is denoted by  $|S|$ . The quantity  $|S|$  is also referred to as the **cardinality** of  $S$ .

A set with no elements in it is called an empty set, which is denoted by  $\emptyset$ .

**Definition 5.1.3.** Given two sets  $S$  and  $T$ , we say that  $T$  is a subset of  $S$ , denoted by  $T \subseteq S$ , if every element in  $T$  is also in  $S$ . It follows that if  $T \subseteq S$ , then  $|T| \leq |S|$ .

For example, let

$$S = \{1, 2, 3, 4, 5\}, \quad T = \{2, 4\}, \quad U = \{2, 4, 6\}$$

Then, we have  $T \subseteq S$  and  $T \subseteq U$ .

**Definition 5.1.4.** Let  $S_1$  and  $S_2$  be two given sets.

The union of  $S_1$  and  $S_2$ , denoted by  $S_1 \cup S_2$ , is the set containing all elements from both  $S_1$  and  $S_2$ .

The intersection of  $S_1$  and  $S_2$ , denoted by  $S_1 \cap S_2$ , is the set containing all elements that are common to both  $S_1$  and  $S_2$ .

We can also see union as the relationship "or", while intersection is the relationship "and".

**Definition 5.1.5.** Now, let  $S$  be a set and  $m \geq 1$  be an integer. A partition of  $S$  into  $m$  parts is a collection of  $m$  subsets of  $S$ , denoted by  $S_1, \dots, S_m$ , with the following properties:

- Exhaustion:  $S = S_1 \cup S_2 \cup \dots \cup S_m$ .
- Non-Overlapping: For  $i \neq j$ , we have  $S_i \cap S_j = \emptyset$

### 5.2 Counting Principle

#### 5.2.1 Addition Principle

Given  $S = \{1, 2, 3, 4, 5\}$ . Let  $S_1 = \{1, 2\}, S_2 = \{3\}, S_3 = \{4, 5\}$  form a partition of  $S$  with 3 parts. Then, the cardinality of  $S$  can be determined by the cardinalities of the constituent parts, i.e.

$$|S| = |S_1| + |S_2| + \dots + |S_m|.$$

**Example** (Count the number of binary strings of length  $n$  with no consecutive 1's.). Let  $S$  be the desired string, then we can apply the addition principle.

Let  $S_0$  be the set of strings in  $S$  that start with 0;  $S_1$  be the set of strings in  $S$  that start with 1. We have  $|S| = |S_0| + |S_1|$ .

We can set up a recurrence relationship. Let  $|S| = T(n)$ . Then, we have  $|S_0| = T(n-1)$  and  $|S_1| = T(n-2)$ . Then we have

$$|S| = T(n) = T(n-1) + T(n-2)$$

Using the initial conditions  $T(1) = 2$  (string starts with 1 or 0) and  $T(2) = 3$  (if the string starts with 1, then we have only one option for the next digit; otherwise, we have two options), one can find the number of the desired binary strings.

### 5.2.2 Multiplication Principle

If each element in  $S$  can be generated by performing an ordered sequence of actions, say,  $A_1, A_2, \dots, A_N$ , and action  $A_i$  has  $p_i$  choices, where  $i = 1, \dots, N$ , then the cardinality of  $S$  can be computed by

$$|S| = p_1 \times p_2 \times \dots \times p_N$$

**Example.** To count the number of integers between 0 and 9999 that have exactly one digit equal to 5, let  $S$  be the set of such integers. We can then partition this set to  $S_1, S_2, S_3, S_4$ , where  $S_1$  is the set of integers in  $S$  with "5" appearing in the first position from the right, and so on. For example,  $2051 \in S_2$ . By the addition principle, we have

$$|S| = |S_1| + |S_2| + |S_3| + |S_4|$$

Since every integer in  $S_1$  takes the form  $xxx5$ , with each " $x$ " having 9 choices. We then have  $|S_1| = 9^3$ . It holds true for the other sets. Hence, we have  $|S| = 4 \times 9^3 = 2916$

**Example.** Let us count the number of odd integers between 1000 and 9999 that have all distinct digits. Let  $S$  be the set of such integers.

**Method 1:** Observe that each integer in  $S$  can be generated by the following ordered sequence of action:

action	choices	no. of choices
$A_1$ : pick the unit digit	$\{1, 3, 5, 7, 9\}$	5
$A_2$ : pick the tens digit	$\{0, 1, \dots, 9\}$ except the digit chosen in $A_1$	9
$A_3$ : pick the hundreds digit	$\{0, 1, \dots, 9\}$ except the digit chosen in $A_1, A_2$	8
$A_4$ : pick the thousands digit	$\{1, \dots, 9\}$ except the digit chosen in $A_1, A_2, A_3$	?

We cannot find the choice for  $A_4$  directly since 0 may or may not be chosen in the previous cases. Thus, we can partition  $S$  into

$S_1$ : set of integers in  $S$  whose tens digit is 0  $\Rightarrow |S_1| = 5 \times 1 \times 8 \times 7 = 280$ ,

$S_2$ : set of integers in  $S$  whose hundreds digit is 0  $\Rightarrow |S_2| = 5 \times 8 \times 1 \times 7 = 280$ ,

$S_3$ : set of integers in  $S$  with no 0  $\Rightarrow |S_3| = 5 \times 8 \times 7 \times 6 = 1680$ .

Then we have  $|S| = 280 + 280 + 1680 = 2240$ .

**Method 2:** Consider a different sequence of actions:

action	choices	no. of choices
$A_1$ : pick the unit digit	$\{1, 3, 5, 7, 9\}$	5
$A_2$ : pick the thousands digit	$\{1, \dots, 9\}$ except the digit chosen in $A_1$	8
$A_3$ : pick the tens digit	$\{0, 1, \dots, 9\}$ except the digit chosen in $A_1, A_2$	8
$A_4$ : pick the hundreds digit	$\{0, 1, \dots, 9\}$ except the digit chosen in $A_1, A_2, A_3$	7

Then we have  $|S| = 5 \times 8 \times 8 \times 7 = 2240$ .

### 5.2.3 Subtraction Principle

**Definition 5.2.1.** Let  $S$  be a set and  $A \subseteq S$  be a subset of  $S$ . The complement of  $A$  in  $S$ , denoted by  $\overline{A}$ , is the set that contains all the elements in  $S$  but not in  $A$ .

$$|S| = |A| + |\overline{A}|$$

**Example.** Consider computer passwords of length 6, each symbol of which is taken from 0, 1, ..., 9 and a, b, ..., z. We would like to count the number of passwords that have repeated symbols. Let  $A$  be the set of such passwords. For instance, we have  $1223aq, bb333k \in A$ , but  $123456 \notin A$ .

We can then use the subtraction principle, i.e. count the number of passwords that contains all distinct symbols. Then, we have

$$|A| = |S| - |\overline{A}| = 36^6 - P(36, 6)$$

### 5.2.4 Division Principle

Division principle states that if  $S$  is partitioned into  $k$  equal-sized parts, then

$$k = \frac{|S|}{\text{number of elements in each part}}.$$

**Definition 5.2.2 (Ceil function).** Given a real number  $x$ , define

$$\lceil x \rceil = \text{least integer } \geq x$$

The symbol  $\lceil x \rceil$  is usually read as "the ceil of  $x$ ".

**Proposition 5.2.1 (Pigeonhole Principle).** Suppose that  $n$  objects are placed into  $k$  boxes. Then, at least one box has at least  $\lceil \frac{n}{k} \rceil$  objects.

**Example.** 51 distinct numbers are chosen from the integers between 1 and 100 inclusively.

Let the 50 pairs of consecutive integers

$$\{1, 2\}, \{3, 4\}, \dots, \{99, 100\}$$

be the pigeonholes and the 51 numbers be the pigeons. Then two of the 51 numbers must be in the same pigeonholes. Therefore, there are 2 consecutive integers among the 51 chosen integers.



## 5.3 Permutation and Combination

### 5.3.1 Permutation

**Example.** Consider a set of  $n$  elements:  $S_0 = \{1, \dots, n\}$ . We call  $S_0$  the ground set. Also, let  $r \geq 1$  be integer. An  $r$ -permutation of the  $n$ -elements ground set  $S_0$  is an ordered selection of  $r$  elements from  $S_0$ . Let  $S$  be the set of all different  $r$ -permutations of the  $n$ -element ground set  $S_0$ . For instance, when  $S_0 = \{1, 2, 3\}$  and  $r = 2$ , we have

$$S = \{(1, 2), (2, 1), (1, 3), (3, 1), (2, 3), (3, 2)\},$$

where the order of the two elements matters.

For general values of  $n$  and  $r$ . Let  $P(n, r)$  denotes this number. By performing the following ordered sequence of action,

action	number of choices
$A_1$ : pick the 1st element	$n$
$A_2$ : pick the 2nd element	$n - 1$
$\vdots$	$\vdots$
$A_r$ : pick the $r$ th element	$n - r + 1$

Thus, we have

$$P(n, r) = n(n - 1) \cdots (n - r + 1) = \frac{n!}{(n - r)!}$$

### 5.3.2 Combination

**Example.** Consider the ground set  $S_0 = \{1, \dots, n\}$  of  $n$  elements. Let  $r \geq 1$  be integer. An  $r$ -combination of the  $n$ -element ground set  $S_0$  is an unordered selection of  $r$  elements from  $S_0$ . Let  $S$  be the set of all different  $r$ -combinations of the  $n$ -element ground set  $S_0$ . For instance, when  $S_0 = \{1, 2, 3\}$  and  $r = 2$ , we have

$$S = \{(1, 2), (1, 3), (2, 3)\}$$

For general values of  $n$  and  $r$ , we can determine  $\binom{n}{r}$  by relating  $r$ -combinations to  $r$ -permutations. Indeed, observe that each  $r$ -permutation of  $S_0$  can be generated via the following ordered sequence of action:

$A_1$ : pick  $r$  elements from  $S_0$

$A_2$ : order the  $r$  elements chosen from  $A_1$  to form the desired  $r$ -permutation.

Note that  $A_1$  has  $\binom{n}{r}$  choices, while  $A_2$  has  $r!$  choices. Hence, by the multiplication principle,

$$P(n, r) = \text{number of } r\text{-permutation of } S_0 = \binom{n}{r} \times r!$$

Since

$$P(n, r) = \frac{n!}{(n - r)!},$$

it follows that

$$\binom{n}{r} = \frac{n!}{r!(n - r)!}$$

**Corollary 5.3.1.**

$$\binom{n}{0} = 1, \binom{n}{1} = n, \binom{n}{n} = 1, \binom{n}{r} = 0 \text{ (whenever } n < r).$$

## Chapter 6

# Binomial Coefficients

### 6.1 Introduction

In this section, we introduce Binomial Coefficients.

#### 6.1.1 Combinations

As it is introduced before,

**Definition 6.1.1.** An  $r$ -combination of the  $n$ -element ground set  $S_0$  is an **unordered selection** of  $r$  elements from  $S_0$ .

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

#### 6.1.2 Permutation

Again, as it is introduced before,

**Definition 6.1.2.** An  $r$ -permutation of the  $n$ -element ground set  $S_0$  is an **ordered selection** of  $r$  elements from  $S_0$ .

$$P(n, r) = \frac{n!}{(n-r)!}$$

#### 6.1.3 Binomial Identities

**Proposition 6.1.1.** For any integers  $m, r \geq 0$  with  $0 \leq r \leq n$ ,

$$\binom{n}{r} = \binom{n}{n-r}$$

We have two ways to prove this proposition, namely Algebraic Proof and Combinatorial Proof.

**Algebraic Proof.**

$$\binom{n}{n-r} = \frac{n!}{(n-r)!(n-n+r)!} = \frac{n!}{r!(n-r)!} = \binom{n}{r}$$

■

**Combinatorial Proof.** Both side of the identity are supposed to be two different ways of solving a counting problem. We can define the counting problem as counting the number of different unordered selections of  $r$  elements from an  $n$ -element ground set. Then, we can define the RHS as the selecting number to be excluded. Then this identity holds. ■

### 6.1.4 Pascal's Identity

**Theorem 6.1.1.** For any integers  $n, r \geq 0$  with  $1 \leq r \leq n - 1$ ,

$$\binom{n}{r} = \binom{n-1}{r} + \binom{n-1}{r-1} \text{ OR } \binom{n+1}{r} = \binom{n}{r} + \binom{n}{r-1}$$

Again, we can prove this theorem by two ways.

**Algebraic Proof.**

$$\begin{aligned} RHS &= \frac{(n-1)!}{r!(n-r-1)!} + \frac{(n-1)!}{(r-1)!(n-1-r+1)!} \\ &= \frac{(n-1)!}{r!(n-r-1)!} + \frac{(n-1)!}{(r-1)!(n-r)!} \\ &= \frac{(n-1)!}{(r-1)!(n-r-1)!} \left( \frac{1}{r} + \frac{1}{n-r} \right) \\ &= \frac{(n-1)!}{(r-1)!(n-r-1)!} \left( \frac{n}{r(n-r)} \right) \\ &= \frac{n!}{r!(n-r)!} \\ &= \binom{n}{r} \end{aligned}$$

■

**Combinatorial Proof.** Recall that  $\binom{n}{k}$  equals the number of subsets with  $k$  elements from a set with  $n$  elements. Suppose one particular element is uniquely labeled  $X$  in a set with  $n$  elements.

To construct a subset of  $k$  elements containing  $X$ , include  $X$  and choose  $k - 1$  elements from the remaining  $n - 1$  elements in the set. There are  $\binom{n-1}{k-1}$  such subsets.

To construct a subset of  $k$  elements **not** containing  $X$ , choose  $k$  elements from the remaining  $n - 1$  elements in the set. There are  $\binom{n-1}{k}$  such subsets.

Every subset of  $k$  elements either contains  $X$  or not. The total number of subsets with  $k$  elements in a set of  $n$  elements is the sum of the number of subsets containing  $X$  and the number of subsets that do not contain  $X$ ,  $\binom{n-1}{k-1} + \binom{n-1}{k}$ .

This equals  $\binom{n}{k}$ .

■

## Chapter 7

# Elements of Discrete Probability

### 7.1