

1. The three models AR(2), MA(1), and ARMA(2,1) are fitted to the following time series:

Year	Quarter 1	Quarter 2	Quarter 3	Quarter 4
2010	112	104	100	96
2011	101	101	105	94
2012	106	106	108	110

The results using R are as follows:

AR(2):			MA(1):	
	ar1	ar2	intercept	
	0.2618	0.0868	104.227	
st. error	0.3289	0.3326	2.372	
$\sigma^2 = 25.02$	log likelihood = -36.4			

  

	ma1	intercept
	0.1899	103.798
st. error	0.2574	1.7373
$\sigma^2 = 25.66$	log likelihood = -36.51	

  

	ar1	ar2	ma1	intercept
	-0.5547	0.4437	0.9779	104.4542
st. error	0.3329	0.33267	0.235	2.4584
$\sigma^2 = 21.31$	log likelihood = -36.02			

The models are ranked using Akaike Information Criterion, Corrected for bias (AICC) (see §11.3 of Lecture Notes or slide 32 of week 5).

$AICC = -2 \times \log\text{-likelihood} + 2 \times \frac{rn}{n-r-1}$  where  $n$  is the sample size and  $r$  is the number of free parameters, most often  $r = p + q + 1$ .

Determine the order from the best to worst model. Give full explanation on how you arrived to your answer. Show calculations.

- AR(2), MA(1), ARMA (2,1)
- AR(2), ARMA (2,1), MA(1)
- MA(1), AR(2), ARMA (2,1)
- MA(1), ARMA (2,1), AR(2)
- ARMA (2,1), AR(2), MA(1).

**2.** In modeling the weekly sales of a certain commodity over the past six months, the time series model  $X_t - \phi_1 X_{t-1} = Z_t + \theta_1 Z_{t-1}$  was thought to be appropriate. Suppose the model was fitted and the autocorrelations of the residuals were:

k	1	2	3	4	5	6	7	8
$\hat{\rho}_{\hat{W}}(k)$	-.04	-.50	.03	-.01	.01	.02	.03	-.01
st. dev $\hat{\rho}_{\hat{W}}(k)$	.08	.10	.11	.11	.11	.11	.11	.11

Is the assumed model really appropriate? If not, how would you modify the model? Explain.

*Hint: Check slides 8 - 9 of Lecture 11 for the 95% confidence intervals for autocorrelation function of the fitted residuals. You might also find slide 26 of Week 2 and slide 15 of Lecture 11 useful.*

**3.** Suppose that in a sample of size 100 from an AR(1) process with mean  $\mu$ ,  $\phi = 0.6$ , and  $\sigma^2 = 2$ , we obtain  $\bar{x}_{100} = 0.271$ . Construct an approximate 95% confidence interval for  $\mu$ . Are the data compatible with the hypothesis that  $\mu = 0$ ?

*Help: (i) Here  $\sigma^2 \equiv \sigma_Z^2$  is the variance of noise  $Z_t$ .*

*(ii) Large sample distribution of  $\bar{X}_n$  is given as follows (See, for example, slide 48 or §10.1 of Week 4): For  $n$  large, distribution of the sample mean  $\bar{X}_n$  is approximately normal with mean  $\mu \equiv EX_t$  and variance  $n^{-1}v$ , where  $v \approx \gamma_X(0) + 2 \sum_{h=1}^{\infty} \gamma_X(h)$ .*

*(iii) For ACVF for AR(1) process see slide 43 of Week 2.*

**The following problem is for students enrolled in PSTAT 274 ONLY**

**G1.** Two hundred observations of a time series,  $X_1, \dots, X_{200}$ , gave the following sample statistics: sample mean:  $\bar{x}_{200} = 3.82$ ; sample variance:  $\hat{\gamma}(0) = 1.15$ ; sample ACF:  $\hat{\rho}(1) = .427$ ,  $\hat{\rho}(2) = .475$ ,  $\hat{\rho}(3) = .169$ .

(a) Based on these sample statistics, is it reasonable to suppose that  $\{X_t - \mu\}$  is a white noise?

(b) Assuming that  $\{X_t - \mu\}$  can be modeled as the AR(2) process,

$$X_t - \mu - \phi_1(X_{t-1} - \mu) - \phi_2(X_{t-2} - \mu) = Z_t,$$

where  $Z_t \sim \text{IID}(0, \sigma_Z^2)$ , find estimates of  $\mu, \phi_1, \phi_2$ , and  $\sigma_Z^2$ .

(c) Assuming that the data was generated from an AR(2) model, derive estimates of the PACF for all lags  $h \geq 1$ .