

# Permutations and Combinations: Takeaways



by Dataquest Labs, Inc. - All rights reserved © 2019

## Concepts

- If we have an experiment  $E_1$  (like flipping a coin) with **a** outcomes, followed by an experiment  $E_2$  (like rolling a die) with **b** outcomes, then the total number of outcomes for the composite experiment  $E_1E_2$  can be found by multiplying **a** with **b** (this is known as the **rule of product**):

$$\text{Number of outcomes} = a \times b$$

- If we have an experiment  $E_1$  with **a** outcomes, followed by an experiment  $E_2$  with **b** outcomes, followed by an experiment  $E_n$  with **z** outcomes, the total number of outcomes for the composite experiment  $E_1E_2 \dots E_n$  can be found by multiplying their individual outcomes:

$$\text{Number of outcomes} = a \times b \times \dots \times z$$

- There are two kinds of arrangements:
  - Arrangements where the order matters, which we call **permutations**.
  - Arrangements where the order doesn't matter, which we call **combinations**.
- To find the number of permutations when we're sampling with replacement, we can use the formula:

$$\text{Permutation} = n!$$

- To find the number of permutations when we're sampling without replacement and taking only **k** objects from a group of **n** objects, we can use the formula:

$${}_nP_k = \frac{n!}{(n-k)!}$$

- To find the number of combinations when we're sampling without replacement and taking only **k** objects from a group of **n** objects, we can use the formula:

$$\begin{equation} {}_nC_k = \{n \text{ choose } k\} = \frac{n!}{k!(n-k)!} \end{equation}$$

## Resources

- [A tutorial on calculating combinations when sampling with replacement](#), which we haven't covered in this mission
- [An easy-to-digest introduction to permutations and combinations](#)



Takeaways by Dataquest Labs, Inc. - All rights reserved © 2019