

MATH 104

REZA PAKZAD

*Davis Foote**

University of California, Berkeley

August 27th, 2015 – December 10th, 2015

CONTENTS

1	Natural Numbers	1
2	Real Numbers	2
2.1	Ordered Fields	2
2.2	Properties of the Real Numbers	4

1 NATURAL NUMBERS

*Lecture 1
August 27th, 2015*

1.1 DEFINITION. Peano axioms for the set of natural numbers:

(N1) $1 \in \mathbb{N}$

(N2) $n \in \mathbb{N} \Rightarrow \exists n + 1 \in \mathbb{N}$, called the **successor** of n

(N3) 1 is not the successor of any element of \mathbb{N}

(N4) $n + 1 = m + 1 \Rightarrow n = m$

(N5) A subset of \mathbb{N} containing 1 and containing $n + 1$ whenever it contains n must be the entire set \mathbb{N} .

*Lecture 2
September 1st, 2015*

*djfoote@berkeley.edu

2. REAL NUMBERS

There are some intuitions about the natural numbers which are not represented directly by these axioms. For example, we know that any natural number which is not 1 is the successor of some natural number.

1.2 THEOREM. $\forall n \in \mathbb{N} : n \neq 1 \Rightarrow \exists m \in \mathbb{N} : n = m + 1$

Proof. Let $n \in \mathbb{N}$ s.t. $n \neq 1$. Suppose $\forall m \in \mathbb{N}, n \neq m + 1$. Let $S = \mathbb{N} \setminus \{n\}$. Let $q \in S$. Then $q \in \mathbb{N}$ and $q \neq n$. Since $q + 1 \in \mathbb{N}$ by N2 and $q + 1 \neq n$ (since n is not the successor of any natural number), then $q + 1 \in S$. Since $n \neq 1, 1 \in S$. Therefore $S = \mathbb{N}$ by N5. But $n \in \mathbb{N}$ and $n \notin S$. Contradiction. \square

1.3 THEOREM (Well-Ordering Principle). *Any subset of the natural numbers admits a "least element." Logically,*

$$\forall S \subseteq \mathbb{N} : \exists n_0 \in S : \forall n \in S : n_0 \leq n + 1$$

TODO: Proof of WOP based on these Peano postulates

1.4 DEFINITION. For some $S \subseteq \mathbb{N}$, if

1. $1 \in S$
2. Whenever $\{1, 2, \dots, n\} \subset S$, then $n + 1 \in S$

then $S = \mathbb{N}$. This is called **strong induction**.

2 REAL NUMBERS

2.1 Ordered Fields

Lecture 3
September 3rd, 2015

Nicholas Bourbaki: school of thought putting forth that there are three main types of structures in mathematics:

- Algebraic structures $\xrightarrow{\text{binary operations}}$ Algebra
- Order structures $\xrightarrow{\text{inequalities}}$ Analysis
- Topological structures $\xrightarrow{\text{continuums, stretches}}$ Geometry/Topology

Goal: identify the "optimal" sets of axioms (related to the above three structures) which will uniquely determine the set of real numbers.

TODO: Incorporate notes from when I left early (on groups and fields)

Lecture 4
September 8th, 2015

2.1 DEFINITION. An **ordered field** is a tuple $(F, +, \cdot, \leq)$ with axioms:

- $(F, +)$ is an abelian group.
- $(F \setminus \{0\})$ is an abelian group.
- \cdot distributes over $+$.
- \leq is a total ordering on F (i.e. it is reflexive, antisymmetric, transitive, and total).
- $\forall a, b, c \in F : a \leq b \implies a + c \leq b + c$.
- $\forall a, b, c \in F : a \leq b \wedge 0 \leq c \implies a \cdot c \leq b \cdot c$

2.2 FACT. \mathbb{C} cannot be an ordered field for any ordering \leq .

The following are true in any ordered field:

- 1) $\forall a, b \in F : a \leq b \implies -b \leq -a$.
- 2) $\forall a, b \in F : a \leq b \wedge c \leq 0 \implies b \cdot c \leq a \cdot c$.

2.3 DEFINITION. Let F be a field. For all $a \in F$, define

$$|a| := \begin{cases} a & a \geq 0 \\ -a & a < 0 \end{cases}$$

This is called the **absolute value** of a .

Properties of the absolute value:

- (i) $\forall a \in F, |a| \geq 0$, and $|a| > 0$ if and only if $a \neq 0$.
- (ii) $\forall a, b \in F, |a \cdot b| = |a| \cdot |b|$.
- (iii) $\forall a, c \in F$ with $c \geq 0, |a| \leq c \iff -c \leq a \leq c$.
- (iv) $\forall a, b \in F, |a + b| \leq |a| + |b|$. This is the **triangle inequality**.

A useful consequence of the triangle inequality:

$$||a| - |b|| \leq |a - b|$$

2.4 DEFINITION. We say a is a **maximum** for $A \subset F$ if and only if $a \in A$ and $\forall x \in A, x \leq a$. A **minimum** is defined similarly.

2.5 DEFINITION. $a \in F$ is an **upper bound** for the set $A \subset F$ if and only if $\forall x \in A, x \leq a$. If such a bound exists, we say A is **bounded above**. Definitions for **lower bound** and **bounded below** are similar.

2. REAL NUMBERS

2.6 DEFINITION. $s \in F$ is a **supremum** (least upper bound) for the set A if and only if

- (i) s is an upper bound for A .
- (ii) For all upper bounds a for A , we have $s \leq a$.

We then say $s = \sup F$.

An **infimum** of F is defined identically as the greatest lower bound. If i is an infimum of F we say $i = \inf F$.

2.2 Properties of the Real Numbers

2.7 DEFINITION (Completeness Axiom). Any nonempty subset $A \subseteq \mathbb{R}$ which is bounded above admits a supremum in \mathbb{R} .

\mathbb{R} is identified as the only possible “complete” ordered field. Any other ordered field that satisfies the completeness axiom is isomorphic to \mathbb{R} .

2.8 PROPOSITION. Consider the open interval $S = (-3, 2]$ in \mathbb{R} . No minimum exists in S .

Proof. Assume $a = \min S$. Then $-3 < a \leq 2 \implies 0 < a + 3$. $1 < 1 + 1 = 2 \implies 1^{-1} > 2^{-1} \implies \frac{1}{2} < 1 \implies \frac{a+3}{2} < a + 3$. Let $b = a - \frac{a+3}{2} = \frac{a-3}{2}$. Then $-3 < b < a$. \square

TODO: Why does that last sentence hold?

2.9 EXAMPLE. Let $A = \{\frac{1}{n} : n \in \mathbb{N}\} \subseteq \mathbb{R}$. $\max A = 1$ and $\min A$ does not exist. The set of upper bounds $U = [1, +\infty)$. The set of lower bounds $L = (-\infty, 0]$. The last of these needs justification, which we’ll see later.

2.10 DEFINITION. If A admits no upper bound, we say A is **unbounded above** and if A admits no lower bound, we say A is **unbounded below**.

2.11 NOTATION. If $A \neq \emptyset$ and A is unbounded above, we write that $\sup A = +\infty$. This does not mean that $+\infty$ is a number nor that $\sup A$ exists. If $A \neq \emptyset$ is unbounded below, we write $\inf A = -\infty$. We also write that $\sup \emptyset = -\infty$ and $\inf \emptyset = +\infty$.

2.12 THEOREM (Archimedean Principle). \mathbb{N} is unbounded above in \mathbb{R} .

Proof. TODO \square

2.13 THEOREM. If $\sup A$ exists, it is unique. Same for $\inf A$.

Proof. Assume $s_1 = \sup A$ and $s_2 = \sup A$. Fix some $\varepsilon > 0, \varepsilon \in \mathbb{R}$. Then $s_2 - \varepsilon < s_2 \implies s_2 - \varepsilon$ is not an upper bound so there exists some element $x \in A$ such that $x > s_2 - \varepsilon \implies s_2 < x + \varepsilon \leq s_1 + \varepsilon$ because s_1 is an upper bound. Since $\forall \varepsilon > 0, s_2 < s_1 + \varepsilon, \forall \varepsilon > 0, s_2 - s_1 < \varepsilon \implies s_2 - s_1 \leq 0$. This argument is symmetric w.r.t. s_1 and s_2 , so we also have that $s_1 - s_2 \leq 0 \implies s_2 = s_1$. Thus $\sup A$ is unique. The same argument can be applied for $\inf A$. \square

TODO: Get that guy's name Here is an alternate, cleaner proof of the uniqueness of $\sup A$, courtesy of []:

Proof. \square

2.14 THEOREM (Existence of $\sqrt{2}$). *There exists $s_0 \in \mathbb{R}, s_0 > 0$ such that $s_0^2 = 2$.*

Proof. Let $A = \{x \in \mathbb{R} : x > 0, x^2 < 2\}$. The A is bounded above because 2 is an upper bound for A . (If $x \in A$ and $x > 2$, then $x^2 > 2 \cdot 2 = 2 + 2 > 2$. This is a contradiction because $x \in A$ implies $x^2 < 2$.) By the completeness axiom, this means that $\sup A$ must exist. Let $s_0 = \sup A$.

Since s_0 is an upper bound, $\forall x \in A, s_0 \geq x > 0 \implies \forall x \in A, s_0^2 \geq x^2$. Suppose $s_0^2 < 2$. Let $x = (s_0 + \varepsilon)$ for some $\varepsilon > 0$. Then $x^2 = s_0^2 + 2\varepsilon s_0 + \varepsilon^2 = s_0^2 + (2s_0 + \varepsilon)\varepsilon$. Choose $\varepsilon < \min\{1, \frac{2-s_0^2}{2s_0+1}\}$. Then $s_0^2 + (2s_0 + \varepsilon)\varepsilon < s_0^2 + (2s_0 + 1)\varepsilon \leq s_0^2 + (2s_0 + 1)\frac{2-s_0^2}{2s_0+1} = 2 \implies x^2 < 2 \implies x \in A$. But $x = s_0 + \varepsilon > s_0$, which is an upper bound for A . Contradiction. So $s_0^2 \not< 2 \implies s_0^2 \geq 2$.

Since s_0 is the smallest upper bound for A , $\forall \varepsilon > 0, s_0 - \varepsilon$ is not an upper bound and there exists some $x \in A$ such that $x > s_0 - \varepsilon$. $0 < s_0 < x + \varepsilon \implies s_0^2 < (x + \varepsilon)^2 = x^2 + 2\varepsilon x + \varepsilon^2 = x^2 + \varepsilon(2x + \varepsilon) < 2 + \varepsilon(4 + \varepsilon)$. So for all $0 < \varepsilon < 1, s_0^2 < 2 + 5\varepsilon$. Therefore $s_0^2 \leq 2 + 0 = 2$. \square