

Stage IV NSqNSCC, molecular tests negative (EGFR/ALK/ROS1/BRAF/RET/MET/EGFR ex20ins/KRAS G12C/NTRK/HER2), without contraindication for immunotherapy

Oligometastatic

not_Oligometastatic

Systemic therapy & LRT [II, B]

≥ 0 PS and ≤ 1 PS and any_expression_of_PD-L1

≥ 0 PS and ≤ 2 PS and ≥ 50 pdl1_percent

$= 2$ PS and < 50 pdl1_percent

Pembrolizumab-platinum-pemetrexed (4 cycles) followed by pembrolizumab-pemetrexed [I, A; MCBS 4]
Atezolizumab-carboplatin-nab-paclitaxel (4-6 cycles) followed by atezolizumab [I, A; MCBS 3]
Atezolizumab-bevacizumab-carboplatin-paclitaxel (4-6 cycles) followed by atezolizumab-bevacizumab [I, A; MCBS 3]
Nivolumab-ipilimumab + 2 cycles of platinum-doublet ChT followed by nivolumab-ipilimumab [I, A; MCBS 4]
Cemiplimab-platinum-doublet ChT (4 cycles) followed by cemiplimab + pemetrexed maintenance [I, A]
Durvalumab-tremelimumab-platinum-doublet ChT (4 cycles) followed by durvalumab-tremelimumab (tremelimumab one additional dose) + pemetrexed maintenance [I, A; MCBS 4]
Nivolumab-ipilimumab (only for PD-L1 $\geq 1\%$) [I, A; MCBS 4]

Disease progression

≥ 0 PS and ≤ 2 PS

≥ 3 PS and ≤ 4 PS

Pemetrexed [I, B]
Docetaxel [I, B]
Nintedanib-docetaxel [II, B]
Ramucirumab-docetaxel [I, B; MCBS 1]
Re-challenge ICI [III, B]

Best Supportive Care alone [III, A]

≥ 3 PS and ≤ 4 PS

Pembrolizumab [I, A; MCBS 5]
Atezolizumab (also for ICs $\geq 10\%$) [I, A; MCBS 5]
Cemiplimab [I, A; MCBS 4] (for PS 2 for all drugs: [III, B])

Disease progression

≥ 3 PS and ≤ 4 PS

≥ 0 PS and ≤ 2 PS

Platinum-doublet ChT [I, A] (pemetrexed preferred) [II, A; MCBS 4] followed by pemetrexed maintenance [I, A; MCBS 4] if pemetrexed switch maintenance [I, B]
If 4 cycles of gemcitabine-cisplatin: gemcitabine continuation maintenance [I, C]
Platinum-doublet ChT [I, A] (pemetrexed preferred) [II, A; MCBS 4]
Carboplatin-paclitaxel-bevacizumab followed by bevacizumab maintenance [I, A; MCBS 2] Or platinum-pemetrexed-bevacizumab followed by pemetrexed-bevacizumab maintenance [I, A]
Re-challenge ICI [III, B]

Pembrolizumab [I, A; MCBS 5]
Atezolizumab (also for ICs $\geq 10\%$) [I, A; MCBS 5]
Cemiplimab [I, A; MCBS 4] (for PS 2 for all drugs: [III, B])

Platinum-doublet ChT [carboplatin preferred: I, A; pemetrexed preferred: II, A]
Maintenance pemetrexed if improvement to PS 0-1 [MCBS 4]
Single-agent ChT [pemetrexed: II, B; gemcitabine, vinorelbine or docetaxel: I, B]

Disease progression

≥ 3 PS and ≤ 4 PS

≥ 0 PS and ≤ 2 PS

Nivolumab [I, A; MCBS 5]
Atezolizumab [I, A; MCBS 5]
Pembrolizumab (PD-L1 $\geq 1\%$) [I, A; MCBS 5]
Other options are the same as for second-line treatment for PS 0-2 after ChT-ICI