## Simple Plant Hydraulics model development and implementation

Daniel Kennedy - djk2120@columbia.edu Pierre Gentine - pg2328@columbia.edu

February 27, 2018

### 1 Model development

#### 1.1 Recipe

- 1. Solve for maximum stomatal conductance based on the Medlyn model (involves iterating for intercellular  $CO_2$ ).
- 2. Solve for the vegetation water potential (and associated stomatal conductance) that matches plant water supply with Penman-Monteith demand.
- 3. Calculate the photosynthesis based on the stomatal conductance solution (involves iterating for intercellular  $CO_2$ ).

#### 1.2 Plant Water Supply Equations

$$q = \int_{\psi_{soil}}^{\psi_{leaf}} \frac{K(\psi)}{z} d\psi \tag{1}$$

$$K(\psi) = \frac{\psi - p_2}{p_1 - p_2} \cdot K_{max} \tag{2}$$

$$q = \frac{K_{max}}{z} \cdot f(\psi) \cdot (\psi_{soil} - \psi_{leaf} - \rho gz)$$
(3)

$$f(\psi) = \frac{\frac{1}{2} (\psi_{soil} + \psi_{leaf}) - p_2}{p_1 - p_2}$$

$$0 \le f \le 1$$
(4)

#### 1.3 Plant Water Demand Equations

$$g_{c,max} = g_0 + \left(1 + \frac{g_1}{\sqrt{D}}\right) \frac{A}{C_a} \tag{5}$$

$$C_i = C_a - \frac{A}{g_c} \tag{6}$$

$$A = \frac{j/4 \left( C_i - \Gamma \right)}{C_i + 2\Gamma} \tag{7}$$

$$g_c = g_{c,max} \cdot h\left(\psi_{leaf}\right) \tag{8}$$

$$h\left(\psi_{leaf}\right) = \frac{1}{1 + \left(\frac{\psi_{leaf}}{p_{50}}\right)^{a}} \tag{9}$$

$$g_w = 1.6g_c \tag{10}$$

$$E = \frac{\frac{\Delta}{\gamma} \left( R_{net} - G \right) + \rho L_v g_a dq}{L_v \left( 1 + \frac{\Delta}{\gamma} + \frac{g_a}{g_w} \right)}$$
(11)

$$E = q \tag{12}$$

## 1.4 Bucket

$$\theta_1 = \theta_0 - \frac{q\Delta t}{Z_r} \tag{13}$$

$$\psi_{soil}\left(\theta\right) = \psi_{soil,sat}\left(\frac{\theta}{\theta_{sat}}\right)^{-b}$$
(14)

# 2 Why are tall Amazonian forests more resistant to precipitation variability?

#### 2.1 Thread 1

Assume  $\frac{dk}{d\psi} = 0$ 

$$q = k\Delta\psi \tag{15}$$

$$\Delta \psi = \psi_{soil} - \psi_{leaf} \tag{16}$$

$$q = k \left( \psi_{soil} - \psi_{leaf} \right) \tag{17}$$

$$\frac{dq}{d\psi_{soil}} = k \left( 1 - \frac{d\psi_{leaf}}{d\psi_{soil}} \right) \tag{18}$$

$$\sigma = \frac{d\psi_{leaf}}{d\psi_{soil}} \tag{19}$$

$$\frac{dq}{d\psi_{soil}} = k \left( 1 - \sigma \right) \tag{20}$$

Use a reference pressure drop and water flux (e.g. Day 1 of drydown) to infer k:

$$k = \frac{q}{\Delta \psi}$$

$$k = \frac{q_0}{\psi_{soil,0} - \psi_{leaf,0}}$$
(21)

Then we have another way to express  $\frac{dq}{d\psi}$ 

$$\frac{dq}{d\psi_{soil}} = q_0 \frac{(1-\sigma)}{\psi_{soil,0} - \psi_{leaf,0}} \tag{22}$$

This shows that the loss of water supply depends is related to the absolute potential drop from soil-to-root. For taller trees, the potential drop from soil-to-root tends to be larger, whereby a given drop in soil potential due to drydown would have a smaller effect as compared to shorter trees.

Likewise the effect of transpiration demand will be larger for taller trees, because for a given increase in transpiration demand (dq), the effect on the potential drop  $(\Delta \psi)$  will be larger, due to the smaller conductance.

$$\Delta \psi = \frac{q}{k} \tag{23}$$

$$\frac{d\Delta\psi}{dq} = \frac{1}{k} = \frac{\psi_{soil,0} - \psi_{leaf,0}}{q_0} \tag{24}$$

## 2.2 Thread 2

Given the same precipitation sequence, soil potential variability is smaller for taller trees due to deeper roots,