



# FEEDING RATE DIBAQ MICROBAQ T

TILAPIA'S STARTER STRATEGY.

| Weight (grams)   | <0,1                                                      | 0.1 to 0.5 | 0.5 to 1.0   | 1.0 to 4.0   | 4.0 to 10    |
|------------------|-----------------------------------------------------------|------------|--------------|--------------|--------------|
| Caliber (mm)     | Microbaq 0                                                | Microbaq 1 | Microbaq 100 | Microbaq 145 | Microbaq 165 |
| Temperature (°C) | Daily food consumption per 100 kilograms of fish biomass. |            |              |              |              |
| 22 °C            | 4,6                                                       | 4,3        | 3,8          | 3,0          | 2,7          |
| 23 °C            | 5,2                                                       | 4,9        | 4,3          | 3,3          | 3,1          |
| 24 °C            | 5,9                                                       | 5,5        | 4,8          | 3,7          | 3,4          |
| 25 °C            | 6,6                                                       | 6,2        | 5,4          | 4,2          | 3,9          |
| 26 °C            | 7,4                                                       | 7,0        | 6,1          | 4,7          | 4,3          |
| 27 °C            | 8,4                                                       | 7,8        | 6,8          | 5,3          | 4,9          |
| 28 °C            | 9,4                                                       | 8,8        | 7,7          | 6,0          | 5,5          |
| 29 °C            | 10,5                                                      | 9,9        | 8,6          | 6,7          | 6,2          |

This table displays the daily food quantity per 100 kg of biomass, considering the nutritional characteristics of the food and the need for customization to suit individual fish farm conditions.



# FEEDING RATE DIBAQ TILAPIA

## TILAPIA'S PRODUCTION STRATEGY

| Weight (grams)   | 10 a 50                                                   | 50 a 100 | 100 a 250 | 250 a 500 | >500 |
|------------------|-----------------------------------------------------------|----------|-----------|-----------|------|
| Caliber (mm)     | 2                                                         | 2        | 3,5       | 3,5       | 5    |
| Temperature (°C) | Daily food consumption per 100 kilograms of fish biomass. |          |           |           |      |
| < 10 °C          | According to the fish's appetite                          |          |           |           |      |
| 10 °C            | 1,2                                                       | 1,0      | 0,8       | 0,6       | 0,5  |
| 12 °C            | 1,8                                                       | 1,5      | 1,0       | 0,8       | 0,6  |
| 14 °C            | 2,4                                                       | 2,0      | 1,5       | 1,0       | 0,8  |
| 16 °C            | 3,1                                                       | 2,6      | 1,8       | 1,3       | 1,0  |
| 18 °C            | 3,7                                                       | 3,1      | 2,3       | 1,5       | 1,3  |
| 20 °C            | 4,3                                                       | 3,6      | 2,8       | 1,8       | 1,5  |
| 22 °C            | 4,9                                                       | 4,1      | 3,3       | 2,3       | 1,8  |
| 24 °C            | 5,1                                                       | 4,6      | 3,8       | 2,8       | 2,0  |
| 26 °C            | 5,3                                                       | 5,0      | 4,1       | 3,0       | 2,1  |
| 28 °C            | 5,5                                                       | 5,2      | 4,3       | 3,1       | 2,2  |
| 30 °C            | 6,2                                                       | 5,8      | 4,8       | 3,5       | 2,5  |
| 32 °C            | 6,9                                                       | 6,5      | 5,4       | 3,9       | 2,8  |
| <33 °C           | According to the fish's appetite or O2 level              |          |           |           |      |

This table displays the daily food quantity per 100 kg of biomass, considering the nutritional characteristics of the food and the need for customization to suit individual fish farm conditions.