

# Transport formulas

Sean Ballinger

May 13, 2017

## 1 Confinement time

For a scaling relation of the form

$$\tau_E = C I^{\alpha_I} B^{\alpha_B} \bar{n}^{\alpha_n} P^{\alpha_P} R^{\alpha_R} \kappa^{\alpha_\kappa} \epsilon^{\alpha_\epsilon} S_{cr}^{\alpha_S} M^{\alpha_M} [\text{sec}]$$

we have the following popular scaling laws:

Scaling	$C \times 10^3$	$\alpha_I$	$\alpha_B$	$\alpha_n$	$\alpha_P$	$\alpha_R$	$\alpha_\kappa$	$\alpha_\epsilon$	$\alpha_S$	$\alpha_M$
ITERH-98P(y,2)	56.2	0.93	0.15	0.41	-0.69	1.97	0.78	0.58	-	0.19
ITPAH-04P(y,1)	22.8	0.86	0.21	0.40	-0.65	0.32	-	-0.99	0.84	0.08
ITPAH-04P(y,2)	19.8	0.85	0.17	0.26	-0.45	-0.04	-	-1.25	0.82	0.11
ITPAH-04P(y,3)	88.0	0.90	-	0.30	-0.47	1.73	-	0.43	-	-
ITER-89P	38	0.85	0.20	0.10	-0.50	1.50	0.50	0.30	-	0.50
ITERL-96P(th)	23	0.96	0.03	0.40	-0.73	1.83	0.64	-0.06	-	0.20

ITER98y,2 is the most popular choice.

### 1.1 I-mode

Using  $\alpha_R$  from the ITER98y,2 scaling relation, we have that

$$\tau_E = 0.082 \frac{I_M^{0.69} B_0^{0.77} \bar{n}_{20}^{0.017} a^{0.81} R_0^{1.39}}{P_M^{0.29}} [\text{sec}]$$

Source: J.P. Freidberg's re-derivation of a size scaling for I-mode using the Kadomtsev constraint. I-mode has not been replicated on enough machines to have a robust empirical size scaling.

## 2 H-mode power threshold

The L-H transition can only occur if the heating power is greater than the threshold value

$$P_{LH} = 1.38 \bar{n}_{20}^{-0.77} B_0^{0.92} R_0^{1.23} a^{0.76} [\text{MW}]$$

Source: J.P. Freidberg, *Plasma Physics and Fusion Energy* sec. 14.5.2.

### 3 Density limit

Operating above the Greenwald density results in disruptions. This limit is defined as

$$n_G = \frac{I_p}{\pi a^2} [10^{20}/\text{m}^3]$$

Arbitrary limits on the achievable fraction of this density can be found by looking at the database of H-mode shots. A reasonable choice for power plant operation is

$$0.35 < \frac{n}{n_G} < 0.7$$

where the lower limit arises from difficulties in entering H-mode.

### 4 Edge pedestal scaling

$$p_{ped} = 18.3a^{-0.57} \left( \frac{M}{n_{ped}} \right) \left( \frac{I_p}{R} \right)^2 \left( \frac{P_{total}}{P_{LH}} \right)^{0.144} [\text{kPa}]$$

Source: Sugihara, IOP 2003.

### 5 Simple model for $n$ and $T$ profiles in H-mode

$$\begin{aligned} n &= \frac{4}{3} \bar{n} (1 - \rho^2)^{1/3} \\ T &= \frac{5}{3} \bar{T} (1 - \rho^2)^{2/3} \\ \rho^2 &= \frac{x^2}{a^2} + \frac{y^2}{\kappa^2 a^2} \end{aligned}$$

This gives elliptical contours using the maximum, on-axis values of  $\bar{T}$  and  $\bar{n}$ .

Source: J.P. Freidberg, *Plasma Physics and Fusion Energy*.