

Q26 Criteria for admission to hospital post ER

Answered: 12 Skipped: 0

#	RESPONSES	DATE
1	-Unresolving pain, significant bleeding during/post procedure, perforation where soiling possible or confidence of closure lacking, high co-morbidity	1/13/2018 6:40 AM
2	Fever bleeding abdominal pain	12/7/2017 5:48 PM
3	- Any abdominal pain after discharge - Dark stool - Bright red bleeding - Patient feeling faint - Hypotension/ tachycardia/increase in temp noted by a medical practitioner	12/3/2017 9:54 PM
4	* hemodynamic instable patient * signs of ongoing bleeding	11/18/2017 9:37 PM
5	Haevy bleeding Severe or worsening pain Fever Tachycardia Distension Focal or generalized peritonism/tenderness	11/14/2017 6:08 AM
6	1. Patient that lives home alone without supports 2. Significant intra-procedural bleeding needing overnight observation 3. Intra-procedural perforation managed with endoscopic closure 4. Patients needing early reintroduction of anticoagulation (e.g MVR)	11/13/2017 1:44 PM
7	Significant abdominal pain post procedure Intraprocedural complications of major bleeding or perforation that could or could not be managed endoscopically Prophylactic admission post resection of large complex colonic lesion especially in patients living far away from the tertiary care center	11/5/2017 3:14 PM
8	1. On going abdominal discomfort 2. Peritonism	11/5/2017 3:26 AM
9	- instability - blood in your stools every 30 minutes or less - faint - fever with chills - stroke or other - abdominal pain + examination suggestive of perforation	11/3/2017 3:11 PM
10	frequent bloody stool moderate or severe pain not responding to simply analgesia pain increasing in intensity rigours, sustained high temperature abdominal signs of peritonism frequent nausea/vomiting	11/3/2017 12:46 PM
11	Severe pain, significant rectal bleeding, fever	11/1/2017 11:43 AM
12	Untreated perforation. Perforation in poor prep. Pain that requires opiates. Peritonism. Dramatic intra-procedural bleeding likely to need transfusion. Sedation or comorbid disease related complication requiring monitoring. At home alone after sedation.	10/31/2017 11:04 PM