

MiMeS: Misalignment Mechanism Solver

Dimitrios Karamitros

Manchester U.

23/11/2021

El Journal Club más Sabroso

Outline

- 1 **Axion Dark Matter**
 - Why particle dark matter
 - The dark matter particle
 - The axion (like) particle
- 2 **Calculating the Relic Abundance**
 - The axion EOM
 - How hard can it be?
 - Initial conditions
 - (Bad) Analytical approximations
 - Need for accuracy, speed, and automation
- 3 **MiMeS**
 - MiMeS: General
 - MiMeS: Under the hood
 - MiMeS: Notation
 - MiMeS: When do you start and stop integrating?
 - MiMeS: Some (classical) physics
- 4 **Using MiMeS**
 - How to get `MiMeS`
 - Configure (and make)
 - Classes
 - Assumptions
 - What `MiMeS` expects from you
 - Template arguments
 - `MiMeS` from `python`
- 5 **Examples**
 - `C++`
 - `python`
- 6 **Outlook**
 - Summing up
 - Future

Axion Dark Matter

1

Axion Dark Matter

- Why particle dark matter
- The dark matter particle
- The axion (like) particle

2

Calculating the Relic Abundance

- The axion EOM
- How hard can it be?
- Initial conditions
- (Bad) Analytical approximations
- Need for accuracy, speed, and automation

3

MiMeS

- MiMeS: General
- MiMeS: Under the hood
- MiMeS: Notation
- MiMeS: When do you start and stop integrating?
- MiMeS: Some (classical) physics

4

Using MiMeS

- How to get MiMeS
- Configure (and make)
- Classes
- Assumptions
- What MiMeS expects from you
- Template arguments
- MiMeS from python

5

Examples


- C++
- python

6

Outlook

- Summing up
- Future


Why particle dark matter



E. Corbelli and P. Salucci, *Mon. Not. Roy. Astron. Soc.*
311 441 (2000), [arXiv:astro-ph/9909252](#).



M. Markevitch, *ESA Spec. Publ.* **604** (2006) 723,
[astro-ph/0511345](#). Clowe, Bradac, *et. al.* *Astrophys. J.*
648. L109 (2006). [astro-ph/0608407](#)



N. Aghanim *et al.* [Planck Collaboration], [arXiv:1807.06209 \[astro-ph.CO\]](#).

“Εν οἶδα, ὅτι οὐδὲν οἶδα.”

“I know one thing, that I know nothing.”

—Socrates

The dark matter particle

“Εν οἶδα, ὅτι οὐδέν οἶδα.”

“I know one thing, that I know nothing.”

—Socrates

- Gravitational interactions.
- Mostly electrically neutral.
- Stable or very slow decay rate.
- Non-Baryonic.
- Cold/Warm and non-relativistic today.

The axion (like) particle

Notably, the original Axion was originally introduced in order to solve the *strong-CP problem* of the SM. Axion-Like-Particles (ALPs) arise in a number of new physics models, beyond the SM.

Axions and ALPs generally:

- Have suppressed interactions with photons.
- Are (mostly) stable.
- Were non-relativistic around the epoch of structure formation.

The axion (like) particle

Notably, the original Axion was originally introduced in order to solve the *strong-CP problem* of the SM. Axion-Like-Particles (ALPs) arise in a number of new physics models, beyond the SM.

Axions and ALPs generally:

- Have suppressed interactions with photons.
- Are (mostly) stable.
- Were non-relativistic around the epoch of structure formation.
- Non-baryonic (by definition), and of-course interact gravitationally.

The axion (like) particle

Notably, the original Axion was originally introduced in order to solve the *strong-CP problem* of the SM. Axion-Like-Particles (ALPs) arise in a number of new physics models, beyond the SM.

Axions and ALPs generally:

- Have suppressed interactions with photons.
- Are (mostly) stable.
- Were non-relativistic around the epoch of structure formation.
- Non-baryonic (by definition), and of-course interact gravitationally.

Maybe DM has Axionic nature!

Calculating the Relic Abundance

1

Axion Dark Matter

- Why particle dark matter
- The dark matter particle
- The axion (like) particle

2

Calculating the Relic Abundance

- The axion EOM
- How hard can it be?
- Initial conditions
- (Bad) Analytical approximations
- Need for accuracy, speed, and automation

3

MiMeS

- MiMeS: General
- MiMeS: Under the hood
- MiMeS: Notation
- MiMeS: When do you start and stop integrating?
- MiMeS: Some (classical) physics

4

Using MiMeS

- How to get MiMeS
- Configure (and make)
- Classes
- Assumptions
- What MiMeS expects from you
- Template arguments
- MiMeS from python

5

Examples

- C++
- python

6

Outlook

- Summing up
- Future

Axions and ALPs follow a similar equation of motion (EOM):

$$\left(\frac{d^2}{dt^2} + 3H(t) \frac{d}{dt} \right) \theta(t) + \tilde{m}_a^2(t) \sin \theta(t) = 0 ,$$

where $\theta = A f_a$, with A the axion field, and f_a some energy scale that characterises the potential (Peccei-Quinn breaking scale).

How hard can it be?

Hard (in general).

How hard can it be?

Hard (in general).

The classical analogue is the damped pendulum with both frequency (length) and friction being time-dependent:

- There is no closed form solution.
- There are no constants of motion (wait a minute).
- No package/library/program available!

How hard can it be?

Hard (in general).

The classical analogue is the damped pendulum with both frequency (length) and friction being time-dependent:

- There is no closed form solution.
- There are no constants of motion (wait a minute).
- No package/library/program available!

MiMeS simulates the evolution of the Axion/ALP, for (virtually) any cosmological scenario and Axion/ALP (thermal) mass.

Some time at the very early Universe, $\tilde{m}_a \ll H(T)$,¹ with

$$\ddot{\theta} + 3H \dot{\theta} \approx 0.$$

The solution is

$$\theta = \theta_{\text{ini}} + C \int_0^t dt' \left(\frac{a(t' = 0)}{a(t')} \right)^3.$$

So, $\dot{\theta} \sim a^{-3}$. Since we are interested in θ at much later times (once the potential becomes relevant), $\dot{\theta} \approx 0$.² Therefore, we can start integration at some point ($t = t_{\text{ini}}$) with $3H \gg \tilde{m}_a$, and set $\theta(t = t_{\text{ini}}) = \theta_{\text{ini}}$ and $\dot{\theta}(t = t_{\text{ini}}) = 0$.

¹ This is an assumption that MiMeS has to make, for the sake of generality.

² Standard misalignment mechanism. For the kinetic one see [R. T. Co, L. J. Hall and K. Harigaya, Phys. Rev. Lett. **124** \(2020\) no.25, 251802 \[arXiv:1910.14152 \[hep-ph\]\]](#), [C. F. Chang and Y. Cui, Phys. Rev. D **102** \(2020\) no.1, 015003 \[arXiv:1911.11885 \[hep-ph\]\]](#), or [B. Barman, N. Bernal, N. Ramberg and L. Visinelli, \[arXiv:2111.03677 \[hep-ph\]\]](#).

(Bad) Analytical approximations

Once we agree on the initial conditions, we move to the next important things:

³ Defined from $s(T_0) = \gamma a_{\text{osc}}^3 s_{\text{osc}}$.

(Bad) Analytical approximations

Once we agree on the initial conditions, we move to the next important things:

- Assume $\theta \ll 1$, and linearise the EOM. *Not always that bad.*

³ Defined from $s(T_0) = \gamma a_{\text{osc}}^3 s_{\text{osc}}$.

(Bad) Analytical approximations

Once we agree on the initial conditions, we move to the next important things:

- Assume $\theta \ll 1$, and linearise the EOM. *Not always that bad.*
- Assume that at $\tilde{m}_a(T_{\text{osc}}) \approx 3H(T_{\text{osc}})$, the oscillations begin with $\dot{\theta}(T_{\text{osc}}) = 0$. *This is in general quite bad.*

³ Defined from $s(T_0) = \gamma a_{\text{osc}}^3 s_{\text{osc}}$.

(Bad) Analytical approximations

Once we agree on the initial conditions, we move to the next important things:

- Assume $\theta \ll 1$, and linearise the EOM. *Not always that bad.*
- Assume that at $\tilde{m}_a(T_{\text{osc}}) \approx 3H(T_{\text{osc}})$, the oscillations begin with $\dot{\theta}(T_{\text{osc}}) = 0$. *This is in general quite bad.*
- Assume that $\theta_{\text{osc}} \approx \theta_{\text{ini}}$. *Generally quite bad.*

³ Defined from $s(T_0) = \gamma a_{\text{osc}}^3 s_{\text{osc}}$.

(Bad) Analytical approximations

Once we agree on the initial conditions, we move to the next important things:

- Assume $\theta \ll 1$, and linearise the EOM. *Not always that bad.*
- Assume that at $\tilde{m}_a(T_{\text{osc}}) \approx 3H(T_{\text{osc}})$, the oscillations begin with $\dot{\theta}(T_{\text{osc}}) = 0$. *This is in general quite bad.*
- Assume that $\theta_{\text{osc}} \approx \theta_{\text{ini}}$. *Generally quite bad.*

Then, we get the “WKB”-approximate solution

$$\theta(t) \approx \theta_{\text{ini}} \left(\frac{3}{4} \right)^{1/4} \sqrt{\frac{\tilde{m}_a(T_{\text{osc}})}{\tilde{m}_a(T)}} \left(\frac{a}{a_{\text{osc}}} \right)^{-3/2} \cos \left(\int_{t_{\text{osc}}}^t dt' \tilde{m}_a(t') \right) .$$

The advantage of this approximation is that we get an easy formula for the axion/ALP energy density today:

$$\rho_{a,0} = \gamma^{-1} \frac{s_0}{s_{\text{osc}}} \frac{1}{2} f_a^2 m_a \tilde{m}_{a,\text{osc}} \theta_{\text{ini}}^2 ,$$

where γ the amount of entropy injection between T_{osc} and today. ³

³ Defined from $s(T_0) = \gamma a_{\text{osc}}^3 s_{\text{osc}}$.

Need for accuracy, speed, and automation

Serious disadvantages of the approximate results:

- The approximations can be tested against numerical results in a case-by-case basis; there is no way to tell if they will work in new models and cosmological scenarios.
- There is no available tool that can help us reproduce published results obtained by numerical integration; people use their own **private** code.
- If someone wants to simply see if an ALP model is compatible with a cosmological scenario, they have to develop their own private code; the overall effort of the community increases.

MiMeS:

- *Easy* to use; anyone can run it and see if their model can work.
- Reasonably fast; less than 0.05 s for the scenarios tested.
- Tools that can help determine if the algorithm is accurate enough.
- The user provides too much input. This helps the user determine whether convergence of the algorithm is consistent.

1

Axion Dark Matter

- Why particle dark matter
- The dark matter particle
- The axion (like) particle

2

Calculating the Relic Abundance

- The axion EOM
- How hard can it be?
- Initial conditions
- (Bad) Analytical approximations
- Need for accuracy, speed, and automation

3

MiMeS

- MiMeS: General
- MiMeS: Under the hood
- MiMeS: Notation
- MiMeS: When do you start and stop integrating?
- MiMeS: Some (classical) physics

4

Using MiMeS

- How to get MiMeS
- Configure (and make)
- Classes
- Assumptions
- What MiMeS expects from you
- Template arguments
- MiMeS from python

5

Examples

- C++
- python

6

Outlook

- Summing up
- Future

- MiMeS is a C++ header-only library that contains various templated classes; there is no “installation” and no special procedures, just include the header files.

- MiMeS is a C++ header-only library that contains various templated classes; there is no “installation” and no special procedures, just include the header files.
- MiMeS comes with a `python` interface so that everybody can use it.

- MiMeS is a C++ header-only library that contains various templated classes; there is no “installation” and no special procedures, just include the header files.
- MiMeS comes with a `python` interface so that everybody can use it.
- MiMeS is distributed under the MIT license; you can do whatever you want with it, and I am *not* responsible.

- MiMeS is a C++ header-only library that contains various templated classes; there is no “installation” and no special procedures, just include the header files.
- MiMeS comes with a `python` interface so that everybody can use it.
- MiMeS is distributed under the MIT license; you can do whatever you want with it, and I am *not* responsible.

MiMeS relies on `NaBBODES`⁴ for the numerical integration, and `SimpleSplines`⁵ for the various interpolations.

Advantages:

- You only need to have the standard C++ library.
- The two libraries are developed by myself, so their integration with MiMeS is seamless.
- There is always going to be a compatible version of these libraries that works with MiMeS.

Disadvantages:

- These are not well tested libraries.
- No community of contributors; if it doesn't work, I have to fix it.
- Slow development.

⁴ <https://github.com/dkaramit/NaBBODES>.

⁵ <https://github.com/dkaramit/SimpleSplines>.

MiMeS uses a notation suitable (any) underlying cosmology, since it is up to the user to define the cosmological evolution. First, we define

$$u \equiv \log(a/a_{\text{ini}}) ,$$

with a_{ini} some initial value of the scale factor.⁶ in order to express the time derivatives as

$$\frac{d}{dt} \rightarrow H \frac{d}{du} , \quad \frac{d^2}{dt^2} \rightarrow H^2 \left(\frac{d^2}{du^2} + \frac{1}{2} \frac{d \log H^2}{du} \frac{d}{du} \right) .$$

Then, we express the EOM as a system of first order ordinary differential equations

$$\frac{d\zeta}{du} + \left[\frac{1}{2} \frac{d \log H^2}{du} + 3 \right] \zeta + \left(\frac{\tilde{m}_a}{H} \right)^2 \sin \theta = 0 .$$
$$\frac{d\theta}{du} - \zeta = 0 .$$

Observe that, by definition, $\zeta = d\theta/du$. The initial conditions are $\zeta(0) = 0$ and $\theta(0) = \theta_{\text{ini}}$.

⁶ Only the ratios a/a_{ini} appear in the calculations. So, the choice does not matter as long as it is consistent.

MiMeS: When do you start and stop integrating?

The choice of a good starting point is important, as need to start at a temperature where $\zeta = 0$ is a good approximation. So you can start at some T_{ini} with a given ratio $3H(T_{\text{ini}})/\tilde{m}_a(T_{\text{ini}}) \gg 1$. This needs to be chosen carefully, as low values of $3H(T_{\text{ini}})/\tilde{m}_a(T_{\text{ini}})$ result in inaccurate result, while high values may result in a slow calculation. *Advice:* Use various values of that ratio, and find where the relic abundance becomes T_{ini} -independent.

The stopping condition is more difficult. You should stop at some point where the axion/ALP evolves "adiabatically". Find a quantity that becomes constant as the system relaxes, and use this to determine when adiabaticity is reached. Once θ starts to evolve adiabatically, the amplitude of its oscillation is known at later times!

If a system exhibits closed orbits, the quantity

$$J \equiv C \oint p \, d\theta \, ,$$

is the adiabatic invariant. In this case, it becomes

$$J = a^3 \, \tilde{m}_a \, \theta_{\text{peak}}^2 \, f(\theta_{\text{peak}}) \, ,$$

with

$$f(\theta_{\text{peak}}) = \frac{2\sqrt{2}}{\pi \theta_{\text{peak}}^2} \int_{-\theta_{\text{peak}}}^{\theta_{\text{peak}}} d\theta \sqrt{\cos \theta - \cos \theta_{\text{peak}}} \, ,$$

the so-called anharmonic factor.

MiMeS: Some (classical) physics

If a system exhibits closed orbits, the quantity

$$J \equiv C \oint p \, d\theta \, ,$$

is the adiabatic invariant. In this case, it becomes

$$J = a^3 \, \tilde{m}_a \, \theta_{\text{peak}}^2 \, f(\theta_{\text{peak}}) \, ,$$

with

$$f(\theta_{\text{peak}}) = \frac{2\sqrt{2}}{\pi \theta_{\text{peak}}^2} \int_{-\theta_{\text{peak}}}^{\theta_{\text{peak}}} d\theta \sqrt{\cos \theta - \cos \theta_{\text{peak}}} \, ,$$

the so-called anharmonic factor.

Important: θ_{peak} is the peak of the oscillation. So, J can be used to determine how θ_{peak} changes with time. By definition, at $\theta = \theta_{\text{peak}}$, $p \sim \dot{\theta} = 0$. This means that we can find $\rho_{a,0}$ on the peak of today's θ , as

$$\rho_{a,0} = \gamma^{-1} \frac{s_0}{s_*} m_a \, \tilde{m}_{a,*} \frac{1}{2} f_a^2 \theta_{\text{peak},*}^2 \, f(\theta_{\text{peak},*}) \, ,$$

where T_* the temperature at which adiabaticity was reached, and γ the entropy injection between T_* and today (*i.e.* $s_0 = \gamma a_*^3 s_*$).

Using MiMeS

1

Axion Dark Matter

- Why particle dark matter
- The dark matter particle
- The axion (like) particle

2

Calculating the Relic Abundance

- The axion EOM
- How hard can it be?
- Initial conditions
- (Bad) Analytical approximations
- Need for accuracy, speed, and automation

3

MiMeS

- MiMeS: General
- MiMeS: Under the hood
- MiMeS: Notation
- MiMeS: When do you start and stop integrating?
- MiMeS: Some (classical) physics

4

Using MiMeS

- How to get MiMeS
- Configure (and make)
- Classes
- Assumptions
- What MiMeS expects from you
- Template arguments
- MiMeS from python

5

Examples

- C++
- python

6

Outlook

- Summing up
- Future

How to get MiMeS

There are several ways you can get a stable version of MiMeS:

1

`git clone -b stable https://github.com/dkaramit/MiMeS.git`.
This is the preferred way, as it is guaranteed to be the latest stable version.

2

Go to mimes.hepforge.org/downloads, and download it.

3

Go to github.com/dkaramit/MiMeS/releases, and download a released version.

You can get the most up-to-date code – not always the most stable one – including the latest version of NaBBODES and SimpleSplines, by running

```
1  git clone https://github.com/dkaramit/MiMeS.git
2  cd MiMeS
3  git submodule init
4  git submodule update --remote
```

Configure (and make)

There is no need to install anything if you are going to use MiMeS in a C++ program. The only thing you *must* do is run

```
1    bash configure.sh
```

Alter that, you can include the header file `MiMeS/MiMeS.hpp`, and you are good to go.

However, you can also run

- `make lib`, in order to produce the (shared) libraries. This is needed in order to run the `python` interface.
- `make examples`, in order to compile the examples in `MiMeS/UserSpace/Cpp`.
- `make exec`, in order to produce some test executables (in `MiMeS/exec`). You just need to run then in order to see if you get any segfaults.

There are three classes useful to the user.⁷

- `mimes::Cosmo<LD>`: interpolation of relativistic degrees of freedom of the plasma. By default it uses the EOS2020⁸ data. The user can choose another file easily.

⁷ There are various arguments that need to be passed to the constructors, and they are all listed and explained in the Appendix of the documentation.

⁸ K. Saikawa and S. Shirai, JCAP **08** (2020), 011 [arXiv:2005.03544 [hep-ph]].

⁹ S. Borsanyi, Z. Fodor, J. Guenther, K. H. Kampert, S. D. Katz, T. Kawanai, T. G. Kovacs, S. W. Mages, A. Pasztor and F. Pittler, *et al.* Nature **539** (2016) no.7627, 69-71 [arXiv:1606.07494 [hep-lat]].

There are three classes useful to the user.⁷

- `mimes::Cosmo<LD>`: interpolation of relativistic degrees of freedom of the plasma. By default it uses the EOS2020⁸ data. The user can choose another file easily.
- `mimes::AxionMass<LD>`: definition of axion/ALP mass as a function of the temperature and f_a . MiMeS is shipped with data from Lattice calculation⁹ of the QCD axion mass.

⁷ There are various arguments that need to be passed to the constructors, and they are all listed and explained in the Appendix of the documentation.

⁸ K. Saikawa and S. Shirai, JCAP **08** (2020), 011 [arXiv:2005.03544 [hep-ph]].

⁹ S. Borsanyi, Z. Fodor, J. Guenther, K. H. Kampert, S. D. Katz, T. Kawanai, T. G. Kovacs, S. W. Mages, A. Pasztor and F. Pittler, *et al.* Nature **539** (2016) no.7627, 69-71 [arXiv:1606.07494 [hep-lat]].

There are three classes useful to the user.⁷

- `mimes::Cosmo<LD>`: interpolation of relativistic degrees of freedom of the plasma. By default it uses the EOS2020⁸ data. The user can choose another file easily.
- `mimes::AxionMass<LD>`: definition of axion/ALP mass as a function of the temperature and f_a . MiMeS is shipped with data from Lattice calculation⁹ of the QCD axion mass.
- `mimes::Axion<LD, Solver, Method>`: This is responsible for actually solving the EOM.

⁷ There are various arguments that need to be passed to the constructors, and they are all listed and explained in the Appendix of the documentation.

⁸ K. Saikawa and S. Shirai, JCAP **08** (2020), 011 [arXiv:2005.03544 [hep-ph]].

⁹ S. Borsanyi, Z. Fodor, J. Guenther, K. H. Kampert, S. D. Katz, T. Kawanai, T. G. Kovacs, S. W. Mages, A. Pasztor and F. Pittler, *et al.* Nature **539** (2016) no.7627, 69-71 [arXiv:1606.07494 [hep-lat]].

Assumptions

MiMeS is designed to make as few assumption as possible. However, it still assumes that:

Assumptions

MiMeS is designed to make as few assumption as possible. However, it still assumes that:

- 1 H/\tilde{m}_a increases monotonically with the temperature.

Assumptions

MiMeS is designed to make as few assumption as possible. However, it still assumes that:

- 1 H/\tilde{m}_a increases monotonically with the temperature.
- 2 $\zeta(0) = 0$. This will be changed in the future.

MiMeS is designed to make as few assumption as possible. However, it still assumes that:

- 1 H/\tilde{m}_a increases monotonically with the temperature.
- 2 $\zeta(0) = 0$. This will be changed in the future.
- 3 The energy density of the axion/ALP is always subdominant.

MiMeS is designed to make as few assumption as possible. However, it still assumes that:

- 1 H/\tilde{m}_a increases monotonically with the temperature.
- 2 $\zeta(0) = 0$. This will be changed in the future.
- 3 The energy density of the axion/ALP is always subdominant.
- 4 Only the EOM determines the energy density (no annihilations, no strings, etc.).

What MiMeS expects from you

Apart from θ_{ini} and f_a , keep in mind that MiMeS needs:

- 1 The mass of the axion/ALP. A data file or an actual function.
- 2 Data file with $\log a/a_i$ (a_i is some arbitrary value; MiMeS rescales it appropriately), T , and $\log H$ of the underlying cosmology.

What MiMeS expects from you

Apart from θ_{ini} and f_a , keep in mind that MiMeS needs:

- 1 The mass of the axion/ALP. A data file or an actual function.
- 2 Data file with $\log a/a_i$ (a_i is some arbitrary value; MiMeS rescales it appropriately), T , and $\log H$ of the underlying cosmology.
- 3 Value for $3H/\tilde{m}_a \gg 1$, which defines the point where integration begins.

What MiMeS expects from you

Apart from θ_{ini} and f_a , keep in mind that MiMeS needs:

- 1 The mass of the axion/ALP. A data file or an actual function.
- 2 Data file with $\log a/a_i$ (a_i is some arbitrary value; MiMeS rescales it appropriately), T , and $\log H$ of the underlying cosmology.
- 3 Value for $3H/\tilde{m}_a \gg 1$, which defines the point where integration begins.
- 4 Relative difference of J between a given number of peaks at which we consider adiabaticity to have been reached.

What MiMeS expects from you

Apart from θ_{ini} and f_a , keep in mind that MiMeS needs:

- 1 The mass of the axion/ALP. A data file or an actual function.
- 2 Data file with $\log a/a_i$ (a_i is some arbitrary value; MiMeS rescales it appropriately), T , and $\log H$ of the underlying cosmology.
- 3 Value for $3H/\tilde{m}_a \gg 1$, which defines the point where integration begins.
- 4 Relative difference of J between a given number of peaks at which we consider adiabaticity to have been reached.
- 5 Other input, related to the algorithm, that might confuse you; e.g. temperature at which integration stops *no matter what!*

Template arguments

You need to choose what numeric type to use. This is done by the template argument `LD` which should be `double` (fast) or `long double` (accurate).¹⁰

You also need to tell `MiMeS` which integration strategy to use. This is done by choosing template arguments:

- `Solver` can be set to 1 for Rosenbrock (semi-implicit Runge-Kutta). The `Method` argument in this case can be:
 - `RODASPR2<LD>` (4th order).
 - `ROS34PW2<LD>` (3rd order).
 - `ROS3W<LD>` (2rd order, *very bad*).
- `Solver` can be set to 2 for explicit RK. The `Method` argument can be:
 - `DormandPrinceRK45<LD>` (7th order)
 - `CashKarpRK45<LD>` (5th order, *very bad*).
 - `RK45<LD>` (5th order, *very bad*).

¹⁰ You could choose `float`, but we live in 2021.

In order to call the `python` interface of MiMeS, we need to first call `make lib` in the root directory of MiMeS.

Before that, we can take some time to decide what the template arguments and compilation options should be. In the file `MiMeS/Definitions.mk`, you can change the variables:

- `LONG=long` will compile the library with `long double` numeric types. `LONG=` will compile the library with `double` numeric types.
- `SOLVER` and `METHOD`, as in the template arguments.

Also, in the same file, you can change compilation options:

- **Compiler:**
 - `CC=g++` in order to use the GNU C++ compiler.
 - `CC=clang -lstdc++` in order to use the `clang` C++ compiler.
- **Optimization level:**
 - `OPT=O0`: No optimization.
 - `O=O1, O2, or O3` (marginally faster, default): some optimization occurs (read the compiler documentation for more information).
 - `OPT=Ofast`: full optimization (fast, but dangerous).

Examples

1

Axion Dark Matter

- Why particle dark matter
- The dark matter particle
- The axion (like) particle

2

Calculating the Relic Abundance

- The axion EOM
- How hard can it be?
- Initial conditions
- (Bad) Analytical approximations
- Need for accuracy, speed, and automation

3

MiMeS

- MiMeS: General
- MiMeS: Under the hood
- MiMeS: Notation
- MiMeS: When do you start and stop integrating?
- MiMeS: Some (classical) physics

4

Using MiMeS

- How to get MiMeS
- Configure (and make)
- Classes
- Assumptions
- What MiMeS expects from you
- Template arguments
- MiMeS from python

5

Examples

- C++
- python

6

Outlook

- Summing up
- Future

Define everything and solve in just a few lines of code!

```

1 // use chi_PATH to interpolate the axion mass.
2 mimes::AxionMass<long double> axionMass(chi_PATH,0,mimes::Cosmo<long double>::mP);
3
4 /*set  $\tilde{m}_a^2$  for  $T \geq T_{\max}$ */
5 long double TMax=axionMass.getTMax(), chiMax=axionMass.getChiMax();
6
7 axionMass.set_ma2_MAX(
8     [&chiMax,&TMax](long double T, long double fa){
9         return chiMax/fa/fa*std::pow(T/TMax,-8.16);
10     }
11 );
12
13 /*set  $\tilde{m}_a^2$  for  $T \leq T_{\min}$ */
14 long double TMin=axionMass.getTMin(), chiMin=axionMass.getChiMin();
15
16 axionMass.set_ma2_MIN(
17     [&chiMin,&TMin](long double T, long double fa){
18         return chiMin/fa/fa;
19     }
20 );
21
22 /*this path contains the cosmology*/
23 std::string inputFile = std::string(rootDir)+
24     std::string("/UserSpace/InputExamples/MatterInput.dat");
25
26 /*declare an instance of Axion*/
27 mimes::Axion<long double, 1, RODASPR2<long double> > ax(
28     0.1, 1e16, 500, 1e-4, 1e3, 15, 1e-2, inputFile, &axionMass,
29     1e-1, 1e-8, 1e-1, 1e-11, 1e-11, 0.9, 1.2, 0.8, int(1e7)
30 );
31
32 /*solve the EOM!*/
33 ax.solveAxion();

```

Outlook

1

Axion Dark Matter

- Why particle dark matter
- The dark matter particle
- The axion (like) particle

2

Calculating the Relic Abundance

- The axion EOM
- How hard can it be?
- Initial conditions
- (Bad) Analytical approximations
- Need for accuracy, speed, and automation

3

MiMeS

- MiMeS: General
- MiMeS: Under the hood
- MiMeS: Notation
- MiMeS: When do you start and stop integrating?
- MiMeS: Some (classical) physics

4

Using MiMeS

- How to get MiMeS
- Configure (and make)
- Classes
- Assumptions
- What MiMeS expects from you
- Template arguments
- MiMeS from python

5

Examples

- C++
- python

6

Outlook

- Summing up
- Future

Summing up

Commend from D: blah blah, what you did

Commend from D: Freeze-out/in, kinematic misalignment mechanism, automatically compare with searches.

Thank you!

Language	files	blank	comment	code
C/C++ Header	35	677	444	1595
C++	20	483	198	1106
Python	22	483	367	1000
SUM:	77	1643	1009	3701

Backup

(equations, derivations, tables)

Adiabatic invariant

Input – `mimes::AxionMass<LD>`

Input – `mimes::Axion<LD, Solver, Method>`