

Leading Causes of Death by Age Group, American Indian or Alaska Native Males-United States, 2012*

Age in Groups

Rank	1-4	5-9	10-14	15-19	20-24	25-34	35-44	45-54	55-64	65+	65-74	75-84	85+	All Ages
1	Unintentional injuries 51.9%	Unintentional injuries 82.4%	Unintentional injuries 29.0%	Unintentional injuries 42.7%	Unintentional injuries 44.4%	Unintentional injuries 39.6%	Unintentional injuries 29.5%	Unintentional injuries 18.3%	Cancer 23.0%	Heart disease 25.2%	Cancer 28.7%	Heart disease 27.2%	Heart disease 26.4%	Heart disease 19.1%
2	Homicide 11.1%	Cancer (2) 2.9%	Suicide 25.8%	Suicide 30.1%	Suicide 25.7%	Suicide 14.7%	Chronic liver disease 13.7%	Heart disease 17.5%	Heart disease 21.1%	Cancer 23.3%	Heart disease 23.2%	Cancer 21.1%	Cancer 15.0%	Cancer 17.2%
3	Cancer 7.4%	Influenza & pneumonia (2) 2.9%	Heart disease 12.9%	Homicide 9.8%	Homicide 12.3%	Homicide 11.1%	Heart disease 12.0%	Cancer 13.2%	Unintentional injuries 9.5%	Chronic lower respiratory diseases 7.0%	Diabetes 7.9%	Chronic lower respiratory diseases 8.4%	Chronic lower respiratory diseases 7.3%	Unintentional injuries 14.1%
4	Birth defects 5.6%	Homicide (2) 2.9%	Cancer 9.7%	Heart disease 3.5%	Cancer 2.1%	Heart disease 7.3%	Suicide 9.1%	Chronic liver disease 11.8%	Chronic liver disease 7.8%	Diabetes 6.9%	Chronic lower respiratory diseases 5.7%	Diabetes 7.0%	Stroke 5.9%	Chronic liver disease 5.7%
5	Stroke 1.9%		Birth defects 6.5%	Cancer 2.1%	Heart disease (5) 1.8%	Chronic liver disease 6.0%	Cancer 5.7%	Diabetes 5.7%	Diabetes 7.6%	Stroke 4.6%	Unintentional injuries 4.6%	Stroke 4.1%	Diabetes 4.5%	Diabetes 5.7%
6			Benign neoplasms 3.2%	Birth defects 1.4%	Chronic liver disease (5) 1.8%	Cancer 2.9%	Homicide 4.2%	Suicide 4.2%	Chronic lower respiratory diseases 4.4%	Unintentional injuries 3.5%	Stroke 4.3%	Unintentional injuries 2.6%	Influenza & pneumonia 3.5%	Suicide 4.3%
7				Diabetes (7) 0.7%	Benign neoplasms (7) 1.1%	Influenza & pneumonia 1.3%	Diabetes 3.8%	Homicide 2.4%	Stroke 2.9%	Kidney disease 2.2%	Chronic liver disease 2.9%	Influenza & pneumonia 2.5%	Kidney disease (7) 2.8%	Chronic lower respiratory diseases 4.1%
8					Legal intervention (7) 1.1%	Diabetes 1.0%	HIV disease 1.2%	Stroke 2.1%	Suicide 1.7%	Influenza & pneumonia 2.1%	Kidney disease 1.9%	Kidney disease 2.4%	Unintentional injuries (7) 2.8%	Stroke 2.9%
9					Hypertension (9) 0.7%	Septicemia (9) 0.7%	Kidney disease 1.1%	Viral hepatitis 1.6%	Influenza & pneumonia 1.7%	Chronic liver disease 2.0%	Septicemia 1.7%	Alzheimer's disease 1.8%	Alzheimer's disease 2.5%	Homicide 2.3%
10					Kidney disease (9) 0.7%	Birth defects (9) 0.7%	Septicemia 1.0%	Chronic lower respiratory diseases 1.3%	Kidney disease 1.6%	Septicemia 1.6%	Influenza & pneumonia 1.3%	Septicemia 1.7%	Hypertension 2.0%	Influenza & pneumonia 1.6%
						Legal intervention (9) 0.7%								

*Percentages represent total deaths in the age group due to the cause indicated. Numbers in parentheses indicate tied rankings. The white, black, American Indian/Alaska Native, and Asian/Pacific Islander race groups include persons of Hispanic and non-Hispanic origin may be of any race. Some terms have been shortened from those used in the National Vital Statistics Report. See the next page for a listing of the shortened terms in the table and their full unabridged equivalents used in the report. To learn more, visit Mortality Tables at http://www.cdc.gov/nchs/nvss/mortality_tables.htm or <http://www.cdc.gov/nchs/deaths.htm> (HHS, CDC, NCHS).

Short and Full Terms for Leading Causes of Death (Males)

Some terms in the leading causes of death table have been shortened from those used in the National Vital Statistics Report. Below is a listing of the shortened terms used in the table and their full, unabridged equivalents used in the report.

SHORT TERMS

Aortic aneurism
Benign neoplasms
Birth defects
Bronchitis
Cancer
C. difficile infection
Chronic liver disease
Chronic lower respiratory diseases

Diabetes
Gallbladder disorders
Heart disease
HIV disease
Homicide
Hypertension
Kidney disease
Kidney infection
Legal Intervention*

Medical & surgical care complications
Operations of War
Perinatal conditions
Pneumonitis
Stroke
Suicide
Unintentional injuries

FULL TERMS

Aortic aneurism and dissection
In situ neoplasms and neoplasms of uncertain or unknown behavior
Congenital malformations, deformations and chromosomal abnormalities
Acute bronchitis and bronchiolitis
Malignant neoplasms
Enter colitis due to clostridium
Chronic liver disease and cirrhosis
Includes bronchitis, emphysema, asthma, bronchiectasis, and other chronic obstructive pulmonary disease
Diabetes mellitus
Cholelithiasis and other disorders of gallbladder
Diseases of the heart
Human immunodeficiency virus (HIV) disease
Assault (homicide)
Essential (primary) hypertension and hypertensive renal disease
Nephritis, nephrotic syndrome and nephrosis
Infections of kidney
Physical or other force used by police or other law-enforcing agents, including military on duty, in the course of arresting or attempting to arrest lawbreakers, suppressing disturbances, maintaining order, and other legal action. Includes legal execution and excludes citizen arrest.

Complications of medical and surgical care
Operations of War and their sequelae
Certain conditions originating in the perinatal period
Pneumonitis due to solids and liquids
Cerebrovascular diseases
Intentional self-harm
Accidents (unintentional injuries)

*World Health Organization. International statistical Classification of Diseases and Related Health Problems, Tenth Revision (ICD-10). Geneva: World Health Organization, 1992. Available at <http://www.who.int/classifications/icd/en/>. Accessed April 15, 2010.