

18.701: Problem Set 9

Dmitry Kaysin

July 2020

Problem 1

Find an orthonormal basis for the vector space P_2 of all real polynomials of degree at most 2 with the symmetric form defined by

$$\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx.$$

Proof. We first check that the form $\langle \cdot, \cdot \rangle$ is indeed symmetric. This is the case since integration is linear and multiplication of real polynomials is commutative.

To find an orthonormal basis with the given symmetric form we will apply the Gram-Schmidt procedure. We start with an arbitrary basis of P_2 :

$$B = \{1, x, x^2\}.$$

We will determine orthogonal basis B' of P_2 first.

We start with the first vector $v_1 = 1$ of B . This will be the first vector of B' .

We notice that

$$\langle 1, x \rangle = \int_{-1}^1 x dx = \frac{x^2}{2} \Big|_{-1}^1 = 0,$$

which means that $v_2 = x$ is orthogonal to $v_1 = w_1 = 1$. We take $w_2 = x$ as the second vector of B' .

We proceed by finding the orthogonal projection of the third vector $v_3 = x^2$ onto the subspace spanned by the orthogonal vectors $\{w_1, w_2\}$:

$$\pi(v_3) = w_1 \frac{\langle w_1, v_3 \rangle}{\langle w_1, w_1 \rangle} + w_2 \frac{\langle w_2, v_3 \rangle}{\langle w_2, w_2 \rangle},$$

where

$$\begin{aligned} \langle w_1, v_3 \rangle &= \int_{-1}^1 x^2 dx = \frac{x^3}{3} \Big|_{-1}^1 = \frac{2}{3}, & \langle w_1, w_1 \rangle &= \int_{-1}^1 dx = x \Big|_{-1}^1 = 2, \\ \langle w_2, v_3 \rangle &= \int_{-1}^1 x^3 dx = \frac{x^4}{4} \Big|_{-1}^1 = 0, & \langle w_2, w_2 \rangle &= \int_{-1}^1 x^2 dx = \frac{x^3}{3} \Big|_{-1}^1 = \frac{2}{3}. \end{aligned}$$

Thus

$$\pi(v_3) = w_1 \frac{\langle w_1, v_3 \rangle}{\langle w_1, w_1 \rangle} + w_2 \frac{\langle w_2, v_3 \rangle}{\langle w_2, w_2 \rangle} = \frac{1}{3}.$$

Since $v_3 - \pi(v_3)$ is orthogonal to both w_1 and w_2 , we take it as the third vector of B' :

$$w_3 = x^2 - \frac{1}{3}.$$

We now have an orthogonal basis $B' = \{1, x, x^2 - \frac{1}{3}\}$. We normalize each vector in B' :

$$\begin{aligned} w'_1 &= \frac{w_1}{\langle w_1, w_1 \rangle^{1/2}} = \frac{\sqrt{2}}{2}, \\ w'_2 &= \frac{w_2}{\langle w_2, w_2 \rangle^{1/2}} = \sqrt{\frac{3}{2}}x, \\ w'_3 &= \frac{w_3}{\langle w_3, w_3 \rangle^{1/2}} = \frac{3\sqrt{5}}{2\sqrt{2}} \left(x^2 - \frac{1}{3} \right), \end{aligned}$$

where

$$\begin{aligned} \langle w_3, w_3 \rangle &= \left\langle x^2 - \frac{1}{3}, x^2 - \frac{1}{3} \right\rangle = \int_{-1}^1 \left(x^2 - \frac{1}{3} \right)^2 dx = \int_{-1}^1 \left(x^4 - \frac{2}{3}x^2 + \frac{1}{9} \right) dx \\ &= \left(\frac{x^5}{5} - \frac{2}{9}x^3 + \frac{1}{9}x \right) \Big|_{-1}^1 = \frac{2}{5} - \frac{4}{9} + \frac{2}{9} = \frac{8}{45}. \end{aligned}$$

Thus, $\left\{ \frac{\sqrt{2}}{2}, \sqrt{\frac{3}{2}}x, \frac{3\sqrt{5}}{2\sqrt{2}} \left(x^2 - \frac{1}{3} \right) \right\}$ is an orthonormal basis of P_2 .

□

Problem 2

Let V be the real vector space of 3×3 matrices with the bilinear form $\langle A, B \rangle = \text{trace } A^t B$, and let W be the subspace of skew-symmetric matrices. Compute the orthogonal projection to W with respect to this form, of the matrix

$$v = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 3 & 0 \end{pmatrix}.$$

Real 3×3 skew-symmetric matrices have the form

$$\begin{pmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{pmatrix}.$$

Thus vector space W has dimension 3 with one basis given as the set of vectors

$$w_1 = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad w_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}.$$

One can easily verify by direct computation that all three vectors w_1, w_2, w_3 are orthogonal and $\langle w_1, w_1 \rangle = \langle w_2, w_2 \rangle = \langle w_3, w_3 \rangle = 2$.

Therefore, orthogonal projection of the matrix v to the subspace W can be calculated as follows:

$$\begin{aligned} \pi_W(v) &= \frac{\langle w_1, v \rangle}{\langle w_1, w_1 \rangle} w_1 + \frac{\langle w_2, v \rangle}{\langle w_2, w_2 \rangle} w_2 + \frac{\langle w_3, v \rangle}{\langle w_3, w_3 \rangle} w_3 \\ &= \frac{2}{2} w_1 + \frac{-1}{2} w_2 + \frac{-2}{2} w_3 = w_1 - \frac{1}{2} w_2 - w_3, \end{aligned}$$

which corresponds to the matrix

$$\begin{pmatrix} 0 & 1 & -\frac{1}{2} \\ -1 & 0 & -1 \\ \frac{1}{2} & 1 & 0 \end{pmatrix}.$$

Problem 3

Let W be a two-dimensional subspace of \mathbb{R}^3 , and consider the orthogonal projection π of \mathbb{R}^3 onto W . Let $(a_i, b_i)^t$ be the coordinate vector of $\pi(e_i)$, with respect to a chosen orthonormal basis of W . Prove that (a_1, a_2, a_3) and (b_1, b_2, b_3) are orthogonal unit vectors.

Proof. Let $B = \{u, v\}$ be the chosen orthonormal basis of W and suppose, u has coordinates $(u_1, u_2, u_3)^t$ and w has coordinates $(w_1, w_2, w_3)^t$ in the standard basis.

Applying the projection formula:

$$\pi(e_1) = \langle u, e_1 \rangle u + \langle v, e_1 \rangle v = \left\langle \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right\rangle u + \left\langle \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right\rangle v = u_1 u + w_1 v.$$

Therefore, $a_1 = u_1$, $b_1 = w_1$.

Similarly we find coordinates for projections of other standard basis vectors and combine them into row vectors

$$\begin{aligned} (a_1, a_2, a_3) &= (u_1, u_2, u_3) = u^t, \\ (b_1, b_2, b_3) &= (w_1, w_2, w_3) = w^t. \end{aligned}$$

Since u and w are orthogonal, u^t and w^t are also orthogonal.

□

Problem 4

Prove that the signature of the form does not depend on the choice of the orthogonal basis (Sylvester's Law).

We first prove that if W^+ and W^- are subspaces of V and if the form is positive definite on W^+ and negative definite on W^- , then W^+ and W^- are independent.

Proof. It suffices to show that $W^+ \cap W^- = 0$. Consider vector $w \in W^+ \cap W^-$. Then both

$$\langle w, w \rangle \leq 0 \quad \text{and} \quad \langle w, w \rangle \geq 0$$

must be true. Since the form is nondegenerate on V , we conclude that $w = 0$. □

Let V be complex vector space equipped with a Hermitian form that is nondegenerate on V . Let $B = \{v_1, \dots, v_k, v_{k+1}, \dots, v_{k+n}\}$ be an orthonormal basis of V such that

$$I_{k,n} = \begin{pmatrix} I_k & \\ & -I_n \end{pmatrix}$$

is the matrix of the form with respect to basis B . Denote $W^+ = \text{span}\{v_1, \dots, v_k\}$, and denote $W^- = \text{span}\{v_{k+1}, \dots, v_{k+n}\}$. Choose $B' = \{w_1, \dots, w_{k+n}\}$, another orthonormal basis of V such that matrix of the form with respect to basis B' is diagonal with entries 1, -1. We will prove that matrix of the form with respect to B' is the same as the matrix of the form with respect to basis B .

Proof. Every vector of B' is either an element of W^+ or W^- . There are exactly k vectors of B' that are elements of W^+ since W^+ is of dimension k and vectors of B' are independent. Similarly, there are exactly n vectors of B' that are elements of W^- . If w_i is in W^+ , then $\langle w_i, w_i \rangle = 1$ since B' is orthonormal. Conversely, if w_i is in W^- , then $\langle w_i, w_i \rangle = -1$. Therefore, matrix of the form with respect to basis B' (after rearranging basis vector appropriately) is as follows:

$$I_{k,n} = \begin{pmatrix} I_k & \\ & -I_n \end{pmatrix}.$$

□

Problem 5

Let T be a linear operator on $V = \mathbb{R}^n$ whose matrix A is a real symmetric matrix.

a) Prove that V is the orthogonal sum $V = (\ker T) \oplus (\text{im } T)$.

Proof.

□

b) Prove that T is an orthogonal projection onto $\text{im } T$ if and only if, in addition to being symmetric, $A^2 = A$.

Proof.

□