

Test Unit 2
PCHA 2020-21 / Dr. Kessner

Name:

No calculator, no notes – just your brain! Have fun!

1. Evaluate the following:

a) $\sec \frac{5\pi}{3} = \frac{1}{\frac{1}{2}} = 2$




b) $\sin(-\frac{2\pi}{3}) = -\frac{\sqrt{3}}{2}$

c) $\cos^{-1}(\underbrace{\cot(-\frac{5\pi}{4})}_{-1}) = \pi$

d) $\sin^{-1}(\underbrace{\cos(-\frac{\pi}{2})}_0) = 0$

e) $\tan^{-1}(\underbrace{\cot(\frac{\pi}{6})}_{\sqrt{3}}) = \frac{\pi}{3}$

f) $\sin(-\frac{\pi}{12}) = \sin(\frac{\pi}{4} - \frac{\pi}{3})$
 $= \sin \frac{\pi}{4} \cos \frac{\pi}{3} + \cos \frac{\pi}{4} \sin(-\frac{\pi}{3})$
 $= \frac{\sqrt{2}}{2} \frac{1}{2} + \frac{\sqrt{2}}{2} (-\frac{\sqrt{3}}{2})$
 $= \frac{\sqrt{2} - \sqrt{6}}{4}$



2. Write down all the relevant properties (period, amplitude, shifts/scales, asymptotes) of the following trig functions, and then graph by hand.


$$f(x) = 4 + 4 \cos\left(\frac{\pi}{6}x - \frac{\pi}{2}\right)$$

$$= 4 + 4 \cos\left(\frac{\pi}{6}(x - 3)\right)$$

period $\frac{2\pi}{(\pi/6)} = 12$

shift right 3, up 4

amplitude 4




$$g(x) = 5 \cot\left(\frac{\pi}{8}(x - 4)\right)$$

period $\frac{\pi}{(\pi/8)} = 8$

shift left 4

vertical scale 5



Bonus Write f as a transformed sin and g as a transformed tan.

$$4 + 4 \sin\left(\frac{\pi}{6}x\right) \quad - 5 \tan\left(\frac{\pi}{8}x\right)$$

3. Prove the identities:

$$\frac{\cos x + \sin x}{\cos x - \sin x} = \sec 2x + \tan 2x$$

$$\begin{aligned} & \frac{\cos x + \sin x}{\cos x - \sin x} \cdot \frac{\cos x + \sin x}{\cos x + \sin x} \\ &= \frac{(\cos x + \sin x)^2}{\cos^2 x - \sin^2 x} \\ &= \frac{\cos^2 x + \sin^2 x + 2 \sin x \cos x}{\cos^2 x - \sin^2 x} \\ &= \frac{1 + \sin 2x}{\cos 2x} \\ &= \sec 2x + \tan 2x \quad \checkmark \end{aligned}$$

$$\sin(\pi - x) = \sin x$$

$$\begin{aligned} \sin(\pi - x) &= \underbrace{\sin \pi}_{=0} \cos x - \underbrace{\cos \pi}_{=-1} \sin x \\ &= \sin x \quad \checkmark \end{aligned}$$

Bonus Prove this using cofactor identities.

4. Find all solutions of $\sin 2\theta + \cos \theta = 0$.


$$2 \sin \theta \cos \theta + \cos \theta = 0$$

$$\cos \theta (2 \sin \theta + 1) = 0$$

$$\cos \theta = 0 \quad \text{or} \quad \sin \theta = -\frac{1}{2}$$

$$\theta = \frac{\pi}{2} + \pi k \quad \theta = -\frac{\pi}{6} + 2\pi k$$

$$-\frac{5\pi}{6} + 2\pi k$$



Derive the following half angle formula from the relevant double angle formula:

$$\cos u = \pm \sqrt{\frac{1 + \cos 2u}{2}}$$


$$\cos 2u = \cos^2 u - \sin^2 u$$

$$= 2\cos^2 u - 1$$

$$2\cos^2 u = 1 + \cos 2u$$

$$\cos^2 u = \frac{1 + \cos 2u}{2}$$

$$\cos u = \pm \sqrt{\frac{1 + \cos 2u}{2}}$$




Use the half angle formula above to find $\cos(-\frac{5\pi}{12})$.

$$\cos\left(-\frac{5\pi}{12}\right) = + \sqrt{\frac{1 + \cos\left(-\frac{5\pi}{6}\right)}{2}}$$

$$= \sqrt{\frac{1 + (-\sqrt{3}/2)}{2}}$$

$$= \frac{1}{2} \sqrt{2 - \sqrt{3}}$$

5. Solve the following triangle: $A = \frac{\pi}{4}$, $b = 5$, $c = 5\sqrt{2}$.



$$\begin{aligned} a^2 &= b^2 + c^2 - 2bc \cos A \\ &= 25 + 50 - 2 \cdot 5 \cdot 5\sqrt{2} \cos \frac{\pi}{4} \\ &= 25 + 50 - 50 \quad \underbrace{\cos \frac{\pi}{4} = \frac{\sqrt{2}}{2}} \\ &= 25 \end{aligned}$$

$$\begin{aligned} a &= 5 \\ \sin C &= \frac{c \sin A}{a} = \frac{5\sqrt{2} \sin \frac{\pi}{4}}{5} = 1 \\ &\Rightarrow C = \frac{\pi}{2} \end{aligned}$$

$$B = \frac{\pi}{4}$$