

The Coastal Carbon Library and Atlas: Open source soil data and tools supporting blue carbon research and policy

James R. Holmquist¹  | David Klinges^{1,2} | Michael Lonneman¹ | Jaxine Wolfe¹ |
 Brandon Boyd³ | Meagan Eagle⁴ | Jonathan Sanderman⁵ | Kathe Todd-Brown²  |
 E. Fay Belshe² | Lauren N. Brown⁶ | Samantha Chapman⁷ | Ron Corstanje⁸ |
 Christopher Janousek⁹ | James T. Morris¹⁰ | Gregory Noe¹¹ | André Rovai¹²  |
 Amanda Spivak¹³  | Megan Vahsen¹⁴ | Lisamarie Windham-Myers¹⁵ | Kevin Kroeger⁴ |
 J. Patrick Megonigal¹ 

¹Smithsonian Environmental Research Center, Edgewater, Maryland, USA

²University of Florida, Gainesville, Florida, USA

³Army Corps of Engineers, Washington, DC, USA

⁴USGS, Woods Hole, Massachusetts, USA

⁵Woods Hole Research Center, Woods Hole, Massachusetts, USA

⁶Bowling Green State University, Bowling Green, Ohio, USA

⁷Villanova University, Villanova, Pennsylvania, USA

⁸Cranfield University, Cranfield, UK

⁹Oregon State University, Corvallis, Oregon, USA

¹⁰University of South Carolina, Columbia, South Carolina, USA

¹¹USGS, Wetland Ecosystem Ecology and Biogeochemistry Laboratory, Reston, Virginia, USA

¹²Louisiana State University, Baton Rouge, Louisiana, USA

¹³University of Georgia, Athens, Georgia, USA

¹⁴Notre Dame University, Notre Dame, Indiana, USA

¹⁵USGS, Water Missions Area, Menlo Park, California, USA

Correspondence

James R. Holmquist, Smithsonian Environmental Research Center, 647 Contees Wharf Road, Edgewater, MD 21037, USA.
 Email: holmquistj@si.edu

Funding information

Department of Energy, Grant/Award Number: COMPASS-FME; Division of Environmental Biology, Grant/Award Number: DEB-1655622; National Oceanic and Atmospheric Administration, Grant/Award Number: NA22OAR4310148; National Science Foundation, Grant/Award Number: DGE-1842473; Pew Charitable Trusts

Abstract

Quantifying carbon fluxes into and out of coastal soils is critical to meeting greenhouse gas reduction and coastal resiliency goals. Numerous 'blue carbon' studies have generated, or benefitted from, synthetic datasets. However, the community those efforts inspired does not have a centralized, standardized database of disaggregated data used to estimate carbon stocks and fluxes. In this paper, we describe a data structure designed to standardize data reporting, maximize reuse, and maintain a chain of credit from synthesis to original source. We introduce version 1.0.0. of the Coastal Carbon Library, a global database of 6723 soil profiles representing blue carbon-storing systems including marshes, mangroves, tidal freshwater forests, and seagrasses. We also present the Coastal Carbon Atlas, an R-shiny application that can

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 Smithsonian Institution and The Authors. *Global Change Biology* published by John Wiley & Sons Ltd. This article has been contributed to by U.S. Government employees and their work is in the public domain in the USA.

be used to visualize, query, and download portions of the Coastal Carbon Library. The majority (4815) of entries in the database can be used for carbon stock assessments without the need for interpolating missing soil variables, 533 are available for estimating carbon burial rate, and 326 are useful for fitting dynamic soil formation models. Organic matter density significantly varied by habitat with tidal freshwater forests having the highest density, and seagrasses having the lowest. Future work could involve expansion of the synthesis to include more deep stock assessments, increasing the representation of data outside of the U.S., and increasing the amount of data available for mangroves and seagrasses, especially carbon burial rate data. We present proposed best practices for blue carbon data including an emphasis on disaggregation, data publication, dataset documentation, and use of standardized vocabulary and templates whenever appropriate. To conclude, the Coastal Carbon Library and Atlas serve as a general example of a grassroots F.A.I.R. (Findable, Accessible, Interoperable, and Reusable) data effort demonstrating how data producers can coordinate to develop tools relevant to policy and decision-making.

KEY WORDS

blue carbon, carbon sequestration, coastal wetland, open data, sea-level rise, soil organic carbon, synthesis

1 | INTRODUCTION

Globally, wetland soils are estimated to contain 500–680 Petagrams of carbon (Poulter et al., 2021), equivalent to 60%–82% of all carbon present in the atmosphere (Ciais et al., 2014). Coastal wetlands such as tidal marshes, tidal freshwater forests, referred to throughout as swamps, mangroves, and seagrasses are known to sequester carbon on decadal to millennial time scales (Chmura et al., 2003; Howard et al., 2017; Mcleod et al., 2011; Ouyang & Lee, 2014). So-called blue carbon habitats have been garnering attention in recent years because of the capacity for management interventions to avoid greenhouse gas emissions by conserving wetlands or re-establish greenhouse gas removals by restoring them (Eagle et al., 2022; Kroeger et al., 2017; Lovelock & Duarte, 2019; Wylie et al., 2016).

Coastal wetland soils are unique in that they store carbon in situ as a dynamic response to sea-level rise (Morris et al., 2002) via mineral trapping and organic soil mass production (Morris et al., 2002; Turner et al., 2006). These dynamics mean that factors such as relative sea-level rise (Rogers et al., 2019), plant community (Doughty et al., 2015; Schile et al., 2017), and elevation (Callaway et al., 2012; Peck et al., 2020b), are of great importance to predicting carbon stocks and burial rates. Given the fact that tidal wetlands were underrepresented in previous efforts (Holmquist et al., 2018b), the requirements of specialist and interdisciplinary knowledge represented in tidal wetland datasets, and their distinctness from terrestrial soils, we propose that soils from blue carbon habitats require their own dedicated synthesis effort for disaggregated data.

Data syntheses for coastal wetland ecosystems have been undertaken previously. An original synthesis by Chmura et al. (2003)

established coastal wetland carbon storage as a major carbon sink. A subsequent expansion by Ouyang and Lee (2014) documented covariates associated with carbon burial. These foundational studies provided data reported or estimated from literature values but did not present original disaggregated information, original measurements rather than summary statistics, limiting their reuse for other purposes. Later synthesis datasets greatly expanded the available disaggregated carbon stocks data available for seagrasses (Fourqurean et al., 2012), tidal marshes (Holmquist et al., 2018b), tidal swamps (Krauss et al., 2018), and mangroves (Donato et al., 2011; Sanderman, 2017). These studies have many elements that a comprehensive database would require, but none were designed to be a living resource for the communities they helped establish.

Sharing of disaggregated data is vital to discovery in coastal carbon studies. Morris et al. (2016) documented a relationship between soil organic content and density which limits vertical accretion capacity. Holmquist et al. (2018b) independently validated soil carbon stock mapping strategies, uncovering an overestimate in carbon stock when relying on conventional maps. Rogers et al. (2019) utilized an earlier version of the database presented herein to show that the geography of relative sea-level rise explained variability in carbon density at a global scale. All of these important studies would not be possible without the sharing of disaggregated data from researcher to researcher. However, data sharing between researchers is much rarer in practice than is implied through data availability statements in academic peer-reviewed journal articles (Tedesoo et al., 2021).

In this paper, we describe the creation and implementation of a set of standards and introduce the Coastal Carbon Library v1.0.0. This database is unique from previous efforts in that it spans multiple



habitats, pays special attention to gathering site characteristics and methodological meta-data required to re-interpret data for new uses, and it contains disaggregated data, rather than summary statistics. We then present the Coastal Carbon Atlas (https://shiny.si.edu/coastal_carbon_atlas), a web tool for visualizing, querying, and downloading the Coastal Carbon Library. We also discuss the strengths of this database in the face of rapidly changing coastal ecosystems. Finally, we discuss potential future directions and proposed best practices.

2 | METHODS

Belowground carbon studies use soil depth series data to quantify belowground carbon stocks and estimate carbon burial rates (Chmura et al., 2003; Needelman et al., 2018; Ouyang & Lee, 2014). Typically cores are extracted using a tube or auger, and sub-sectioned into multiple depth increments (Smeaton et al., 2020). Organic matter density or carbon density is usually assessed as the product of dry bulk density (Blake, 1965), the organic matter fraction using loss-on-ignition (Dean, 1974) and/or fraction organic carbon (Callaway et al., 2012; Craft et al., 1991; Poppe & Rybczyk, 2018). Multiple methods are also employed for creating models of the accumulation or accretion rate of the profile including lead-210 (^{210}Pb ; Appleby & Oldfield, 1978; DeLaune et al., 1987), radiocarbon (^{14}C ; Redfield, 1972), radiocesium (^{137}Cs ; DeLaune et al., 1978), as well as tying traces of pollution (Gerlach et al., 2017), pollen (Gerlach et al., 2017; Kearney & Ward, 1986), or sediment quality (Abbott, Elsey-Quirk, & DeLaune, 2019; Drexler et al., 2019) to dated historical events.

2.1 | Goals and scope

F.A.I.R. (Findable, Accessible, Interoperable, and Reusable) data principles are a set of best practices, seeking to maximize the re-usability of data. To meet goals of F.A.I.R. data, the goals of the data structure were four-fold, and grounded in best practices for data management (Wilkinson et al., 2016; Wilson et al., 2017). First, we preserved data in as much detail as is practical to maximize versatility. Second, we were transparent, relying on stable repositories. Third, we explicitly connect individual datasets within the synthesis to their original sources through one more associated citations. Fourth, we emphasized simplicity.

For scope, we focused on vegetated coastal tidal wetlands. We focused on emergent vegetation (marsh), scrub/shrub, and forested wetlands including mangroves and tidal swamps. We included seagrasses and tidal flats, but not kelp beds, coral reefs, or oyster reefs (Howard et al., 2017). These included saltwater, brackish, and freshwater ecosystems in the tidal zone and subtidal seagrasses. In addition to current tidal wetlands, we also included former tidal wetlands that had been modified by human impact and uplands or non-tidal wetlands if that comparison was a key aspect of the original study. Finally, while we did classify datasets based on their completeness or utility, we did not exclude any datasets based on methods used, origins, or lack of attributes beyond a few key requirements such as

basic positional and depth information, as well as at least one relevant measured depth series attribute (Data S1).

2.2 | Data structure

Coastal Carbon Library data formatting (Holmquist et al., 2023) follows practices (Wilson et al., 2017) for maximizing data reuse such as descriptive attribute names, consistently formatted 'no data' values, multiple quality control attributes with consistent formatting, and multiple places for explanatory comments written in prose. Data files are long rather than wide, designating each row as a single observation, each column as a single variable, and each table as a set of observations and variables (Wickham, 2014). Data files are 'flat', all information is stored as simple, uniform, text. The structure consists of seven data tables (Figure 1):

1. Site: contains site-level information (Table 1).
2. Cores: contains core-level information (Table 2).
3. Depth series: provides the disaggregated information collected across depth increments of each soil profile (Table 3).
4. Methods: contains information about the methodology used for a collection of sampling and measurement events (Table 4).
5. Species: provides the plant species observed proximal to the sampling location (Table 5).
6. Impacts: provides a controlled classification of anthropogenic impacts associated with the sampling location (Table 6).
7. Bibliography: associated information of sources associated with datasets (Table 7).

All tables are relational with common linking variables study_id, site_id, methods_id, core_id, and bibliography_id. The site, core, depth series, species, impacts, and methods tables are hierarchical. Depth increments are nested within cores. Cores are nested within sites. Species and impacts are nested within cores, or within sites, if site-level data are the highest resolution provided by the study (Figure 1). Categorical variables have defined vocabulary listed in Table 8.

2.3 | Data sourcing

Data entered the Coastal Carbon Library through two tracks, with and without curatorial assistance by Coastal Carbon Network personnel. If a data source was already available through a data publication then the original data and associated metadata were downloaded to the Library repository (Figure 2). Ingestion scripts were then written in R code (R Core Team, 2021) to reshape, and reformat so that datasets could be merged (Figure 2).

If a dataset was unpublished, Coastal Carbon Network personnel first assisted the data submitter in creating a data publication on Figshare (Figure 2). Data releases included a title, abstract, keywords, authors, associated publications, acknowledgment of funding sources, data files, and ecological meta-data language (EML)-style metadata, an HTML version of the metadata, and a




FIGURE 1 Relationships between the tables that make up the Coastal Carbon Library. Attributes tracked in each table are listed. Bold and italicized text indicates common attributes that can be used to join tables. The structure of the library is mostly hierarchical (a site contains multiple cores, and a core contains multiple depth increments in a series). One coring location can also have multiple anthropogenic impacts and plant species associated with it. A methods table links directly to depth increments as more than one set of methods may be used to analyze depth increments within a core.

citable digital object identifier (DOI). We published 27 datasets in this way (Abbott, Quirk, & Delaune, 2019; Allen et al., 2022; Arias-Ortiz, Masqué, et al., 2021; Belshe et al., 2019b; Boyd et al., 2019; Breithaupt et al., 2019, 2020; Buffington et al., 2020; Callaway et al., 2019; Carlin et al., 2021; Doughty et al., 2019; Ensign et al., 2021; Fell et al., 2021; Kauffman et al., 2020b; Kemp et al., 2020; Keshta et al., 2020; Lagomasino et al., 2020; McTigue et al., 2020; Messerschmidt & Kirwan, 2020; Nolte, 2020; Peck et al., 2020a; Poppe & Rybczyk, 2019; Sanbor & Darwyn, 2020;

St. Laurent et al., 2020b; Thom, 2019; Vaughn et al., 2021; Weston et al., 2022).

2.4 | Data quality control and quality assessment

Data were quality-controlled using automated and visual checks. Automated quality tests were performed at both the individual study and whole synthesis levels and included (but were not limited to):

TABLE 1 Attribute information for site table.

| Attribute name | Definition | Data type | Unit |
|--------------------|---|-------------|--------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text | |
| site_id | Site identification code unique to each study | Free text | |
| site_description | Site description includes relevant study details and political geographic units. Some of these descriptions can be automated by the ingestion code. | Free text | |
| site_latitude_max | Maximum latitude defining a bounding box for the site in decimal degree World Geodetic System of 1984 (WGS84) | Numeric | Degree |
| site_latitude_min | Minimum latitude defining a bounding box for the site in decimal degree WGS84 | Numeric | Degree |
| site_longitude_max | Maximum longitude defining a bounding box for the site in decimal degree WGS84 | Numeric | Degree |
| site_longitude_min | Minimum longitude defining a bounding box for the site in decimal degree WGS84 | Numeric | Degree |
| site_boundaries | As an alternative to submitting or automatically generating a bounding box, submitters can include a shapefile (.shp) or keyhole markup language (.kml) documenting the geographic boundaries of the site. This can be converted to and stored in well-known text format. | Free text | |
| salinity_class | Code based on submitter field observation or measurement indicating average annual salinity | Categorical | |
| salinity_method | Indicate whether salinity_class was determined using a field observation or a measurement | Categorical | |
| salinity_notes | Any relevant submitter-generated notes on how salinity_class was determined | Free text | |
| vegetation_class | Code based on submitter field observations or measurements indicating dominant wetland vegetation type | Categorical | |
| vegetation_method | Indicate whether vegetation_class was determined using a field observation or a measurement | Categorical | |
| vegetation_notes | Any relevant submitter-generated notes on how vegetation_class and/or dominant species were determined | Free text | |
| inundation_class | Code based on submitter field observation or measurement indicating how often the coring location is inundated | Categorical | |
| inundation_method | Indicate whether inundation_class was determined using a field observation or a measurement | Categorical | |
| inundation_notes | Any relevant submitter-generated notes on how inundation was determined | Free text | |

- checking that the controlled attributes and variable names match guidelines.
- checking for the presence of mandatory and conditionally mandatory attributes (Data S1).
- verifying the uniqueness of core identifiers for a site and study.
- verifying the uniqueness of core locations in the overall database.
- verifying numeric attributes expressed as fractions were bound between 0 and 1.
- enforcing consistent spelling of taxonomic names using a global name resolver (Chamberlain & Szöcs, 2013).

During visual checks, Coastal Carbon Network personnel mapped the provided coordinates of sampling locations, the relationship between dry bulk density and organic matter, organic matter and organic carbon (Callaway et al., 2012; Craft et al., 1991; Morris et al., 2016), as well as depth-series of any depth profiles of soil measurements or radioisotope values (Arias-Ortiz et al., 2018).

2.5 | Data post-processing

We built a post-processing workflow that added geography attributes, detailed habitat classifications, and classified profiles based

on the data's utility, completeness, and quality. Following initial processing, cores were assigned geographic units, using spatial joins in the sf package (Pebesma, 2018) in R. Geographic units include countries (ESRI Data and Maps, 2015b), internationally recognized exclusive economic zones (Flanders Marine Institute, 2019) and level-1 administrative districts such as states and provinces (ESRI Data and Maps, 2015a).

Explicit habitat classification was not a part of the initial versions of this database, only descriptions of vegetation type (emergent, scrub/shrub, and forested) and salinity, as is compatible with the U.S. National Wetlands Inventory (FGDC Wetlands Subcommittee, 2009). In response to community demand, we provided a consistent classification of coastal ecosystems (e.g., marshes, mangroves, swamps, seagrasses, and scrub-shrub wetlands). This information was typically present but spread across multiple linked tables such as the core and species table, and multiple attributes such as vegetation_class, salinity_class, core_id, or site_id within the core table. For the version of the Library presented herein, we improved the classification of blue carbon habitats by writing a decision-making workflow to classify each core based on contextual study-specific information, salinity and vegetation descriptions, presence of indicator species, and author-provided core and site descriptions:

TABLE 2 Attribute information for the core table.

| Attribute name | Definition | Data type | Unit |
|--------------------|---|-------------|-------------------------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text | |
| site_id | Site identification code unique to each study | Free text | |
| core_id | Core identification code unique to each site | Free text | |
| year | Year of collection | Numeric | Numerical year |
| month | Numeric month of year of collection | Numeric | Numerical month of year |
| day | Numeric day of month of collection | Numeric | Numerical day of month |
| core_notes | Any other relevant submitter-generated notes on how cores were collected. | Free text | |
| latitude | Positional latitude of the core in decimal degree WGS84 | Numeric | Degree |
| longitude | Positional longitude of the core in decimal degree WGS84 | Numeric | Degree |
| position_accuracy | Accuracy of latitude and longitude measurement, if determined and recorded | Numeric | Meter |
| position_method | Code indicating how latitude and longitude were determined | Categorical | |
| position_notes | Any relevant submitter generated notes on how latitude and longitude were determined | Free text | |
| elevation | Surface elevation of the core relative to defined datum | Numeric | Meter |
| elevation_datum | The datum relative to which the core elevation was measured against (For a complete list of datum names and aliases please refer to the ISO Geodetic Registry https://iso.registry.bespire.eu/register/geodetic/VerticalDatum) | Categorical | |
| elevation_accuracy | Accuracy of elevation measurement, if determined and recorded | Numeric | Meter |
| elevation_method | Code indicating how elevation was determined | Categorical | |
| elevation_notes | Any relevant submitter-generated notes on how elevation was determined | Free text | |
| salinity_class | Code based on submitter field observation or measurement indicating average annual salinity | Categorical | |
| salinity_method | Indicates whether salinity_class was determined using a field observation or a measurement | Categorical | |
| salinity_notes | Any relevant submitter-generated notes on how salinity_class was determined | Free text | |
| vegetation_class | Code based on submitter field observations or measurement indicating dominant wetland vegetation type | Categorical | |
| vegetation_method | Indicates whether vegetation_class was determined using a field observation or a measurement | Categorical | |
| vegetation_notes | Any relevant submitter-generated notes on how vegetation_class and/or dominant species were determined | Free text | |
| habitat | Habitat classification of the sampled location based on description or dominant vegetation | Categorical | |
| inundation_class | Code based on submitter field observation or measurement indicating how often the coring location is inundated | Categorical | |
| inundation_method | Indicates whether inundation_class was determined using a field observation or a measurement | Categorical | |
| inundation_notes | Any relevant submitter-generated notes on how inundation was determined | Free text | |
| core_length_flag | Indicates whether or not the coring team believes they recovered a full sediment profile, down to bedrock, or other non-marsh interface | Categorical | |
| stocks_qual_code | Code indicating the quality of information present for a core containing data relevant for carbon stock assessment | Categorical | |



TABLE 2 (Continued)

| Attribute name | Definition | Data type | Unit |
|---------------------|--|-------------|------------|
| dates_qual_code | Code indicating the quality of information present for a core containing dated stratigraphy | Categorical | |
| elevation_qual_code | Code indicating the quality of information present for a dated core that contains elevation data | Categorical | |
| max_depth | Maximum depth of the sampled soil profile | Numeric | Centimeter |
| country | Country in which data were collected | Free text | |
| admin_division | Administrative division in which data were collected | Free text | |

1. We detected whether the entry originates from a habitat-specific synthesis for mangroves (Sanderman, 2017) or seagrasses (Fourqurean et al., 2012).
2. Non-forested wetlands were classified into marshes, seagrasses, or tidal flat based on vegetation_class.
3. If the vegetation class was not descriptive, then we cross-referenced species present at the coring location with the USDA Plants database (USDA, NRCS, 2022) to determine the growth form of the plant. If multiple growth forms were reported for a species, then we classified it based on the tallest reported. We used habitat-specific lists of taxa (Larkum et al., 2006; Tomlinson, 2016) to determine whether or not a species had any special designation as seagrass or mangrove. Any graminoids, forbs, or ferns not otherwise classified as seagrasses were classified as marshes. Any trees not otherwise classified as mangroves were classified as swamps.
4. If previous steps were inconclusive, we used partial string matching to detect whether any descriptive language such as 'mangrove' or 'seagrass' was present in site_id or core_id.
5. A small number of estuarine forested wetland locations from the United States (Nahlik & Fennessy, 2016) could not be sorted into swamps and mangroves using the previously outlined methods. We assumed sites were mangrove if they occurred south of 29.75° latitude, the maximum northern extent of mangroves observed on the U.S. Atlantic coast of the U.S. (Cavanaugh et al., 2014). Conversely, we assumed estuarine forested sites north of this threshold were tidal swamps.

We developed an automated workflow to classify cores based on the type of study the data can be used for, including carbon stock assessments, calculation of carbon burial rates, and fitting of models of future carbon sequestration and wetland resilience. If dry bulk density and either organic matter or organic carbon were present in-depth series, then the core meets the minimal inclusion criteria for carbon stocks (C). If the core was confirmed to reach the bottom of the profile then it was considered a high-quality core (C1). If not, it was classified as of lower utility (C2). Here, the wetland sediment profile is considered complete if the contact point between wetland sediment and bedrock or non-wetland sediment is reached.

If any profile age-related or disaggregated data associated with these techniques were present in a core, then the core met the minimal inclusion criteria for calculating carbon burial rates (B). If any

radioisotope activity levels were reported, then we required that the associated instrument error be reported for dating information to be considered complete (B1). This included ^{137}Cs , total ^{210}Pb , excess ^{210}Pb , ^{14}C , as well as any isotopes used to estimate supported ^{210}Pb , ^{226}Ra via its proxies ^{214}Bi and/or ^{214}Pb in gamma analysis. If the depth of ^{137}Cs peaks were reported, then we required a ^{137}Cs activity depth profile for completeness. If excess ^{210}Pb was reported, then we required that total ^{210}Pb be reported as well for the dataset to be considered complete. For ^{14}C ages, the material dated needed to be specified for the dataset to be considered complete. Any missing radioisotope errors or conditional attributes resulted in the dataset being classified as having incomplete data reporting (B2).

Cores with any age-depth data and elevation data met the minimum inclusion criteria for accretion modeling (A; Schile et al., 2014). We differentiated between elevations that were interpolated from remotely sensed or spatially interpolated digital elevation models (A3) and those that were directly measured using precise real-time kinematic GPS data or better (A1 or A2), which typically only have errors of a few centimeters. Cores with both a full suite of dating information and precise elevation were classified as the highest utility for data-model integration (A1), and cores with precise elevation but some missing dating information were graded lower (A2).

2.6 | Versioning, data use policy, and citation

The version of the database presented herein is v.1.0.0. and current as of January 12, 2023 (Coastal Carbon Network, 2023). Preliminary versions of this database were summarized in previous publications (Malhotra et al., 2019; Todd-Brown et al., 2022). New versions of this database will adopt standardized semantic versioning with the first digit indicating major changes to database structure, the second minor changes and additions of new datasets, and the third, backward-compatible changes. We anticipate regular updates to the Coastal Carbon Library as new datasets are submitted, approximately quarterly. It has a digital object identifier and a web link that will automatically route to the most up-to-date version.

The Coastal Carbon Library is licensed under CC-BY-4, meaning that it can be used without restriction other than attribution. Full attribution of the Coastal Carbon Library requires three things, first, citing the original primary datasets, papers or data publications, when they are reused for other purposes. In cases which an original

TABLE 3 Attribute information for depth-series table.

| Attribute name | Definition | Data type | Unit |
|--------------------------|---|-----------|-------------------------------------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text | |
| site_id | Site identification code unique to each study | Free text | |
| core_id | Core identification code unique to each site | Free text | |
| method_id | Unique identification code used to distinguish cores or sampling intervals that were collected or processed using different methods | Free text | |
| depth_min | Minimum depth of a sampling increment | Numeric | Centimeter |
| depth_max | Maximum depth of a sampling increment | Numeric | Centimeter |
| representative_depth_min | Minimum depth of a soil horizon of which the sample is representative | Numeric | Centimeter |
| representative_depth_max | Maximum depth of a soil horizon of which the sample is representative | Numeric | Centimeter |
| sample_id | This is a sample identification unique to the core. This should be used in the case that there are relevant lab specific sample codes, or in the case that there are multiple replicate samples. | Free text | |
| dry_bulk_density | Dry mass per unit volume of a soil sample. This does not include ash-free bulk density | Numeric | Grams per cubic centimeter |
| fraction_organic_matter | This is the mass of organic matter relative to sample dry mass. Ash-free bulk density should not be used here but should be expressed as a loss on ignition fraction. | Numeric | Dimensionless |
| fraction_carbon | Mass of carbon relative to sample dry mass | Numeric | Dimensionless |
| compaction_fraction | Fraction of the sample depth interval reduced due to compaction | Numeric | Dimensionless |
| compaction_notes | Any submitter generated notes on compaction | Free text | |
| cs137_peak_age | Age of documented radiocesium (^{137}Cs) peak | Numeric | Year |
| cs137_activity | Radioactivity counts per unit dry weight for ^{137}Cs | Numeric | Specified in ^{137}Cs unit |
| cs137_activity_se | 1 standard error of uncertainty associated with cs137_activity | Numeric | Specified in ^{137}Cs unit |
| cs137_unit | Reported unit for sample interval's ^{137}Cs activity measurements | Free text | Specified in ^{137}Cs unit |
| excess_pb210_activity | Excess radioactivity counts per unit dry weight for excess lead 210 (^{210}Pb) | Numeric | Specified in ^{210}pb unit |
| excess_pb210_activity_se | 1 standard error of uncertainty associated with excess_pb210_activity | Numeric | Specified in ^{210}pb unit |
| total_pb210_activity | Total radioactivity counts per unit dry weight for excess ^{210}Pb | Numeric | Specified in ^{210}pb unit |
| total_pb210_activity_se | 1 standard error of uncertainty associated with total_pb210_activity | Numeric | Specified in ^{210}pb unit |
| pb210_unit | Reported unit for sample interval's ^{210}Pb activity measurements | Free text | |
| ra226_activity | This is the total radioactivity counts per unit dry weight for Radium-226 (^{226}Ra), also referred to as supported ^{210}Pb . This can be measured either using the proxy lead-214 (^{214}Pb), bismuth-214 (^{214}Bi), or the two of them averaged. Authors also have the option to report ^{214}Pb and ^{214}Bi in separately under pb214_activity and bi214_activity respectively. | Numeric | Specified in ^{226}ra unit |
| ra226_activity_se | 1 standard error of uncertainty associated with ra226_activity | Numeric | Specified in ^{226}ra unit |
| ra226_unit | Reported unit for sample interval's ^{226}Ra activity measurements | Free text | Specified in ^{226}ra unit |
| pb214_activity | This is the radioactivity counts per unit dry weight for lead-214 (^{214}Pb), a proxy for Radium-226 (^{226}Ra), also referred to as supported ^{210}Pb . Radioactivity should be reported here, rather than ra226_activity, either if the author also measures bismuth-214 and ra226_activity is a composite of the two measurements, or if they want to specify the proxy used for ^{226}Ra . | Numeric | Specified in ^{214}pb unit |
| pb214_activity_se | Estimated uncertainty in pb214_activity | Numeric | Specified in ^{214}pb unit |
| pb214_unit | Reported unit for sample interval's ^{214}Pb activity measurements | Free text | |

TABLE 3 (Continued)

| Attribute name | Definition | Data type | Unit |
|----------------------|--|-------------|-------------------------------------|
| bi214_activity | This is the radioactivity counts per unit dry weight for bismuth-214 (^{214}Bi), a proxy for Radium-226 (^{226}Ra), also referred to as supported ^{210}Pb . Radioactivity should be reported here, rather than ra226_activity , either if the author also measures lead-214 and ra226_activity is a composite of the two measurements, or if they want to specify the proxy used for ^{226}Ra . | Numeric | Specified in ^{214}Bi unit |
| bi214_activity_se | Reported standard error for radioactivity counts per unit dry weight for ^{214}Bi | Numeric | Specified in ^{214}Bi unit |
| bi214_unit | Reported unit for sample interval's ^{214}Bi activity measurements. | Free text | |
| c14_age | Radiocarbon (^{14}C) age as estimated from AMS measurements | Numeric | Radiocarbon year |
| c14_age_se | Estimated uncertainty in c14_age | Numeric | Radiocarbon year |
| c14_material | Description of the material selected for ^{14}C dating | Free text | |
| c14_notes | Any relevant submitter generated notes on ^{14}C dating process | Free text | |
| delta_c13 | The isotopic signature of ^{13}C , oftentimes measured along with ^{14}C age and can be useful for analyzing carbon liability and provenance | Numeric | Parts per million |
| be7_activity | Radioactivity counts per unit dry weight for beryllium-7 (^{7}Be) | Numeric | Specified in ^{7}Be unit |
| be7_activity_se | Estimated uncertainty in be_7_activity | Numeric | Specified in ^{7}Be unit |
| be7_unit | Reported unit for sample interval's ^{7}Be activity measurements | Free text | |
| marker_date | The age of any other dated depth horizon such as an artificial marker, pollen horizon, pollution horizon, etc. | Date | Numerical year |
| marker_date_se | The standard error of the age of any other dated depth horizon such as an artificial marker, pollen horizon, pollution horizon, etc. | Date | |
| marker_type | Code indicating the type of marker | Categorical | |
| marker_notes | Any other submitter-generated notes about the origin of the marker | Free text | |
| age | Most likely, median, or mean age of the depth interval from submitter generated age-depth model | Numeric | Numerical year |
| age_min | Minimum age of the depth interval from submitter generated age-depth model | Numeric | Numerical year |
| age_max | Maximum age of the depth interval from submitter generated age-depth model | Numeric | Numerical year |
| age_se | Standard error of age estimate from submitter generated age-depth model | Numeric | Numerical year |
| depth_interval_notes | Any other submitter-generated notes specific to the depth interval | Free text | |

source entered the Coastal Carbon Library via another synthesis for example, Sanderman et al. (2018) and Fourqurean et al. (2012), both the original source and synthesis should be cited. Second and third, the Coastal Carbon Library itself (Coastal Carbon Network, 2023) and, this manuscript should be cited in addition to primary and synthesis sources. Importantly, when data are downloaded and analyzed for new purposes, original primary sources need to always be cited to comply with the policy; citation of the Coastal Carbon Library and this paper alone would not be sufficient.

Because relationships between original data sources and the Coastal Carbon Library are complex, we include a relational table connecting study_id's to bibliographic information (Figure 1). The Coastal Carbon Altas (Figure 3), automatically formats bibliographic information based on the subset of datasets downloaded.

2.7 | Quantification of data coverage

We compared the makeup of cores in the Coastal Carbon Library to the global allocation of blue carbon area based on habitat and countries. For habitats, we reference literature for global area estimates

for seagrasses, tidal marshes, and mangroves (Bunting et al., 2022; McKenzie et al., 2020; Murray et al., 2022; Table 9). For countries, we used a probabilistic map of intertidal area (Murray et al., 2022), counting tidal wetland pixels with greater than a 50% chance of being classified as tidal flat, tidal marsh, or mangrove in 1999. Country borders included both the land borders (ESRI Data and Maps, 2015b) and exclusive economic zones (Flanders Marine Institute, 2019).

3 | ACCESSING DATA VIA THE COASTAL CARBON ATLAS

The Coastal Carbon Atlas (https://shiny.si.edu/coastal_carbon_atlas; Figure 3) is an R-shiny web application that interfaces with the Coastal Carbon Library allowing users to explore, query, and directly download data. This tool has been adapted to suit user needs through community feedback. The Coastal Carbon Atlas consists of a map interface (Figure 3a) displaying sampling locations. The tabular view allows for a direct visual review of the site, core, and depth series tables (Figure 3b). The Coastal Carbon Atlas allows for the sub-setting of datasets by habitat type, geography, data availability,

TABLE 4 Attribute information for methods table.

| Attribute name | Definition | Data type | Unit |
|--------------------------------|--|-------------|------------------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text | |
| method_id | Unique identification code used to distinguish cores or sampling intervals that were collected or processed using different methods | Free text | |
| coring_method | Code indicating what type of device was used to collect soil depth profiles | Categorical | |
| roots_flag | Code indicating whether live roots were included or excluded from carbon assessments | Categorical | |
| sediment_sieved_flag | Code indicating whether or not sediment was sieved prior to carbon measurements | Categorical | |
| sediment_sieve_size | If sediment was sieved, the size of sieve used | Numeric | Millimeters |
| compaction_flag | Code indicating how the authors qualified or quantified compaction of the core | Categorical | |
| dry_bulk_density_temperature | This is the temperature at which samples were dried to measure dry bulk density. This can include either samples that were freeze dried or oven dried. | Numeric | Degree celsius |
| dry_bulk_density_time | Time over which samples were dried to measure dry bulk density | Numeric | Hour |
| dry_bulk_density_sample_volume | Sample volume used for bulk density measurements, if held constant | Numeric | Cubic centimeter |
| dry_bulk_density_sample_mass | Sample mass used for bulk density measurements, if held constant | Numeric | Gram |
| dry_bulk_density_flag | Any notable codes regarding how the authors quantified dry bulk density | Categorical | |
| loss_on_ignition_temperature | Temperature at which samples were combusted to estimate fraction organic matter | Numeric | Degree celsius |
| loss_on_ignition_time | Time over which samples were combusted to estimate fraction organic matter | Numeric | Hour |
| loss_on_ignition_sample_volume | Sample volume used for loss on ignition, if held constant | Numeric | Cubic centimeter |
| loss_on_ignition_sample_mass | Sample mass used for loss on ignition, if held constant | Numeric | Gram |
| loss_on_ignition_flag | Common codes regarding loss on ignition methodology | Categorical | |
| carbonates_removed | Whether or not carbonates were removed prior to calculating fraction organic carbon | Categorical | |
| carbonate_removal_method | The method used to remove carbonates prior to measuring fraction carbon | Categorical | |
| fraction_carbon_method | Code indicating the method for which fraction carbon was measured | Categorical | |
| fraction_carbon_type | Code indicating whether fraction_carbon refers to organic or total carbon | Categorical | |
| carbon_profile_notes | Any other submitter-defined notes describing methodologies for determining dry bulk density, organic matter fraction, and carbon fraction | Free text | |
| pb210_counting_method | Code indicating the method used for determining lead 210 activity | Categorical | |
| excess_pb210_rate | Code indicating the mass or accretion rate used in the excess_pb_210_model | Categorical | |
| excess_pb210_model | Code indicating the model used to estimate accumulation rates and/or age-depth chronologies | Categorical | |
| ra226_assumption | Code indicating the assumption used to estimate the core's supported radium-226-specific activities | Categorical | |
| c14_counting_method | Code indicating the method used for determining radiocarbon activity | Categorical | |
| dating_notes | Any submitter defined notes elaborating on the process of dating the core not yet made clear by the coding | Free text | |
| age_depth_model_reference | Code indicating the reference or 0 year of the age depth model | Categorical | |
| age_depth_model_notes | Any submitter defined notes on how the age depth model was created | Free text | |

TABLE 5 Attribute information for species table.

| Attribute name | Definition | Data type |
|----------------|--|-------------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text |
| site_id | Site identification code unique to each study | Free text |
| core_id | Core identification code unique to each site | Free text |
| species_code | Code associated with a species or a vegetation assemblage | Free text |
| code_type | Defines whether the code refers to a description, or level of plant taxonomy | Categorical |
| habitat | Habitat classification of the sampled location based on description or dominate vegetation | Categorical |

TABLE 6 Attribute information for impacts table.

| Attribute name | Definition | Data type |
|----------------|--|-------------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text |
| site_id | Site identification code unique to each study | Free text |
| core_id | Core identification code unique to each site | Free text |
| impact_class | Code indicating any major anthropogenic impacts historically and currently affecting the coring location | Categorical |

core depth, study, human impacts, plant species present, and salinity descriptor. Finally, the Coastal Carbon Atlas provides options for limited post-processing and for downloading the data (Figure 3c). The user can specify custom depth intervals to download quantitative attributes summarized as depth-weighted averages of the depth increments specified.

Data downloads (Figure 3d) are accompanied by a bibliography of original sources and associated publications, metadata, and a text file that outlines the data use policy. To facilitate data users in providing credit to original sources, bibliographic information is included in a text file with formatted citations, a BibTex file that can be imported into reference management software, and a comma-separated value file with disaggregated reference information.

4 | CURRENT STATE OF THE COASTAL CARBON LIBRARY

The Coastal Carbon Library synthesis contained 6723 coastal wetland soil profiles from around the globe. These represent 48,870 depth increments, and 126,629 individual measurements, from 64 countries (Figures 3 and 4). Overall there were 305 studies represented in the database, related to 299 bibliographic entities (Abbott, Elsey-Quirk, & DeLaune, 2019; Abbott, Quirk, & Delaune, 2019; Abed-Navandi & Dworschak, 2005; Abino et al., 2014; Adame et al., 2013, 2015; Agawin et al., 1996; Ajonina et al., 2014; Allen et al., 2007, 2021, 2022; Alongi, 1996; Alongi et al., 1999, 2001, 2004, 2005; Alongi, Tirendi, Clough, 2000; Alongi, Tirendi, Trott, et al., 2000; Alongi, Trott, Rachmansyah, et al., 2008; Alongi, Trott, Undu, et al., 2008; Al-rousan et al., 2005; Amon & Herndl, 1991; Andreetta et al., 2014, 2016; Arias-Ortiz, Masqué, et al., 2021; Arias-Ortiz, Oikawa, et al., 2021; Arriola & Cable, 2017; Ashton & Macintosh, 2002; Banerjee et al., 2012;

Barrón et al., 2004; Baustian, Stagg, Perry, Moss, & Carruthers, 2021; Baustian, Stagg, Perry, Moss, Carruthers, et al., 2021; Belshe et al., 2019a, 2019b; Bernhardt et al., 2018; Borg et al., 2010; Boschker et al., 2000; Bouillon et al., 2003; Boyd, 2012; Boyd et al., 2017, 2019; Boyd & Sommerfield, 2016; Breithaupt et al., 2014, 2017, 2019, 2020; Brunskill et al., 2002, 2004; Buffington et al., 2020; Bukoski et al., 2017; Bulmer et al., 2016; Burden et al., 2018, 2019; Burns & Swart, 1992; Buzzelli, 1998; Cahoon & Lynch, 1997; Callaway et al., 1997, 2012, 2019; Calleja et al., 2007; Campbell et al., 2014; Cardona & Botero, 1998; Carlin et al., 2021; Carruthers et al., 2005; Chambers et al., 2019; Chanda et al., 2013; Chen et al., 2010, 2012, 2014, 2016; Chen & Twilley, 1999; Chmura et al., 2003; Coastal Protection and Restoration Authority, 2015; Cochran et al., 1998; Cotner et al., 2004; Craft, 2007; Crooks et al., 2014; da Silva et al., 2009; Danovaro, 1996; Danovaro et al., 1994; Danovaro & Fabiano, 1995; Danovaro & Gambi, 2002; De Falco et al., 2006; de Jongh et al., 1995; De Troch et al., 2006; DelVecchia et al., 2014; Devereux et al., 2011; Dissanayake & Chandrasekara, 2014; Donato et al., 2011, 2012; Doughty et al., 2015, 2019; Drake et al., 2015; Drexler et al., 2009, 2013, 2019; Dung et al., 2016; Dutta et al., 2013; Elsey-Quirk et al., 2011; Ensign et al., 2015, 2021; EPA, 2016a, 2016b; Erftemeijer & Middelburg, 1993; Eyre & Ferguson, 2002; Ezcurra et al., 2016; Fell et al., 2021; Ferreira et al., 2007; Fonseca et al., 2011; Fourqurean et al., 2010, 2012; Fujimoto et al., 1999; Gacia et al., 2002; Gerlach et al., 2017; Giblin et al., 2018; Gillis et al., 2017; Gleason & Ewel, 2002; Gomes et al., 2016; Gonzea et al., 2004, 2018; Grady, 1981; Grenz et al., 2003; He et al., 2002; Hebert et al., 2006; Hemminga et al., 1994; Hill & Anisfeld, 2015; Holmer et al., 2001, 2006, 2009; Holmer & Duarte, 2003; Holmer & Frederiksen, 2007; Holmquist et al., 2018a; Howe et al., 2009; Iacono et al., 2008; Isaksen & Finster, 1996; Jennerjahn & Ittekkot, 2002; Johnson et al., 2007; Jones et al., 2016, 2017; Kairis & Rybcyzk, 2010; Kamp-Nielsen et al., 2002; Kao et al., 2002; Kauffman et al., 2020a,

TABLE 7 Attribute information for bibliography table.

| Attribute name | Definition | Data type |
|------------------|--|-------------|
| study_id | Unique identifier for the study made up of the first author's family name, as well as the second author's family name or et al. if more than three, then publication year separated by underscores | Free text |
| bibliography_id | Bibliography identification code unique to each study | Free text |
| publication_type | Code indicating the type of publication the study originates from | Categorical |
| bibtype | BibTeX entry type | Free text |
| title | The title of the work | Free text |
| author | The author(s) of the publication | Free text |
| school | The school where the thesis was written | Free text |
| doi | Digital object identifier associated with the work | Free text |
| url | Permanent web address where the work can be located | Free text |
| year | The year of publication or, if unpublished, the year of creation | Free text |
| month | The month of publication (or, if unpublished, the month of creation) | Free text |
| publisher | The publisher's name | Free text |
| volume | The volume of a journal or multi-volume book | Free text |
| number | The (issue) number of a journal, magazine, or tech-report, if applicable (most publications have a volume, but no number field) | Free text |
| pages | Page numbers, separated either by commas or double-hyphens | Free text |
| journal | The journal or magazine the work was published in | Free text |
| copyright | The intellectual rights of the publication | Free text |
| editor | The name(s) of the editor(s) | Free text |
| booktitle | The title of the book, if only part of it is being cited | Free text |
| institution | The institution that was involved in the publishing, but not necessarily the publisher | Free text |
| language | The language a publication was written in | Free text |

2020b; Kemp et al., 2012, 2020; Kenig et al., 1990; Keshta et al., 2017, 2020, 2021; Keuskamp et al., 2013; Koch & Madden, 2001; Koepfler et al., 1993; Krause-Jensen et al., 2011; Krauss et al., 2018; Kristensen et al., 1992, 2000; Kulawardhana et al., 2015; Lacerda et al., 1995; Lagomasino et al., 2013, 2020; Lallier-Verges et al., 1998; Lang et al., 2014; Larned, 2003; Leduc & Probert, 2010; Lee et al., 2005; Lee & Baker, 1972; Leopold et al., 2013; Lewis et al., 2007; Lillebø et al., 2006; Liu et al., 2008, 2014; Livesley & Andrusiak, 2012; Loi & Ni, 2015; Lovelock et al., 2010; Luk et al., 2020, 2021; MacKenzie et al., 2016; Maher & Eyre, 2010; Mao et al., 2011, 2012; Marchand et al., 2003, 2004; Marchio et al., 2016; Mateo et al., 1997; Mateo & Romero, 1997; Matsui and Naohiro, 1998; McClellan, 2021; McClellan et al., 2021; McFadden et al., 2016; McGlathery et al., 2012; McTigue et al., 2019, 2020; Mellors et al., 2002; Merrill, 1999; Messerschmidt & Kirwan, 2020; Mfilinge et al., 2002; Middelburg et al., 1996; Miller et al., 2022; Miyajima et al., 1998; Murdiyarsa et al., 2015; Nahlik & Fennessy, 2016; Naidoo, 1980; Naidoo & Raiman, 1982; Nam et al., 2016; Neubauer et al., 2002; Noe et al., 2012, 2016; Nolte, 2020; Nolte et al., 2013; Nsommbo et al., 2016; Nuttle, 1996; Nyman et al., 1993; Oakes & Connolly, 2004; O'keefe Suttles, Eagle, Mann, & Kroeger, 2021; O'keefe Suttles, Eagle, Mann, Moseman-Valtierra, et al., 2021; O'keefe Suttles, Eagle, Mann, Spivak, et al., 2021; O'keefe Suttles, Eagle, Mann, Wang, et al., 2021; O'keefe Suttles, Wigand, Eagle, Branoff, et al., 2021; Ooi et al., 2011; Orem et al., 1999; Orson & Simpson, 1990; Osland, 2017; Osland et al., 2012; Otero et al., 2009;

Pastore et al., 2017; Paula et al., 2001; Pazi et al., 2016; Peck et al., 2020a, 2020b; Pedersen et al., 1997; Piazza et al., 2011; Poppe, 2015; Poppe & Rybczyk, 2018, 2019, 2021, 2022; Prentice et al., 2020; Pulich Jr., 1987; Qu et al., 2006; Radabaugh et al., 2017; Rigollet, 2004; Rodriguez & Molly, 2022; Rosenfeld, 1979; Sahu et al., 2016; Saintilan et al., 2013; Sanbor & Darwyn, 2020; Sanderman, 2017; Sanders et al., 2008, 2014; Schile & Megonigal, 2017; Sfriso & Marcomini, 1999; Smith, 2012; Smith et al., 2015; Smoak et al., 2013; Spivak, 2020; Spivak et al., 2009; Spruzen et al., 2007; St. Laurent et al., 2020a, 2020b; Stoner et al., 1998; Stringer et al., 2015; Tam & Wong, 1998; Thimdee et al., 2003; Thom, 1992, 2019; Thompson et al., 2014; Thorne et al., 2018; Townsend & Fonseca, 1998; Trettin et al., 2017; Tue et al., 2014; Twilley et al., 1997; Unger et al., 2016; U.S. Geological Survey and Thorne, 2015; Van Engeland, 2010; Vasconcelos et al., 2014; Vaughn et al., 2020, 2021; Vichkovitten & Holmer, 2005; Volkman et al., 2008; Wang et al., 2013; Ward, 2021; Ward et al., 2021; Watson & Byrne, 2013; Weis et al., 2001; Weis & Anthony, 1999; Weston et al., 2022; White et al., 2020; Wigand et al., 2021; Windham-Myers et al., 2010; Woodroffe, 1985; Wooller et al., 2003; Xia et al., 2015; Xu et al., 2012; Yamamuro et al., 1993; Yando et al., 2016; Yang et al., 2013; Yarbro & Carlson, 2008; Zhang et al., 2012).

The majority of soil profiles came from tidal marsh (43.7%), followed by mangrove (29.5%), and seagrass ecosystems (11.7%). Lower proportions were from unvegetated surfaces (8.5%), not specified

TABLE 8 Definitions of controlled vocabulary for attributes with categorical data.

| Attribute name | Variable name | Definition |
|-----------------------|---------------------------------|--|
| corresponding_author | TRUE | The author should be contacted with any further questions. |
| corresponding_author | FALSE | The author should not be contacted with any further questions. |
| publication_type | Combined dataset and manuscript | Primary dataset and associated manuscript are from the same source. |
| publication_type | Primary dataset | Primary published dataset |
| publication_type | Associated source | Source associated with primary dataset |
| publication_type | Synthesis dataset | Published dataset associated with the data synthesis |
| publication_type | Synthesis source | Source associated with data synthesis contribution |
| coring_method | Cryo core | A technique involving collecting a core by freezing soil using liquid nitrogen to a copper tube |
| coring_method | Gouge auger | A half cylinder coring device in which the coring section is open, not sealed off by a fin |
| coring_method | Hargas corer | A large diameter (>10 cm) coring device consisting of a tube, piston, and a cutting head |
| coring_method | Mcauley corer | A half cylinder coring device with the coring section sealed off by a fin attached to a rotating pivot point |
| coring_method | Mccaffrey peat cutter | U-shaped blade that extracts a core by cutting down through peat |
| coring_method | None specified | No coring device was specified |
| coring_method | Other shallow corer | Any other type of coring device typically taking cores shallower than 30 cm |
| coring_method | Piston corer | A device that extrudes core into tube upward with a plunger |
| coring_method | Push core | Any number of coring types involving driving a tube into the sediment to recover a core |
| coring_method | Pvc and hammer | PVC pipe was driven into the sediment with a hammer to recover a core |
| coring_method | Russian corer | A half cylinder coring device with the coring section sealed off by a fin attached to a rotating pivot point |
| coring_method | Vibracore | A technique involving collecting a core by sinking a continuous pipe into sediment attaching a source of vibration, then recovering using a winch and pulley |
| coring_method | Surface sample | A technique involving collecting a core shallower than ~5 cm using a circular metal cutter |
| coring_method | Soil pit | A pit is dug from the surface of the soil to the underlying bedrock |
| roots_flag | Roots and rhizomes included | Roots and rhizomes were included in dry bulk density and or organic matter and carbon measurements |
| roots_flag | Roots and rhizomes separated | Roots and rhizomes were separated from soil before dry bulk density and or organic matter and carbon measurements |
| sediment_sieved_flag | Sediment sieved | Sediment was sieved prior to analysis for organics |
| sediment_sieved_flag | Sediment not sieved | Sediment was not sieved prior to analysis for organics |
| compaction_flag | Compaction qualified | Compaction was at least qualified and noted by the authors |
| compaction_flag | Compaction quantified | Compaction was quantified and corrected for in core based measurements |
| compaction_flag | Corer limits compaction | Authors specified that the coring device's design minimized compaction |
| compaction_flag | No obvious compaction | Authors observed no obvious compaction |
| compaction_flag | Not specified | Compaction was not specified |
| dry_bulk_density_flag | Air dried to constant mass | Methodology specified that samples were air dried to a constant mass |
| dry_bulk_density_flag | Freeze dried | Bulk density was measured on freeze dried samples |
| dry_bulk_density_flag | Not specified | No additional details regarding bulk density methodology were provided |
| dry_bulk_density_flag | Removed non structural water | Bulk density methodology did not specify drying temperature or length, only that non-structural water was removed |
| dry_bulk_density_flag | Time approximate | Bulk density time recorded herein is an approximate estimate |
| dry_bulk_density_flag | To constant mass | Bulk density methodology did not specify drying temperature or length, only that samples were dried to a constant mass |

(Continues)

TABLE 8 (Continued)

| Attribute name | Variable name | Definition |
|---------------------------|------------------------|--|
| loss_on_ignition_flag | Time approximate | Loss on ignition time recorded herein is an approximate estimate |
| loss_on_ignition_flag | Not specified | No additional details regarding loss on ignition methodology or time were provided |
| carbonates_removed | FALSE | Carbonates were not removed before measuring organic carbon |
| carbonates_removed | TRUE | Carbonates were removed before measuring organic carbon |
| carbonate_removal_method | Direct acid treatment | Carbonates were removed using direct application of dilute acid |
| carbonate_removal_method | Acid fumigation | Carbonates were removed by fumigating with concentrated acid |
| carbonate_removal_method | Low carbonate soil | Organic carbon fraction was measured without removing carbonates assuming carbonate content of the soil type was minimal |
| carbonate_removal_method | Carbonates not removed | Carbonates were not removed and low carbonate soil was not specified |
| carbonate_removal_method | None specified | Carbonate removal methodology was not specified |
| fraction_carbon_method | EA | Each sample presented was measured using Elemental Analysis |
| fraction_carbon_method | Kjeldahl digestion | Each sample was measured kjeldahl digestion method |
| fraction_carbon_method | Not specified | No additional details were provided regarding fraction carbon methodologies |
| fraction_carbon_method | Wet oxidation | Each sample was measured using a wet oxidation method |
| fraction_carbon_type | Organic carbon | Author specified that fraction carbon measurements were of organic carbon |
| fraction_carbon_type | Total carbon | Author specified that fraction carbon measurements were of total carbon |
| pb210_counting_method | Alpha | Alpha counting method was used |
| pb210_counting_method | Gamma | Gamma counting method was used |
| excess_pb210_rate | Mass accumulation | Excess ^{210}Pb modeled using mass accumulation rate |
| excess_pb210_rate | Accretion | Excess ^{210}Pb modeled using vertical accretion rate |
| excess_pb210_rate | Cumulative mass | Model run as excess ^{210}Pb against cumulative mass |
| excess_pb210_rate | Depth | Model run as excess ^{210}Pb against depth |
| excess_pb210_model | CRS | Constant rate of supply model used to estimate mass and/or accretion rates from age profile or chronology |
| excess_pb210_model | CIC | Constant initial concentration model used |
| excess_pb210_model | CFCS | Constant flux constant sedimentation model used |
| ra226_assumption | Each sample | ^{226}Ra was measured for each sample |
| ra226_assumption | Total core | ^{226}Ra was measured for the total core |
| ra226_assumption | At asymptote | ^{226}Ra was measured at the asymptote of the ^{210}Pb profile |
| ra226_assumption | Selected samples | ^{226}Ra measured by spectrometry at selected samples along the soil profile |
| c14_counting_method | AMS | Accelerator mass spectroscopy used |
| c14_counting_method | Beta | Beta counting used |
| age_depth_model_reference | YBP | Year zero is defined as years before present, 1960 CE |
| age_depth_model_reference | CE | Year zero is set according to Common Era and Before Common Era standards |
| age_depth_model_reference | Core collection date | Year zero is set as the core's collection year |

TABLE 8 (Continued)

| Attribute name | Variable name | Definition |
|-------------------|---------------------------|---|
| salinity_class | Estuarine | 0.5–35 ppt |
| salinity_class | Brine | >50 ppt |
| salinity_class | Saline | 30–50 ppt |
| salinity_class | Brackish | 0.5–30 ppt |
| salinity_class | Fresh | <0.5 ppt |
| salinity_class | Mixoeuhaline | 30–40 ppt |
| salinity_class | Polyhaline | 18–30 ppt |
| salinity_class | Mesohaline | 5–18 ppt |
| salinity_class | Oligohaline | 0.5–5 ppt |
| salinity_class | Estuarine C-CAP | 5–35 parts per thousand salinity (ppt) according to the coastal change analysis program |
| salinity_class | Palustrine C-CAP | <5 ppt according to the coastal change analysis program |
| salinity_method | Field observation | Salinity inferred by field observation such as vegetation |
| salinity_method | Measurement | Salinity observed from local instrument |
| vegetation_class | Emergent | Describes wetlands dominated by persistent emergent vascular plants |
| vegetation_class | Scrub shrub | Describes wetlands dominated by woody vegetation < or equal to 5 m in height |
| vegetation_class | Forested | Describes wetlands dominated by woody vegetation >5 m in height |
| vegetation_class | Forested to shrub | Dominated by forested to scrub/shrub biomass |
| vegetation_class | Forested to emergent | Dominated by forest and underlying marsh |
| vegetation_class | Seagrass | Describes tidal or subtidal communities dominated by rooted vascular plants |
| vegetation_method | Field observation | Vegetation inferred by field observation |
| vegetation_method | Measurement | Vegetation measured by counts or plots. |
| inundation_class | High | Study-specific definition of an elevation relatively high in the tidal frame, typically defined by vegetation type |
| inundation_class | Mid | Study-specific definition of an elevation in the relative middle of the tidal frame, typically defined by vegetation type |
| inundation_class | Low | Study-specific definition of an elevation in relatively low in the tidal frame, typically defined by vegetation type |
| inundation_class | Levee | Study-specific definition of a relatively high elevation zone built up on the edge of a river, creek, or channel |
| inundation_class | Back | Study-specific definition of a relatively low elevation zone behind a levee |
| inundation_method | Field observation | Inundation inferred by field observation such as vegetation |
| inundation_method | Measurement | Inundation class assessed from elevation and nearby tide gauge or other similar method |
| position_method | RTK | Real-time kinematic global position system (GPS) |
| position_method | Handheld | Conventional Commercially available hand-held GPS |
| position_method | Other high resolution | Any other technique resulting in positional error <1 m |
| position_method | Other moderate resolution | Any other technique resulting in positional error <30 m |
| position_method | Other low resolution | Any other technique resulting in positional error >30 m |
| elevation_datum | NAVD88 | A gravity-based geodetic datum, North American Vertical Datum of 1988 |
| elevation_datum | MSL | A tidal datum, Mean Sea Level as measured against a local tide gauge |
| elevation_datum | MTL | A tidal datum, Mean Tidal Level as measured against a local tide gauge |
| elevation_datum | MHW | A tidal datum, Mean High Water as measured against a local tide gauge |
| elevation_datum | MHHW | A tidal datum, Mean Higher High Water as measured against a local tide gauge |
| elevation_datum | MHHWS | A tidal datum, Mean Higher High Water for Spring Tides as measured against a local tide gauge |
| elevation_datum | MLW | A tidal datum, Mean Low Water as measured against a local tide gauge |

(Continues)


TABLE 8 (Continued)

| Attribute name | Variable name | Definition |
|------------------|---------------------------------------|--|
| elevation_datum | MLLW | A tidal datum, Mean Lower Low Water as measured against a local tide gauge |
| elevation_datum | NHN | Normaal Amsterdams Peil, a vertical datum used in many areas of Western Europe |
| elevation_datum | NAP | Normalhoehennull, a vertical datum used in Germany that represents height above mean sea level |
| elevation_method | RTK | Real-time kinematic GPS |
| elevation_method | Other high resolution | Any other technique resulting in positional error <5 cm of random error |
| elevation_method | LiDAR | Handheld GPS matched to lidar-based digital elevation model |
| elevation_method | DEM | Handheld GPS matched to another digital elevation model |
| elevation_method | Other low resolution | Any other technique resulting in positional error >5 cm of random error |
| core_length_flag | Core depth limited by length of corer | The total depth of the core was limited by the length of the coring device |
| core_length_flag | Core depth represents deposit depth | Authors report that the depth of the core represents the depth of the wetland soil deposit. |
| core_length_flag | Not specified | Authors did not specify whether or not the depth of the core represents the depth of the wetland soil deposit |
| marker_type | Artificial horizon | Horizon was added to the surface artificially by using materials such as feldspar, glitter, or rare earth elements |
| marker_type | Pollen | Pollen analysis was used to tie horizon to the timing of vegetation change such as the arrival of invasives, or the beginning of local agriculture |
| marker_type | Pollution | Chemical analysis was used to tie the horizon to the timing of a pollution event |
| marker_type | Tsunami | Sediment analysis was used to tie the horizon to the timing of a tsunami event |
| impact_class | Tidally restricted | Tidal flow is muted or blocked by built structures |
| impact_class | Impounded | Water level is raised artificially by a tidal restriction, resulting in ponding of water on the wetland and or upland surface |
| impact_class | Salt impacted | Wetland has been salinized |
| impact_class | Natural | No disturbance or management has occurred |
| impact_class | Managed impounded | Wetland is impounded seasonally, and other times natural or semi natural hydrology occurs |
| impact_class | Ditched | Tidal hydrology is altered because artificial ditches have been cut to promote tidal flooding and drainage |
| impact_class | Diked and drained | The wetland has been diked and drained, with or without flapper gates, pumping, or other means |
| impact_class | Farmed | Managed impoundment or drainage in which wetland has been converted to agricultural land |
| impact_class | Tidally restored | Tidal flow has been restored by removing an artificial obstruction |
| impact_class | Revegetated | Wetland vegetation has been reintroduced by replanting on unvegetated surfaces |
| impact_class | Restored | Some method of restoration (potentially unspecified) has occurred |
| impact_class | Invasive plants removed | Natural plant communities have been restored by the active removal of invasive plant species |
| impact_class | Invasive herbivores removed | Tidal wetland vegetation has been managed by the removal of invasive herbivores |
| impact_class | Sediment added | Elevation has been managed by artificially adding sediment to the site using techniques such as thin layering or sediment diversion |
| impact_class | Wetlands built | Constructed wetland using sediments such as dredge spoils or other sediment source |
| habitat | Mangrove | Tropical and dominated by specialized shrubs or trees |
| habitat | Swamp | Predominantly freshwater and dominated by trees |
| habitat | Scrub/shrub | Wetland dominated by shrubs |

TABLE 8 (Continued)

| Attribute name | Variable name | Definition |
|---------------------|---------------|---|
| habitat | Marsh | Wetland dominated by emergent vegetation, graminoids or forbs |
| habitat | Seagrass | Intertidal to subtidal and dominated by specialized predominantly submerged grasses |
| habitat | Algal mat | Dominated by algae |
| habitat | Unvegetated | Unvegetated |
| habitat | Tidal flat | Describes unvegetated areas exposed and flooded by the tides |
| habitat | Upland | Predominately above sea level and dominated by trees |
| code_type | Description | Cover description not related to plant taxonomy |
| code_type | Family | Refers to taxonomic family without indicating a genus or species |
| code_type | Genus | Refers to a taxonomic genus without indicating a species |
| code_type | Genus species | Refers to taxonomic Genus and species |
| stocks_qual_code | C2 | Carbon stock data complete, not confirmed to be a complete profile |
| stocks_qual_code | C1 | Carbon stock data complete, confirmed to be a complete profile |
| dates_qual_code | B2 | Dating information present, but not complete |
| dates_qual_code | B1 | Dating information present and complete |
| elevation_qual_code | A3 | Elevation data present but of low quality, dating info present |
| elevation_qual_code | A2 | Elevation data are of high quality, and dating info present but incomplete |
| elevation_qual_code | A1 | Elevation data are of high quality, and dating info is complete |

FIGURE 2 A flowchart shows the interactions between people (data creators, project personnel, and data users), products (data publications, the Coastal Carbon Library, and Coastal Carbon Atlas), and processes, outlined in this manuscript. Data creators can publish their data independently or through Coastal Carbon Network. Public data enters the Coastal Carbon Library, a centralized data synthesis, after multiple stages of data reshaping and quality control. Data users can access the library directly or through the Coastal Carbon Atlas.



(2.8%), or swamp ecosystems (1.8%). Algal mat, scrub/shrub, adjacent upland, and tidal flat ecosystems made up <1% of the database. Also, 93.6% of cores in the database are either classified as 'natural' or do not specify any anthropogenic impacts.

The most common types of soils data for marsh, mangrove, seagrass, and swamp ecosystems included dry bulk density, fraction organic matter, and fraction carbon measurements (Figure 4). Soil cores with a suite of basic soil carbon stock information included 4815 profiles (Figure 5). However, 90.4% were not confirmed to reach a contact point between wetland sediment and bedrock or

non-wetland sediment. Only 19.5% of cores were greater than or equal to 1 m long, the target depth of soil maps used in some organic carbon stock assessments (Holmquist et al., 2018b).

Of the marsh, mangrove, seagrass, and swamp cores with sufficient data to calculate carbon stocks (4815), a subset of those had data sufficient to calculate carbon accumulation rates (533, Figure 5). Of these, the majority (77.8%) have fully transparent age-depth information traceable back to the original measurements (Figure 5). Of these dated cores, 326 also have associated elevation data, with 317 having both high-quality elevation and age-depth information (Figure 5).




FIGURE 3 The Coastal Carbon Atlas: (a) Shows the map view with warmer colors indicating denser clusters of data. Clusters change and become more detailed as a user zooms in on a particular region or location. When zoomed all the way in users can identify individual coring locations. (b) An example of the Atlas's tabular view, which allows users to browse the raw data stored in the Coastal Carbon Library and accessible for download. (c) Shows the download screen including summary statistics of a query and options of post-processing depth series to output as depth weighted averages. (d) Shows a file tree of downloads including file summary in 'readMe', attribute and variable definitions in the 'data dictionary' file, the bibliographic information in three file formats, the site, core, depth series, methods, impacts, and species table, as well as the derived normalized weighted averages depth series table in the 'standardized depth series' table. The interactive application is available online (https://shiny.si.edu/coastal_carbon_atlas). Map lines do not necessarily depict accepted national boundaries.

TABLE 9 We compared global area estimates for three major blue carbon ecosystems, tidal marshes, mangroves, and seagrasses, compared with the representativeness of data appropriate for carbon stock assessments and burial rates when only the subset of those habitats is considered. Multiple estimates were found, so we present ranges based on the maximal scenario for mangroves and minimal for marshes, as well as maximal for marshes and minimal for mangroves. Despite variation in estimates of areas based on source, the current iteration of the Coastal Carbon Library clearly overrepresents marshes.

| Habitat | Global area | | | Coastal carbon library representation | |
|----------|-------------|----------------------------------|-----------|---------------------------------------|------------------------|
| | Source | Area estimate (km ²) | Area (%) | Carbon stocks (%) | Carbon burial rate (%) |
| Marsh | a,b | 90,800–152,361 | 17.0–32.7 | 61.3 | 92.9 |
| Mangrove | a,c | 152,604–284,803 | 32.8–53.1 | 25.2 | 4.98 |
| Seagrass | a | 160,387 | 30.0–34.5 | 7.04 | 2.11 |

^aMcKenzie et al. (2020).

^bMurray et al. (2022).

^cBunting et al. (2022).

When subsetting the major three habitats, marsh, mangrove, and seagrasses, and comparing to global estimates of area, it is clear that the current database over-represents tidal marshes and under-represents mangroves and seagrasses for carbon stock, but especially for carbon burial rate data (Table 9). Depending on the area estimate used, tidal marshes may represent between 17 and 32% of intertidal area, but tidal marshes dominate ecosystem representation

in the database, at 61.3% of the carbon stock data and 92.9% of the carbon burial data. Seagrass and mangrove area are at least equal to, and very likely greater than, tidal marsh area (Table 9).

Data in the Coastal Carbon Library originated from 64 coastal countries, but the majority (59.9%) were from the United States despite the fact that the United States represented 8.6% of global tidal ecosystem extent (Murray et al., 2022; Figure 6). Countries such as

Indonesia, Canada, Brazil, and Papua New Guinea were undersampled despite having relatively large areas (Figure 6).

Median organic matter density was highest in swamps and marshes, lower in mangroves, and lowest in seagrasses (Table 10).




FIGURE 4 Summary of the count of cores with key measured attributes associated with them. Dry bulk density, fraction organic matter, and fraction carbon were most commonly measured. Data associated with core elevation and stratigraphic dating were less common.

Dry bulk density, organic matter fraction, and their products, organic matter density, were non-normally distributed (Figure 7). All varied significantly based on habitat type according to a non-parametric Kruskal-Wallis rank-sum test ($p < 2.2e-16$).

5 | DISCUSSION

5.1 | Strengths of current effort

The Coastal Carbon Library and Atlas's strengths are in providing an example of how to build F.A.I.R. data in the earth and environmental sciences (Wilkinson et al., 2016). Providing support in issuing persistent digital object identifiers helps make data more findable. Accessibility was increased by preparing new data releases and distributing the Coastal Carbon Library all under creative commons open source licenses. Accessibility was further improved by creating the Coastal Carbon Atlas so that data could be queried by those without specialized data management and coding skills. The data were made interoperable by the creation and adoption of a controlled data structure and vocabulary. The data were made reusable by providing them in a highly disaggregated form. For example, for profiles available for calculating accretion rates and carbon burial rates, the majority of these data are fully reported, meaning that these metrics can be derived using existing publicly available age-depth modeling software (Aquino-López et al., 2018; Blaauw & Christen, 2011).

Beyond enacting F.A.I.R. data principles, the strengths of the Coastal Carbon Library and Atlas are in enhancing transparency.




FIGURE 5 Summary of data quality and completeness for three different purposes: carbon stock assessments, estimating carbon burial rates, and parameterizing models of future carbon sequestration. We define sediment profiles as complete if reaching the contact point between wetland sediment and bedrock or lower non-wetland sediment.




FIGURE 6 (a) Points show country by country, the makeup of the Coastal Carbon Library as percentages, with respect to the global area of tidal ecosystems (Murray et al., 2022). Countries in the top 6 of database representation and/or tidal area are labeled. The black line represents an idealized one-to-one relationship with countries above the line being over-sampled and those below, under-sampled. Note the log-10 scale. Countries with no representation in the Coastal Carbon Library are plotted at the bottom of the y-axis. The United States is over-represented, while Indonesia, Brazil, Papua New Guinea, and Canada were underrepresented. (b) Maps show same information as (a), with colors visualizing the degree of over- or under-representativeness. Here we define over- or under-representation as actual minus ideal representation. Ideal representation is based on based on tidal habitat coverage (x-axis of a), actual on database representation (y-axis of a). Map lines do not necessarily depict accepted national boundaries.

The ability to independently replicate derivative calculations and link measurements to original studies are both important to the validity of greenhouse gas inventories (Crooks et al., 2018) and carbon market verification (Needelman et al., 2018).

When collating data for use by a broad swathe of researchers and managers, the importance of grassroots approaches cannot be understated. The fact that this was a domain-specific effort allowed us to grow the database by leveraging professional networks, generate enthusiasm with a community that understood the immediate utility of the synthesis, and build trust based on relationships. Building trust took an understanding of both academic incentives around sharing data and the personal nature of many datasets. In shifting the burden of curation from data producers

to dedicated staff this effort has helped data creators comply with open data standards (Tedesco et al., 2021), and rescued data that otherwise would have never been made public (Todd-Brown et al., 2022). Our ambition is to continue providing some data curatorial services depending on the availability of funding through new projects.


5.2 | Future improvements

While there are myriad strengths to the current effort, there are also limitations to estimating deep carbon stocks and assessing carbon burial rates outside of marsh ecosystems. Most of the dated or deep

TABLE 10 Summary statistics of organic matter fraction, dry bulk density (g cm^{-3}), and organic matter density (g cm^{-3}). Summary statistics include mean, median, lower and upper 95% quantile, standard deviation (SD) and data point count (n).

| Variable | Habitat | Lower 95% | Median | Mean | Upper 95% | SD | n |
|-------------------------|----------|-----------|--------|------|-----------|------|--------|
| Fraction organic matter | Marsh | 0.02 | 0.15 | 0.2 | 0.66 | 0.17 | 12,746 |
| | Mangrove | 0 | 0.06 | 0.18 | 0.67 | 0.22 | 1433 |
| | Swamp | 0.03 | 0.27 | 0.36 | 0.91 | 0.27 | 594 |
| | Seagrass | 0 | 0.02 | 0.02 | 0.12 | 0.04 | 2805 |
| Dry bulk density | Marsh | 0.09 | 0.48 | 0.56 | 1.47 | 0.37 | 12,746 |
| | Mangrove | 0.13 | 0.75 | 0.75 | 1.61 | 0.46 | 1433 |
| | Swamp | 0.06 | 0.31 | 0.45 | 1.4 | 0.39 | 594 |
| | Seagrass | 0.31 | 0.99 | 1.06 | 1.93 | 0.45 | 2805 |
| Organic matter density | Marsh | 0.01 | 0.06 | 0.07 | 0.19 | 0.04 | 12,746 |
| | Mangrove | 0 | 0.05 | 0.06 | 0.13 | 0.04 | 1433 |
| | Swamp | 0.03 | 0.08 | 0.08 | 0.19 | 0.04 | 594 |
| | Seagrass | 0 | 0.02 | 0.02 | 0.07 | 0.02 | 2805 |

FIGURE 7 Summary of global blue carbon data shown as the frequency distribution of dry bulk density (g cm^{-3}), fraction organic matter, and organic matter density (g cm^{-3}). Note, the y-axes are different for each type of ecosystem in the rows.



cores in the Coastal Carbon Library were from tidal marshes. For carbon stocks, we showed that the majority of datasets did not represent full profiles, reaching a contact point between wetland and deeper non-wetland layer, and in most cases, soils datasets were limited by the depth of the coring device. So, while the characterization of shallow carbon stock is the most widely available application of the data, future work could quantify the effect that profile depth has on total carbon stock assessments, how spatially predictable that

contact point is from existing data, and how predictable deeper carbon stocks are from shallower ones.

Future efforts could also focus on improving the representativeness of data in the repository. The majority of the data were from the U.S. and future efforts could improve the representation of tropical and developing countries (Wylie et al., 2016). Poor representation in some cases is due to lack of data, and other times due to the fact that data has not yet been integrated. Efforts like this allow

the community to take stock of what data are already available and focus on new data collection and synthesis where it would be most impactful.

Further, future efforts could include increasing the representation of mangroves, and seagrasses, especially for carbon burial data. Finally, future efforts could analyze whether the Coastal Carbon Library is biased towards pristine wetlands and underrepresents degraded or restored wetlands.

5.3 | Proposed best practices for blue carbon data dissemination and reuse

This data synthesis effort has resulted in proposed guidelines for both individual research groups as well as journal editors, reviewers, and funders. We suggest that best practices for reporting blue carbon soils data should include distributing disaggregated data associated with the summary statistics presented in journal articles and reports. We propose reporting at the level of original measurements, for example, loss-on-ignition and dry bulk density measurements reported for individual depth increments, and individual radioisotope measurements determining supported and unsupported ^{210}Pb activity profiles in addition to derived age-depth models. We propose that positional information, elevation, wetland management history, salinity, and vegetation composition are all vital to data reinterpretation; we provide multiple ways for coding data resolution and methodology (Tables 1 and 2) in order to represent the original studies with fidelity. We propose that data producers should provide separate files for original measurements and derivative calculations, ideally with an open-source scripting workflow documenting how derivations are made. We propose utilizing dataset templates and consistent vocabulary whenever possible, such as those provided as Data S1 (Holmquist et al., 2023). The templates we provide can be modified and added to. We propose metadata accompanying data releases should define attributes, specify units, and detail study methods.

We also propose best practices for data producers include making data freely available in public data repositories with open-source Creative Commons licenses. We encourage journal editors and funders to require data publication as part of publication and end-of-project reporting, and for reviewers to ensure data releases are analysis-ready and well-documented. Funders have a role in supporting both project-specific data curation, as well as community-wide aggregation efforts. This effort shows the value of dedicated staff in helping data producers meet their open data ambitions.

Having a synthetic dataset can allow researchers to develop additional best practices, and disentangle the effect different methodological choices have on carbon burial and accretion estimates (Holmquist et al., 2021). Fusing data with soil formation models (Morris et al., 2002; Schile et al., 2014) could allow for more standardization surrounding concepts such as accretion, burial, accumulation, and sequestration.

The adoption of open data policies has the potential to improve equity for researchers and communities in the global south (Serwadda et al., 2018). Although adopting open data is not free

from ethical risks on its own, the protocol we propose does provide a way to avoid ‘parachute science’, in which data are collected from a middle-income or developing country, but not made available to the communities from which it came. Data and journal publications offer an avenue for credit towards researchers from the global south by recognizing a taxonomy of roles, including for local people who consulted and physically collected the data (Allen et al., 2014; Serwadda et al., 2018).

We finally propose some best practices for those reusing data for new purposes. Although the CC-BY 4.0 open data allows unrestricted use with attribution, we encourage data users to interact with the Coastal Carbon Library and Atlas products not only as a database but also as the community that built it. This could also include reaching out to original data producers when local knowledge is warranted for data interpretation. Data users can offer co-authorship when consultations result in more substantial contributions (Allen et al., 2014), or recommend data providers as reviewers, notify them of open-peer review periods, notify them of preprints, and provide copies of journal articles when a dataset they provided is vital to a new study.

Our hope is that by providing a centralized open database, building the practice of data publication, and implementing a data reuse policy that centers original data producers, we will contribute towards democratizing the development of coastal ecosystem service science.

6 | CONCLUSIONS

We present the Coastal Carbon Library, an open-source database for disaggregated global tidal wetland soil carbon stock, and accumulation rate data. It is made up of 6723 soil profiles, from 64 countries. In addition to the data itself, the vocabulary, structures, and metadata are all discussed in depth. The strength of these data products lies in their commitment to F.A.I.R. data principles and their transparency. The addition of the Coastal Carbon Atlas, which allows for data visualization, subsetting, and limited post-processing, increases the accessibility of the data for non-specialists. Future work is needed especially to increase the amount of deep carbon stock data across wetlands and calculate carbon burial rates consistently. While the database is global, new efforts are needed to increase the representation of countries outside the U.S. To conclude, we think that any scientific synthesis effort can learn important lessons from the grassroots nature of the Coastal Carbon Library. Data producers were incentivized to be involved by providing data templates, shifting the burden of curation from data producers to dedicated staff, and generating trust through a data use policy that rewards data producers through citation of primary material.

AUTHOR CONTRIBUTIONS

James R. Holmquist: Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology; project administration; software; supervision; visualization;

writing – original draft; writing – review and editing. **David Klings:** Conceptualization; data curation; methodology; software; visualization; writing – review and editing. **Michael Lonneman:** Data curation; methodology; software; visualization; writing – review and editing. **Jaxine Wolfe:** Conceptualization; data curation; methodology; software; visualization; writing – original draft; writing – review and editing. **Brandon Boyd:** Conceptualization; data curation; methodology; writing – review and editing. **Lauren N. Brown:** Conceptualization; methodology; writing – review and editing. **Meagan Eagle:** Conceptualization; data curation; methodology; writing – review and editing. **Jonathan Sanderman:** Conceptualization; data curation; methodology; writing – review and editing. **Kathe Todd-Brown:** Conceptualization; methodology; software; writing – review and editing. **E. Fay Belshe:** Conceptualization; data curation; writing – review and editing. **Samantha Chapman:** Conceptualization; data curation; writing – review and editing. **Ron Corstanje:** Conceptualization; writing – review and editing. **Christopher Janousek:** Conceptualization; data curation; methodology; writing – review and editing. **James T. Morris:** Conceptualization; writing – review and editing. **Gregory Noe:** Conceptualization; data curation; writing – review and editing. **André Rovai:** Conceptualization; writing – review and editing. **Amanda Spivak:** Conceptualization; writing – review and editing. **Megan Vahsen:** Conceptualization; writing – review and editing. **Lisamarie Windham-Myers:** Conceptualization; funding acquisition; supervision; writing – review and editing. **Kevin Kroeger:** Conceptualization; methodology; supervision; writing – review and editing. **J. Patrick Megonigal:** Conceptualization; funding acquisition; methodology; project administration; supervision; writing – review and editing.

ACKNOWLEDGEMENTS

We acknowledge Rose Cheney for providing organizational support in revising this manuscript. Initial work was funded by NSF (DEB-1655622). James Holmquist and Jaxine Wolfe were also supported by awards from Pew Charitable Trusts, the Department of Energy's COMPASS-FME program, and NOAA award (NA22OAR4310148), while drafting this manuscript. David H. Klings was supported by the National Science Foundation Graduate Research Fellowship (DGE-1842473). This work was an outgrowth of a 2017 Science of Blue Carbon meeting in Menlo Park California, sponsored by the U.S. Global Change Research Program and Commission for Environmental Cooperation. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

CONFLICT OF INTEREST STATEMENT

The authors claim no conflicts of interest.

DATA AVAILABILITY STATEMENT

A more detailed version of the Coastal Carbon Library's data structures, as well as data entry templates, are openly available on Smithsonian Libraries Figshare at <https://doi.org/10.25573/serc.21565671.v1>.

23580474.v1, reference number (Holmquist et al. 2023). The Coastal Carbon Library version used in this analysis is also openly available on Smithsonian Libraries' Figshare at <https://doi.org/10.25573/serc.21565671.v1>, reference number (Coastal Carbon Network, 2023) 2023). Ingestion scripts and workflow for generating the Coastal Carbon Library are publicly available on GitHub (<https://github.com/Smithsonian/CCN-Data-Library>). Code used for analyses and figure generation is also available on GitHub (https://github.com/Smithsonian/CCN_database_paper).

ORCID

- James R. Holmquist  <https://orcid.org/0000-0003-2546-6766>
Kathe Todd-Brown  <https://orcid.org/0000-0002-3109-8130>
André Rovai  <https://orcid.org/0000-0003-4117-2055>
Amanda Spivak  <https://orcid.org/0000-0001-6743-0783>
J. Patrick Megonigal  <https://orcid.org/0000-0002-2018-7883>

REFERENCES

- Abbott, K. M., Elsey-Quirk, T., & DeLaune, R. D. (2019). Factors influencing blue carbon accumulation across a 32-year chronosequence of created coastal marshes. *Ecosphere*, 10, e02828. <https://doi.org/10.1002/ecs2.2828>
Abbott, K. M., Quirk, T., & Delaune, R. D. (2019). Dataset: Factors influencing blue carbon accumulation across a 32-year chronosequence of created coastal marshes. https://smithsonian.figshare.com/articles/Dataset_Factors_influencing_blue_carbon_accumulation_across_a_32_year_chronosequence_of_created_coastal_marshes/10005215.
Abed-Navandi, D., & Dworschak, P. (2005). Food sources of tropical thalassinidean shrimps: A stable-isotope study. *Marine Ecology Progress Series*, 291, 159–168. <https://doi.org/10.3354/meps291159>
Abino, A. C., Castillo, J. A., & Lee, Y. J. (2014). Species diversity, biomass, and carbon stock assessments of a natural mangrove forest in Palawan, Philippines. *Pakistan Journal of Botany*, 46, 1955–1962.
Adame, M. F., Kauffman, J. B., Medina, I., Gamboa, J. N., Torres, O., Caamal, J. P., Reza, M., & Herrera-Silveira, J. A. (2013). Carbon stocks of tropical coastal wetlands within the karstic landscape of the Mexican Caribbean. *PLoS One*, 8, e56569. <https://doi.org/10.1371/journal.pone.0056569>
Adame, M. F., Santini, N. S., Tovilla, C., Vázquez-Lule, A., Castro, L., & Guevara, M. (2015). Carbon stocks and soil sequestration rates of tropical riverine wetlands. *Biogeosciences*, 12, 3805–3818. <https://doi.org/10.5194/bg-12-3805-2015>
Agawin, N., Duarte, C., & Fortes, M. (1996). Nutrient limitation of Philippine seagrasses (Cape Bolinao, NW Philippines): In situ experimental evidence. *Marine Ecology Progress Series*, 138, 233–243. <https://doi.org/10.3354/meps138233>
Ajonina, G. N., Ago, E. E., Amoussou, G., Mibog, E. D., Akambi, I. D., & Dossa, E. (2014). Carbon budget as a tool for assessing mangrove forests degradation in the western, coastal wetlands complex (Ramsar site 1017) of Southern Benin, West Africa. In S. Diop, J.-P. Barusseau, & C. Descamps (Eds.), *Estuaries of the world* (pp. 139–149). Springer International Publishing. https://doi.org/10.1007/978-3-319-06388-1_12
Allen, D. E., Dalal, R. C., Rennenberg, H., Meyer, R. L., Reeves, S., & Schmidt, S. (2007). Spatial and temporal variation of nitrous oxide and methane flux between subtropical mangrove sediments and the atmosphere. *Soil Biology and Biochemistry*, 39, 622–631. <https://doi.org/10.1016/j.soilbio.2006.09.013>
Allen, J. R., Cornwell, J. C., & Baldwin, A. H. (2021). Contributions of organic and mineral matter to vertical accretion in tidal wetlands

- across a Chesapeake Bay Subestuary. *Journal of Marine Science and Engineering*, 9, 751. <https://www.mdpi.com/2077-1312/9/7/751>
- Allen, J. R., Cornwell, J. C., & Baldwin, A. H. (2022). Dataset: Contributions of organic and mineral matter to vertical accretion in tidal wetlands across a Chesapeake Bay subestuary. https://smithsonian.figshare.com/articles/dataset/Dataset_Contributions_of_Organic_and_Mineral_Matter_to_Vertical_Accretion_in_Tidal_Wetlands_across_a_Chesapeake_Bay_Subestuary/18130892
- Allen, L., Scott, J., Brand, A., Hlava, M., & Altman, M. (2014). Publishing: Credit where credit is due. *Nature*, 508, 312–313.
- Alongi, D., Pfitzner, J., Trott, L., Tirendi, F., Dixon, P., & Klumpp, D. (2005). Rapid sediment accumulation and microbial mineralization in forests of the mangrove *Kandelia candel* in the Jiulongjiang Estuary, China. *Estuarine, Coastal and Shelf Science*, 63, 605–618. <https://doi.org/10.1016/j.ecss.2005.01.004>
- Alongi, D., Sasekumar, A., Chong, V., Pfitzner, J., Trott, L., Tirendi, F., Dixon, P., & Brunskill, G. (2004). Sediment accumulation and organic material flux in a managed mangrove ecosystem: Estimates of land-ocean-atmosphere exchange in Peninsular Malaysia. *Marine Geology*, 208, 383–402. <https://doi.org/10.1016/j.margeo.2004.04.016>
- Alongi, D., Tirendi, F., & Clough, B. (2000). Below-ground decomposition of organic matter in forests of the mangroves *Rhizophora stylosa* and *Avicennia marina* along the arid coast of Western Australia. *Aquatic Botany*, 68, 97–122. [https://doi.org/10.1016/s0304-3770\(00\)00110-8](https://doi.org/10.1016/s0304-3770(00)00110-8)
- Alongi, D., Tirendi, F., Dixon, P., Trott, L., & Brunskill, G. (1999). Mineralization of organic matter in intertidal sediments of a tropical semi-enclosed delta. *Estuarine, Coastal and Shelf Science*, 48, 451–467. <https://doi.org/10.1006/ecss.1998.0465>
- Alongi, D., Tirendi, F., Trott, L., & Xuan, T. (2000). Benthic decomposition rates and pathways in plantations of the mangrove *Rhizophora apiculata* in the Mekong Delta, Vietnam. *Marine Ecology Progress Series*, 194, 87–101. <https://doi.org/10.3354/meps194087>
- Alongi, D., Trott, L., Undu, M., & Tirendi, F. (2008). Benthic microbial metabolism in seagrass meadows along a carbonate gradient in Sulawesi, Indonesia. *Aquatic Microbial Ecology*, 51, 141–152. <https://doi.org/10.3354/ame01191>
- Alongi, D., Wattayakorn, G., Pfitzner, J., Tirendi, F., Zagorskis, I., Brunskill, G., Davidson, A., & Clough, B. (2001). Organic carbon accumulation and metabolic pathways in sediments of mangrove forests in southern Thailand. *Marine Geology*, 179, 85–103. [https://doi.org/10.1016/s0025-3227\(01\)00195-5](https://doi.org/10.1016/s0025-3227(01)00195-5)
- Alongi, D. M. (1996). The dynamics of benthic nutrient pools and fluxes in tropical mangrove forests. *Journal of Marine Research*, 54, 123–148. <https://doi.org/10.1357/0022240963213475>
- Alongi, D. M., Trott, L. A., Rachmansyah, F. T., McKinnon, A. D., & Undu, M. C. (2008). Growth and development of mangrove forests overlying smothered coral reefs, Sulawesi and Sumatra, Indonesia. *Marine Ecology Progress Series*, 375, 97–109. <https://doi.org/10.3354/meps07661>
- Al-rousan, S. A., Rasheed, M. Y., Khalaf, M. A., & Badran, M. I. (2005). Ecological and geochemical characteristics of bottom habitats at the northern Jordanian coast of the Gulf of Aqaba. *Chemistry and Ecology*, 21, 227–239. <https://doi.org/10.1080/02757540500211277>
- Amon, R. M. W., & Herndl, G. J. (1991). Deposit feeding and sediment. *Marine Ecology*, 12, 163–174. <https://doi.org/10.1111/j.1439-0485.1991.tb00250.x>
- Andreatta, A., Fusi, M., Cameldi, I., Cimò, F., Carnicelli, S., & Cannicci, S. (2014). Mangrove carbon sink. Do burrowing crabs contribute to sediment carbon storage? Evidence from a Kenyan mangrove system. *Journal of Sea Research*, 85, 524–533. <https://doi.org/10.1016/j.seares.2013.08.010>
- Andreatta, A., Huertas, A. D., Lotti, M., & Cerise, S. (2016). Land use changes affecting soil organic carbon storage along a mangrove swamp rice chronosequence in the Cacheu and Oio regions (northern Guinea-Bissau). *Agriculture, Ecosystems & Environment*, 216, 314–321. <https://doi.org/10.1016/j.agee.2015.10.017>
- Appleby, P. G., & Oldfield, F. (1978). The calculation of lead-210 dates assuming a constant rate of supply of unsupported ^{210}pb to the sediment. *Catena*, 5, 1–8.
- Aquino-López, M. A., Blaauw, M., Christen, J. A., & Sanderson, N. K. (2018). Bayesian analysis of ^{210}pb dating. *Journal of Agricultural, Biological and Environmental Statistics*, 23, 317–333.
- Arias-Ortiz, A., Masqué, P., García-Orellana, J., Serrano, O., Mazarrasa, I., Marbà, N., Lovelock, C. E., Lavery, P. S., & Duarte, C. M. (2018). Reviews and syntheses: ^{210}pb -derived sediment and carbon accumulation rates in vegetated coastal ecosystems—setting the record straight. *Biogeosciences*, 15, 6791–6818.
- Arias-Ortiz, A., Masqué, P., Paytan, A., & Baldocch, D. D. (2021). Dataset: Tidal and nontidal marsh restoration: A trade-off between carbon sequestration, methane emissions, and soil accretion. https://smithsonian.figshare.com/articles/dataset/Dataset_Tidal_and_nontidal_marsh_restoration_a_trade-off_between_carbon_sequestration_methane_emissions_and_soil_accretion/15127743
- Arias-Ortiz, A., Oikawa, P. Y., Carlin, J., Masqué, P., Shaham, J., Kanneg, S., Paytan, A., & Baldocch, D. D. (2021). Tidal and nontidal marsh restoration: A trade-off between carbon sequestration, methane emissions, and soil accretion. *Journal of Geophysical Research: Biogeosciences*, 126, e2021JG006573. <https://doi.org/10.1029/2021JG006573>
- Arriola, J. M., & Cable, J. E. (2017). Variations in carbon burial and sediment accretion along a tidal creek in a Florida salt marsh. *Limnology and Oceanography*, 62, S15–S28. <https://doi.org/10.1002/limo.10652>
- Ashton, E. C., & Macintosh, D. J. (2002). Preliminary assessment of the plant diversity and community ecology of the Sematan mangrove forest, Sarawak, Malaysia. *Forest Ecology and Management*, 166, 111–129.
- Banerjee, K., Chowdhury, M. R., Sengupta, K., Sett, S., & Mitra, A. (2012). Influence of anthropogenic and natural factors on the mangrove soil of Indian Sundarbans Wetland. *Archives of Environmental Science*, 6, 80–91.
- Barrón, C., Marbá, N., Terrados, J., Kennedy, H., & Duarte, C. M. (2004). Community metabolism and carbon budget along a gradient of seagrass (*Cymodocea nodosa*) colonization. *Limnology and Oceanography*, 49, 1642–1651. <https://doi.org/10.4319/lo.2004.49.5.1642>
- Baustian, M. M., Stagg, C. L., Perry, C. L., Moss, L. C., & Carruthers, T. J. B. (2021). Long-term carbon sinks in marsh soils of coastal Louisiana are at risk to wetland loss. *Journal of Geophysical Research: Biogeosciences*, 126, e2020JG005832. <https://doi.org/10.1029/2020JG005832>
- Baustian, M. M., Stagg, C. L., Perry, C. L., Moss, L. C., Carruthers, T. J. B., Allison, M. A., & Hall, C. T. (2021). Long-term soil carbon data and accretion from four marsh types in Mississippi River delta in 2015. <https://www.sciencebase.gov/catalog/item/5b3299d4e4b040769c159bb0>
- Belshe, E., Sanjuan, J., Leiva-Dueñas, C., Piñeiro-Juncal, N., Serrano, O., Lavery, P., & Mateo, M. A. (2019a). Modeling organic carbon accumulation rates and residence times in coastal vegetated ecosystems. *Journal of Geophysical Research: Biogeosciences*, 124, 3652–3671. <https://doi.org/10.1029/2019jg005233>
- Belshe, E., Sanjuan, J., Leiva-Dueñas, C., Piñeiro-Juncal, N., Serrano, O., Lavery, P. S., & Mateo, M. A. (2019b). Dataset: Modeling organic carbon accumulation rates and residence times in coastal vegetated ecosystems. https://smithsonian.figshare.com/articles/Datas et_Modeling_organic_carbon_accumulation_rates_and_residence _times_in_coastal_vegetated_ecosystems/9856769
- Bernhardt, C. E., Krauss, K. W., Cormier, N., From, A., Jones, M. C., Conner, W. H., Noe, G., & Duberstein, J. A. (2018). Carbon budget

- assessment of tidal freshwater forested wetland and oligohaline marsh ecosystems along the Waccamaw and Savannah Rivers, USA (2005–2016). <https://www.sciencebase.gov/catalog/item/59e9f79de4b05fe04cd690a1>
- Blaauw, M., & Christen, J. A. (2011). Flexible paleoclimate age-depth models using an autoregressive gamma process. International Society for Bayesian Analysis.
- Blake, G. (1965). Bulk density. In A. Klute (Ed.), *Agronomy monographs* (Vol. 9, pp. 374–390). American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America.
- Borg, J., Rowden, A., Attrill, M., Schembri, P., & Jones, M. (2010). Spatial variation in the composition of motile macroinvertebrate assemblages associated with two bed types of the seagrass *Posidonia oceanica*. *Marine Ecology Progress Series*, 406, 91–104. <https://doi.org/10.3354/meps08534>
- Boschker, H., Wielemaker, A., Schaub, B., & Holmer, M. (2000). Limited coupling of macrophyte production and bacterial carbon cycling in the sediments of *Zostera* spp. meadows. *Marine Ecology Progress Series*, 203, 181–189. <https://doi.org/10.3354/meps03181>
- Bouillon, S., Dahdouh-Guebas, F., Rao, A. V. V. S., Koedam, N., & Frank, D. (2003). Sources of organic carbon in mangrove sediments: Variability and possible ecological implications. *Hydrobiologia*, 495, 33–39.
- Boyd, B. (2012). Comparison of sediment accumulation and accretion in impounded and unimpounded marshes of the Delaware estuary. Master's thesis, University of Delaware. <http://udspace.udel.edu/handle/19716/12831>
- Boyd, B., Sommerfield, C. K., Quirk, T., & Unger, V. (2019). Dataset: Accretion and sediment accumulation in impounded and unimpounded marshes in the Delaware estuary and Barnegat Bay. https://smithsonian.figshare.com/articles/Dataset_Accretion_and_sediment_accumulation_in_impounded_and_unimpounded_marshes_in_the_Delaware_Estuary_and_Barnegat_Bay/9747065
- Boyd, B. M., & Sommerfield, C. K. (2016). Marsh accretion and sediment accumulation in a managed tidal wetland complex of Delaware Bay. *Ecological Engineering*, 92, 37–46. <https://doi.org/10.1016/j.ecoleng.2016.03.045>
- Boyd, B. M., Sommerfield, C. K., & Elsey-Quirk, T. (2017). Hydrogeomorphic influences on salt marsh sediment accumulation and accretion in two estuaries of the U.S. mid-Atlantic coast. *Marine Geology*, 383, 132–145. <https://doi.org/10.1016/j.margeo.2016.11.008>
- Breithaupt, J. L., Smoak, J. M., Rivera-Monroy, V. H., Castañeda-Moya, E., Moyer, R. P., Simard, M., & Sanders, C. J. (2017). Partitioning the relative contributions of organic matter and mineral sediment to accretion rates in carbonate platform mangrove soils. *Marine Geology*, 390, 170–180. <https://doi.org/10.1016/j.margeo.2017.07.002>
- Breithaupt, J. L., Smoak, J. M., Sanders, C. J., Thomas, J., & Smith, I. (2019). Temporal variability of carbon and nutrient burial, sediment accretion, and mass accumulation over the past century in a carbonate platform mangrove forest of the Florida everglades. <https://doi.org/10.25573/serc.11310926>
- Breithaupt, J. L., Smoak, J. M., Smith, T. J., III, & Sanders, C. J. (2014). Temporal variability of carbon and nutrient burial, sediment accretion, and mass accumulation over the past century in a carbonate platform mangrove forest of the Florida everglades. *Journal of Geophysical Research: Biogeosciences*, 119, 2032–2048. <https://doi.org/10.1002/2014JG002715>
- Breithaupt, L. J., Smoak, M. J., Bianchi, S. T., Vaughn, D., Sanders, J. C., Radabaugh, R. K., Osland, M. J., Feher, L. C., Lynch, J. C., Cahoon, D. R., Anderson, G. H., Whelan, K. R., Rosenheim, B. E., Moyer, R. P., & Chambers, L. G. (2020). Dataset: Increasing rates of carbon burial in Southwest Florida coastal wetlands. <https://doi.org/10.25573/data.9894266>
- Brunskill, G., Zagorskis, I., & Pfitzner, J. (2002). Carbon burial rates in sediments and a carbon mass balance for the Herbert River Region of the Great Barrier Reef continental shelf, North Queensland, Australia. *Estuarine, Coastal and Shelf Science*, 54, 677–700. <https://doi.org/10.1006/ecss.2001.0852>
- Brunskill, G. J., Zagorskis, I., Pfitzner, J., & Ellison, J. (2004). Sediment and trace element depositional history from the Ajkwa River Estuarine mangroves of Irian Jaya (West Papua), Indonesia. *Continental Shelf Research*, 24, 2535–2551. <https://doi.org/10.1016/j.csr.2004.07.024>
- Buffington, K., Janousek, C., Thorne, K., & Dugger, B. (2020). Dataset: Carbon stocks and accretion rates for wetland sediment at Miner Slough, Sacramento-San Joaquin delta, California. https://smithsonian.figshare.com/articles/Dataset_Carbon_stocks_and_accretion_rates_for_wetland_sediment_at_Miner_Slough_Sacramento-San_Joaquin_Delta_California/11968740
- Bukoski, J. J., Broadhead, J. S., Donato, D. C., Murdiyarso, D., & Gregoire, T. G. (2017). The use of mixed effects models for obtaining low-cost ecosystem carbon stock estimates in mangroves of the Asia-Pacific. *PLoS One*, 12, e0169096. <https://doi.org/10.1371/journal.pone.0169096>
- Bulmer, R. H., Schwendenmann, L., & Lundquist, C. J. (2016). Carbon and nitrogen stocks and below-ground allometry in temperate mangroves. *Frontiers in Marine Science*, 3, 150. <https://doi.org/10.3389/fmars.2016.00150>
- Bunting, P., Rosenqvist, A., Hilarides, L., Lucas, R. M., Thomas, N., Tadono, T., Worthington, T. A., Spalding, M., Murray, N. J., & Rebelo, L.-M. (2022). Global mangrove extent change 1996–2020: Global mangrove watch version 3.0. *Remote Sensing*, 14, 3657.
- Burden, A., Garbutt, A., & Evans, C. D. (2019). Effect of restoration on saltmarsh carbon accumulation in eastern England. *Biology Letters*, 15, 20180773. <https://doi.org/10.1098/rsbl.2018.0773>
- Burden, A., Garbutt, A., Hughes, S., Oakley, S., & Tempest, J. (2018). Soil biochemical measurements from salt marshes of different ages on the Essex coast, UK (2011). <https://catalogue.ceh.ac.uk/id/0b1faab4-3539-457f-9169-b0b1fdb59bc2>
- Burns, S. J., & Swart, P. K. (1992). Diagenetic processes in holocene carbonate sediments: Florida Bay mudbanks and islands. *Sedimentology*, 39, 285–304. <https://doi.org/10.1111/j.1365-3091.1992.tb01039.x>
- Buzzelli, C. P. (1998). Dynamic simulation of littoral zone habitats in lower Chesapeake Bay. I. Ecosystem characterization related to model development. *Estuaries*, 21, 659. <https://doi.org/10.2307/1353271>
- Cahoon, D. R., & Lynch, C. J. (1997). Vertical accretion and shallow subsidence in a mangrove forest of southwestern Florida, USA. *Mangroves and Salt Marshes*, 1, 173–186.
- Callaway, J., DeLaune, R., & Patrick, W. (1997). Sediment accretion rates from four coastal wetlands along the Gulf of Mexico. *Journal of Coastal Research*, 13, 181–191.
- Callaway, J. C., Borgnis, E. L., Turner, R. E., & Milan, C. S. (2012). Carbon sequestration and sediment accretion in San Francisco Bay tidal wetlands. *Estuaries and Coasts*, 35, 1163–1181. <https://doi.org/10.1007/s12237-012-9508-9>
- Callaway, J. C., Borgnis, E. L., Eugene Turner, R., & Milan, C. S. (2019). Dataset: Carbon sequestration and sediment accretion in San Francisco Bay tidal wetlands. https://smithsonian.figshare.com/articles/Dataset_Carbon_sequestration_and_sediment_accretion_in_San_Francisco_Bay_tidal_wetlands/9693251
- Calleja, M. L., Marbà, N., & Duarte, C. M. (2007). The relationship between seagrass (*Posidonia oceanica*) decline and sulfide porewater concentration in carbonate sediments. *Estuarine, Coastal and Shelf Science*, 73, 583–588. <https://doi.org/10.1016/j.ecss.2007.02.016>
- Campbell, J. E., Lacey, E. A., Decker, R. A., Crooks, S., & Fourqurean, J. W. (2014). Carbon storage in seagrass beds of Abu Dhabi, United Arab Emirates. *Estuaries and Coasts*, 38, 242–251. <https://doi.org/10.1007/s12237-014-9802-9>
- Cardona, P., & Botero, L. (1998). Soil characteristics and vegetation structure in a heavily deteriorated mangrove forest in the Caribbean

- Coast of Colombia. *Biotropica*, 30, 24–34. <https://doi.org/10.1111/j.1744-7429.1998.tb00366.x>
- Carlin, J., Oikawa, P. Y., Arias-Ortiz, A., Kanneg, S., Duncan, T., & Beener, K. (2021). Dataset: Sedimentary organic carbon measurements in a restored coastal wetland in San Francisco Bay, CA, USA. https://smithsonian.figshare.com/articles/dataset/Dataset_Sedimentary_organic_carbon_measurements_in_a_restored_coastal_wetland_in_San_Francisco_Bay_CA_USA/16416684
- Carruthers, T. J. B., Barnes, P. A. G., Jacome, G. E., & Fourqurean, J. W. (2005). Lagoon scale processes in a coastally influenced caribbean system: Implications for the seagrass *Thalassia testudinum*. *Caribbean Journal of Science*, 41, 441–455. https://repository.si.edu/bitstream/handle/10088/3827/Carruthers_Barnes_Jacome_and_Fourqurean_2005.pdf?sequence=1&isAllowed=y
- Cavanaugh, K. C., Kellner, J. R., Forde, A. J., Gruner, D. S., Parker, J. D., Rodriguez, W., & Feller, I. C. (2014). Poleward expansion of mangroves is a threshold response to decreased frequency of extreme cold events. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 723–727.
- Chamberlain, S. A., & Szöcs, E. (2013). taxize: Taxonomic search and retrieval in R. *F1000Research*, 2, 191.
- Chambers, L., Steinmuller, H., Dittmer, K., White, J., Cook, R., & Xue, Z. (2019). Barataria Bay carbon mineralization and biogeochemical properties from nine soil cores. <https://doi.org/10.1575/1912/bco-dmo.775547.1>
- Chanda, A., Akhand, A., Manna, S., Dutta, S., Das, I., Hazra, S., Rao, K. H., & Dadhwala, V. K. (2013). Measuring daytime CO₂ fluxes from the inter-tidal mangrove soils of Indian Sundarbans. *Environmental Earth Sciences*, 72, 417–427. <https://doi.org/10.1007/s12665-013-2962-2>
- Chen, G., Chen, B., Yu, D., Ye, Y., Tam, N. F. Y., & Chen, S. (2016). Soil greenhouse gases emissions reduce the benefit of mangrove plant to mitigating atmospheric warming effect. *Biogeosciences Discussions*, 1–22. <https://doi.org/10.5194/bg-2015-662>
- Chen, G., Tam, N., & Ye, Y. (2010). Summer fluxes of atmospheric greenhouse gases N₂O, CH₄ and CO₂ from mangrove soil in South China. *Science of the Total Environment*, 408, 2761–2767. <https://doi.org/10.1016/j.scitotenv.2010.03.007>
- Chen, G. C., Tam, N. F., & Ye, Y. (2012). Spatial and seasonal variations of atmospheric N₂O and CO₂ fluxes from a subtropical mangrove swamp and their relationships with soil characteristics. *Soil Biology and Biochemistry*, 48, 175–181. <https://doi.org/10.1016/j.soilbio.2012.01.029>
- Chen, G. C., Ulu mudin, Y. I., Pramudji, S., Chen, S. Y., Chen, B., Ye, Y., Ou, D. Y., Ma, Z. Y., Huang, H., & Wang, J. K. (2014). Rich soil carbon and nitrogen but low atmospheric greenhouse gas fluxes from North Sulawesi mangrove swamps in Indonesia. *Science of the Total Environment*, 487, 91–96. <https://doi.org/10.1016/j.scitotenv.2014.03.140>
- Chen, R., & Twilley, R. R. (1999). A simulation model of organic matter and nutrient accumulation in mangrove wetland soils. *Biogeochemistry*, 44, 93–118. <https://doi.org/10.1007/bf00993000>
- Chmura, G. L., Anisfeld, S. C., Cahoon, D. R., & Lynch, J. C. (2003). Global carbon sequestration in tidal, saline wetland soils. *Global Biogeochemical Cycles*, 17, 1111.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., & Thornton, P. (2014). Carbon and other biogeochemical cycles. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M. Midgley (Eds.), *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change* (pp. 465–570). Cambridge University Press.
- Coastal Carbon Network. (2023). Coastal carbon library (version 1.0.0). <https://doi.org/10.25573/serc.21565671>
- Coastal Protection and Restoration Authority. (2015). Crms soil properties. <https://cirms.coastal.louisiana.gov>
- Cochran, J., Hirschberg, D., Wang, J., & Dere, C. (1998). Atmospheric deposition of metals to coastal waters (Long Island Sound, New York, USA): Evidence from saltmarsh deposits. *Estuarine, Coastal and Shelf Science*, 46, 503–522. <https://doi.org/10.1006/ecss.1997.0299>
- Cotner, J. B., Suplee, M. W., Chen, N. W., & Shermann, D. E. (2004). Nutrient, sulfur and carbon dynamics in a hypersaline lagoon. *Estuarine, Coastal and Shelf Science*, 59, 639–652. <https://doi.org/10.1016/j.ecss.2003.11.008>
- Craft, C. (2007). Freshwater input structures soil properties, vertical accretion, and nutrient accumulation of Georgia and U.S. tidal marshes. *Limnology and Oceanography*, 52, 1220–1230. <https://doi.org/10.4319/lo.2007.52.3.1220>
- Craft, C., Seneca, E., & Broome, S. (1991). Loss on ignition and kjeldahl digestion for estimating organic carbon and total nitrogen in estuarine marsh soils: Calibration with dry combustion. *Estuaries*, 14, 175–179.
- Crooks, S., Rybczyk, J., O'Connell, K., Devier, D. L., Poppe, K., & Emmett-Mattox, S. (2014). *Coastal blue carbon opportunity assessment for the Snohomish estuary: The climate benefits of estuary restoration*. Technical Report, Western Washington University. <https://doi.org/10.13140/RG.2.1.1371.6568>
- Crooks, S., Sutton-Grier, A. E., Troxler, T. G., Herold, N., Bernal, B., Schile-Beers, L., & Wirth, T. (2018). Coastal wetland management as a contribution to the US National Greenhouse Gas Inventory. *Nature Climate Change*, 8, 1109–1112.
- da Silva, J. F., Duck, R., & Catarino, J. (2009). Nutrient retention in the sediments and the submerged aquatic vegetation of the coastal lagoon of the Ria de Aveiro, Portugal. *Journal of Sea Research*, 62, 276–285. <https://doi.org/10.1016/j.seares.2009.06.007>
- Danovaro, R. (1996). Detritus-bacteria-metofauna interactions in a seagrass bed (*Posidonia oceanica*) of the NW Mediterranean. *Marine Biology*, 127, 1–13. <https://doi.org/10.1007/bf00993638>
- Danovaro, R., & Fabiano, M. (1995). Seasonal and inter-annual variation of bacteria in a seagrass bed of the Mediterranean sea: Relationship with labile organic compounds and other environmental factors. *Aquatic Microbial Ecology*, 9, 17–26. <https://doi.org/10.3354/ame009017>
- Danovaro, R., Fabiano, M., & Boyer, M. (1994). Seasonal changes of benthic bacteria in a seagrass bed (*Posidonia oceanica*) of the ligurian sea in relation to origin, composition and fate of the sediment organic matter. *Marine Biology*, 119, 489–500. <https://doi.org/10.1007/bf00354310>
- Danovaro, R., & Gambi, C. (2002). Biodiversity and trophic structure of nematode assemblages in seagrass systems: Evidence for a coupling with changes in food availability. *Marine Biology*, 141, 667–677. <https://doi.org/10.1007/s00227-002-0857-y>
- De Falco, G., Baroli, M., Murru, E., Piergallini, G., & Cancemi, G. (2006). Sediment analysis evidences two different depositional phenomena influencing seagrass distribution in the Gulf of Oristano (Sardinia, Western Mediterranean). *Journal of Coastal Research*, 225, 1043–1050. <https://doi.org/10.2112/00129.1>
- de Jongh, H., Wennø, B., & Meelis, E. (1995). Seagrass distribution and seasonal biomass changes in relation to dugong grazing in the Moluccas, East Indonesia. *Aquatic Botany*, 50, 1–19. [https://doi.org/10.1016/0304-3770\(94\)00438-r](https://doi.org/10.1016/0304-3770(94)00438-r)
- De Troch, M., Van Gansbeke, D., & Vincx, M. (2006). Resource availability and meiotauna in sediment of tropical seagrass beds: Local versus global trends. *Marine Environmental Research*, 61, 59–73. <https://doi.org/10.1016/j.marenres.2005.05.003>
- Dean, W. E. (1974). Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition; comparison with other methods. *Journal of Sedimentary Research*, 44, 242–248.

- DeLaune, R., Patrick, W., Jr., & Buresh, R. (1978). Sedimentation rates determined by ¹³⁷Cs dating in a rapidly accreting salt marsh. *Nature*, 275, 532–533.
- DeLaune, R., Smith, C., Patrick, W., Jr., & Roberts, H. (1987). Rejuvenated marsh and Bay-bottom accretion on the rapidly subsiding coastal plain of us gulf coast: A second-order effect of the emerging Atchafalaya delta. *Estuarine, Coastal and Shelf Science*, 25, 381–389.
- DelVecchia, A. G., Bruno, J. F., Benninger, L., Alperin, M., Banerjee, O., & de Dios Morales, J. (2014). Organic carbon inventories in natural and restored Ecuadorian mangrove forests. *PeerJ*, 2, e388. <https://doi.org/10.7717/peerj.388>
- Devereux, R., Yates, D. F., Aukamp, J., Quarles, R. L., Jordan, S. J., Stanley, R. S., & Eldridge, P. M. (2011). Interactions of *Thalassia testudinum* and sediment biogeochemistry in Santa Rosa Sound, NW Florida. *Marine Biology Research*, 7, 317–331. <https://doi.org/10.1080/17451000.2010.515227>
- Dissanayake, N., & Chandrasekara, U. (2014). Effects of mangrove zonation and the physicochemical parameters of soil on the distribution of macrobenthic fauna in Kadolkele mangrove forest, a tropical mangrove forest in Sri Lanka. *Advances in Ecology*, 2014, 1–13. <https://doi.org/10.1155/2014/564056>
- Donato, D., Kauffman, J., Mackenzie, R., Ainsworth, A., & Pfleeger, A. (2012). Whole-island carbon stocks in the tropical pacific: Implications for mangrove conservation and upland restoration. *Journal of Environmental Management*, 97, 89–96. <https://doi.org/10.1016/j.jenvman.2011.12.004>
- Donato, D. C., Kauffman, J. B., Murdiyarno, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, 4, 293–297. <https://doi.org/10.1038/ngeo1123>
- Doughty, C., Adam Langley, J., Walker, W., Feller, I. C., Schaub, R., & Chapman, S. (2019). *Mangroves marching northward: The impacts of rising seas and temperatures on ecosystems at Kennedy space center*. https://smithsonian.figshare.com/articles/Mangroves_marching_northward_the_impacts_of_rising_seas_and_temperatures_on_ecosystems_at_Kennedy_Space_Center/9695918/1
- Doughty, C. L., Langley, J. A., Walker, W. S., Feller, I. C., Schaub, R., & Chapman, S. K. (2015). Mangrove range expansion rapidly increases coastal wetland carbon storage. *Estuaries and Coasts*, 39, 385–396. <https://doi.org/10.1007/s12237-015-9993-8>
- Drake, K., Halifax, H., Adamowicz, S. C., & Craft, C. (2015). Carbon sequestration in tidal salt marshes of the Northeast United States. *Environmental Management*, 56, 998–1008. <https://doi.org/10.1007/s00267-015-0568-z>
- Drexler, J. Z., de Fontaine, C. S., & Brown, T. A. (2009). Peat accretion histories during the past 6,000 years in marshes of the Sacramento-San Joaquin delta, CA, USA. *Estuaries and Coasts*, 32, 871–892. <https://doi.org/10.1007/s12237-009-9202-8>
- Drexler, J. Z., Krauss, K. W., Sasser, M. C., Fuller, C. C., Swarzenski, C. M., Powell, A., Swanson, K. M., & Orlando, J. (2013). A long-term comparison of carbon sequestration rates in impounded and naturally tidal freshwater marshes along the lower Waccamaw river, South Carolina. *Wetlands*, 33, 965–974. <https://doi.org/10.1007/s13157-013-0456-3>
- Drexler, J. Z., Woo, I., Fuller, C. C., & Nakai, G. (2019). Carbon accumulation and vertical accretion in a restored versus historic salt marsh in southern Puget Sound, Washington, United States. *Restoration Ecology*, 27, 1117–1127. <https://doi.org/10.1111/rec.12941>
- Dung, L. V., Tue, N. T., Nhuan, M. T., & Omori, K. (2016). Carbon storage in a restored mangrove forest in can gio mangrove forest park, Mekong delta, Vietnam. *Forest Ecology and Management*, 380, 31–40. <https://doi.org/10.1016/j.foreco.2016.08.032>
- Dutta, M., Chowdhury, C., Jana, T., & Mukhopadhyay, S. (2013). Dynamics and exchange fluxes of methane in the estuarine mangrove environment of the Sundarbans, NE coast of India. *Atmospheric Environment*, 77, 631–639. <https://doi.org/10.1016/j.atmosenv.2013.05.050>
- Eagle, M. J., Kroeger, K. D., Spivak, A. C., Wang, F., Tang, J., Abdul-Aziz, O. I., Ishtiaq, K. S., Suttles, J. O., & Mann, A. G. (2022). Soil carbon consequences of historic hydrologic impairment and recent restoration in coastal wetlands. *Science of the Total Environment*, 848, 157682.
- Elsey-Quirk, T., Seliskar, D. M., Sommerfield, C. K., & Gallagher, J. L. (2011). Salt marsh carbon pool distribution in a mid-Atlantic Lagoon, USA: Sea level rise implications. *Wetlands*, 31, 87–99. <https://doi.org/10.1007/s13157-010-0139-2>
- Ensign, S. H., Noe, G. B., Hupp, C. R., & Skalak, K. J. (2015). Head-of-tide bottleneck of particulate material transport from watersheds to estuaries. *Geophysical Research Letters*, 42, 10671–10679. <https://doi.org/10.1002/2015gl066830>
- Ensign, S. H., Noe, G. B., Hupp, C. R., & Skalak, K. J. (2021). Dataset: Head of tide bottleneck of particulate material transport from watersheds to estuaries. https://smithsonian.figshare.com/articles/dataset/Dataset_Head_of_tide_bottleneck_of_particulate_material_transport_from_watersheds_to_estuaries/13483332
- EPA. (2016a). National wetland condition assessment 2011. <https://www.epa.gov/national-aquatic-resource-surveys/data-national-aquatic-resource-surveys>
- EPA. (2016b). National wetland condition assessment: 2011 technical report. <https://www.epa.gov/national-aquatic-resource-surveys/national-wetland-condition-assessment-2011-technical-report>
- Erfemeijer, P. L. A., & Middelburg, J. J. (1993). Sediment-nutrient interactions in tropical seagrass beds: A comparison between a terrigenous and a carbonate sedimentary environments in South Sulawesi. *Marine Ecology Progress Series*, 102. <https://www.jstor.org/stable/24841993>, 187–198.
- ESRI Data and Maps. (2015a). World administrative divisions: Version 10.3. <http://sandbox.idre.ucla.edu/mapshare/data/world/data/admin.zip>
- ESRI Data and Maps. (2015b). World countries: Version 10.3. <http://sandbox.idre.ucla.edu/mapshare/data/world/data/country.zip>
- Eyre, B., & Ferguson, A. (2002). Comparison of carbon production and decomposition, benthic nutrient fluxes and denitrification in seagrass, phytoplankton, benthic microalgae- and macroalgae-dominated warm-temperate Australian lagoons. *Marine Ecology Progress Series*, 229, 43–59. <https://doi.org/10.3354/meps229043>
- Escurra, P., Ezcurra, E., Garcillán, P. P., Costa, M. T., & Aburto-Oropeza, O. (2016). Coastal landforms and accumulation of mangrove peat increase carbon sequestration and storage. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 4404–4409. <https://doi.org/10.1073/pnas.1519774113>
- Fell, C., Adgie, T., & Chapman, S. (2021). Dataset: Carbon sequestration across northeastern, Florida Coastal Wetlands. https://smithsonian.figshare.com/articles/dataset/Dataset_Carbon_sequestration_across_Northeastern_Florida_Coastal_Wetlands/15043920
- Ferreira, T., Vidal-Torrado, P., Otero, X., & Mací, F. (2007). Are mangrove forest substrates sediments or soils? A case study in southeastern Brazil. *Catena*, 70, 79–91. <https://doi.org/10.1016/j.catena.2006.07.006>
- FGDC Wetlands Subcommittee. (2009). Wetlands mapping standard. *Federal Geographic Data Committee document number FGDC-STD-015-2009*.
- Flanders Marine Institute. (2019). *Maritime boundaries geodatabase: Maritime boundaries and exclusive economic zones (200nm), version 11*. <https://www.marineregions.org/>
- Fonseca, G., Hutchings, P., & Gallucci, F. (2011). Meiofaunal communities of seagrass beds (*Zostera capricorni*) and unvegetated sediments along the coast of New South Wales, Australia. *Estuarine, Coastal and Shelf Science*, 91, 69–77. <https://doi.org/10.1016/j.ecss.2010.10.003>

- Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbá, N., Holmer, M., Mateo, M. A., Apostolaki, E. T., Kendrick, G. A., Krause-Jensen, D., McGlathery, K. J., & Serrano, O. (2012). Seagrass ecosystems as a globally significant carbon stock. *Nature Geoscience*, 5, 505–509. <https://doi.org/10.1038/ngeo1477>
- Fourqurean, J. W., Muth, M. F., & Boyer, J. N. (2010). Epiphyte loads on seagrasses and microphytobenthos abundance are not reliable indicators of nutrient availability in oligotrophic coastal ecosystems. *Marine Pollution Bulletin*, 60, 971–983. <https://doi.org/10.1016/j.marpolbul.2010.03.003>
- Fujimoto, K., Imaya, A., Tabuchi, R., Kuramoto, S., Utsugi, H., & Murofushi, T. (1999). Belowground carbon storage of Micronesian mangrove forests. *Ecological Research*, 14, 409–413. <https://doi.org/10.1046/j.1440-1703.1999.00313.x>
- Gacia, E., Duarte, C. M., & Middelburg, J. J. (2002). Carbon and nutrient deposition in a Mediterranean seagrass (*Posidonia oceanica*) meadow. *Limnology and Oceanography*, 47, 23–32. <https://doi.org/10.4319/lo.2002.47.1.00023>
- Gerlach, M. J., Engelhart, S. E., Kemp, A. C., Moyer, R. P., Smoak, J. M., Bernhardt, C. E., & Cahill, N. (2017). Reconstructing common era relative sea-level change on the Gulf Coast of Florida. *Marine Geology*, 390, 254–269. <https://doi.org/10.1016/j.margeo.2017.07.001>
- Giblin, A., Forbrich, I., & Plum Island Ecosystems LTER. (2018). Pie Iter high marsh sediment chemistry and activity measurements, Nelson Island creek marsh, Rowley, MA. <https://portal.edirepository.org/nis/mapbrowse?packageid=knb-lter-pie.427.1>
- Gillis, L., Belshe, F., Ziegler, A., & Bouma, T. (2017). Driving forces of organic carbon spatial distribution in the tropical seascapes. *Journal of Sea Research*, 120, 35–40. <https://doi.org/10.1016/j.seares.2016.12.006>
- Gleason, S. M., & Ewel, K. C. (2002). Organic matter dynamics on the forest floor of a Micronesian mangrove forest: An investigation of species composition shifts. *Biotropica*, 34, 190–198. <https://doi.org/10.1111/j.1744-7429.2002.tb00530.x>
- Gomes, F. H., Ker, J. C., Ferreira, T. O., dos Santos Moreau, A. M. S., & Moreau, M. S. (2016). Characterization and pedogenesis of mangrove soils from Ilhéus–BA, Brazil. *Revista Ciéncia Agronómica*, 47, 599–608. <https://doi.org/10.5935/1806-6690.20160072>
- Gonneea, M. E., Kroeger, K. D., & Amp, O. (2018). Collection, analysis, and age-dating of sediment cores from salt marshes on the South Shore of Cape Cod, Massachusetts, from 2013 through 2014. <https://www.sciencebase.gov/catalog/item/5a748e35e4b00f54eb19f96c>
- Gonneea, M. E., Paytan, A., & Herrera-Silveira, J. A. (2004). Tracing organic matter sources and carbon burial in mangrove sediments over the past 160 years. *Estuarine, Coastal and Shelf Science*, 61, 211–227. <https://doi.org/10.1016/j.ecss.2004.04.015>
- Grady, J. R. (1981). Properties of sea grass and sand flat sediments from the intertidal zone of St. Andrew Bay, Florida. *Estuaries*, 4, 335. <https://doi.org/10.2307/1352158>
- Grenz, C., Denis, L., Boucher, G., Chauvaud, L., Clavier, J., Fichez, R., & Pringault, O. (2003). Spatial variability in sediment oxygen consumption under winter conditions in a lagoonal system in New Caledonia (South Pacific). *Journal of Experimental Marine Biology and Ecology*, 285–286, 33–47. [https://doi.org/10.1016/s0022-0981\(02\)00518-x](https://doi.org/10.1016/s0022-0981(02)00518-x)
- He, B., Wen, Y., Yuan, X., Liang, H., & Shirong, L. (2002). Studies on soil physical and chemical properties and enzyme activities of different mangrove communities in Yingluo Bay of Guangxi. *Scientia Silvae Sinicae*, 38, 21–26.
- Hebert, A. B., Morse, J. W., & Eldridge, P. M. (2006). Small-scale heterogeneity in the geochemistry of seagrass vegetated and non-vegetated estuarine sediments: Causes and consequences. *Aquatic Geochemistry*, 13, 19–39. <https://doi.org/10.1007/s10498-006-9007-3>
- Hemminga, A. M., Slim, J. F., Kazungu, J., Ganssen, M. G., Nieuwewnhuize, J., & Kruyt, M. N. (1994). Carbon outwelling from a mangrove forest with adjacent seagrass beds and coral reefs. *Marine Ecology Progress Series*, 106. https://pure.knaw.nl/ws/files/474186/Hemminga_ea_0650.pdf, 291–301.
- Hill, T. D., & Anisfeld, S. C. (2015). Coastal wetland response to sea level rise in Connecticut and New York. *Estuarine, Coastal and Shelf Science*, 163, 185–193. <https://doi.org/10.1016/j.ecss.2015.06.004>
- Holmer, M., Andersen, F. Å., Nielsen, S. L., & Boschke, H. T. (2001). The importance of mineralization based on sulfate reduction for nutrient regeneration in tropical seagrass sediments. *Aquatic Botany*, 71, 1–17. [https://doi.org/10.1016/s0304-3770\(01\)00170-x](https://doi.org/10.1016/s0304-3770(01)00170-x)
- Holmer, M., Carta, C., & Anderse, F. (2006). Biogeochemical implications for phosphorus cycling in sandy and muddy rhizosphere sediments of *Zostera marina* meadows (Denmark). *Marine Ecology Progress Series*, 320, 141–151. <https://doi.org/10.3354/meps320141>
- Holmer, M., Duarte, C. M., & Marbá, N. (2003). Sulfur cycling and seagrass (*Posidonia oceanica*) status in carbonate sediments. *Biogeochemistry*, 66, 223–239. <https://doi.org/10.1023/b:biog.0000005326.35071.51>
- Holmer, M., & Frederiksen, M. S. (2007). Stimulation of sulfate reduction rates in Mediterranean fish farm sediments inhabited by the seagrass *Posidonia oceanica*. *Biogeochemistry*, 85, 169–184. <https://doi.org/10.1007/s10533-007-9127-x>
- Holmer, M., Marbà, N., Lamote, M., & Duarte, C. M. (2009). Deterioration of sediment quality in seagrass meadows (*Posidonia oceanica*) invaded by macroalgae (*Caulerpa* sp.). *Estuaries and Coasts*, 32, 456–466. <https://doi.org/10.1007/s12237-009-9133-4>
- Holmquist, J., Klinges, D., Lonneman, M., Wolfe, J., Boyd, B., Eagle, M., Sanderman, J., Todd-Brown, K., Belshe, E. F., Chapman, S., Corstanje, R., Janousek, C., Morris, J. T., Noe, G. B., Rovai, A., Spivak, A. C., Vahsen, M. L., Windham-Myers, L., Kroeger, K. D., & Megonigal, J. P. (2023). Coastal Carbon Network's Data Library Data Structure and Templates (version 1.0.0). https://smithsonian.figshare.com/articles/dataset/Coastal_Carbon_Network_s_Data_Library_Data_Structure_and_Templates_version_1_0_0/_23580474
- Holmquist, J. R., Brown, L. N., & MacDonald, G. M. (2021). Localized scenarios and latitudinal patterns of vertical and lateral resilience of tidal marshes to sea-level rise in the contiguous United States. *Earth's Future*, 9, e2020EF001804.
- Holmquist, J. R., Windham-Myers, L., Bliss, N., Crooks, S., Morris, J. T., Megonigal, J. P., Troxler, T., Weller, D., Callaway, J., Drexler, J., Ferner, M. C., Gonneea, M. E., Kroeger, K. D., Schile-Beers, L., Woo, I., Buffington, K., Boyd, B. M., Breithaupt, J., Brown, L. N., ... Woodrey, M. (2018a). Accuracy and precision of tidal wetland soil carbon mapping in the conterminous United States: Public soil carbon data release. <https://repository.si.edu/handle/10088/35684>
- Holmquist, J. R., Windham-Myers, L., Bliss, N., Crooks, S., Morris, J. T., Megonigal, J. P., Troxler, T., Weller, D., Callaway, J., Drexler, J., Ferner, M. C., Gonneea, M. E., Kroeger, K. D., Schile-Beers, L., Woo, I., Buffington, K., Breithaupt, J., Boyd, B. M., Brown, L. N., ... Woodrey, M. (2018b). Accuracy and precision of tidal wetland soil carbon mapping in the conterminous United States. *Scientific Reports*, 8, 9478.
- Howard, J., Sutton-Grier, A., Herr, D., Kleypas, J., Landis, E., Mcleod, E., Pidgeon, E., & Simpson, S. (2017). Clarifying the role of coastal and marine systems in climate mitigation. *Frontiers in Ecology and the Environment*, 15, 42–50.
- Howe, A., Rodri, J., & Saco, P. (2009). Surface evolution and carbon sequestration in disturbed and undisturbed wetland soils of the Hunter Estuary, Southeast Australia. *Estuarine, Coastal and Shelf Science*, 84, 75–83. <https://doi.org/10.1016/j.ecss.2009.06.006>
- Iacono, C. L., Mateo, M. A., Gracia, E., Guasch, L., Carbonell, R., Serrano, L., Serrano, O., & Dañobeitia, J. (2008). Very high-resolution seismic-acoustic imaging of seagrass meadows (Mediterranean sea): Implications for carbon sink estimates. *Geophysical Research Letters*, 35, 5. <https://doi.org/10.1029/2008gl034773>

- Isaksen, M., & Finster, K. (1996). Sulphate reduction in the root zone of the seagrass *Zostera noltii* on the intertidal flats of a coastal lagoon (Arcachon, France). *Marine Ecology Progress Series*, 137, 187–194. <https://doi.org/10.3354/meps137187>
- Jennerjahn, T. C., & Ittekkot, V. (2002). Relevance of mangroves for the production and deposition of organic matter along tropical continental margins. *Naturwissenschaften*, 89, 23–30. <https://doi.org/10.1007/s00114-001-0283-x>
- Johnson, B. J., Moore, K. A., Lehmann, C., Bohlen, C., & Brown, T. A. (2007). Middle to late holocene fluctuations of C_3 and C_4 vegetation in a northern New England salt marsh, Sprague marsh, Phippsburg Maine. *Organic Geochemistry*, 38, 394–403. <https://doi.org/10.1016/j.orggeochem.2006.06.006>
- Jones, M. C., Bernhardt, C. E., Krauss, K. W., & Noe, G. B. (2017). The impact of late holocene land use change, climate variability, and sea level rise on carbon storage in tidal freshwater wetlands on the southeastern United States coastal plain. *Journal of Geophysical Research: Biogeosciences*, 122, 3126–3141. <https://doi.org/10.1002/2017jg004015>
- Jones, T. G., Ratsimba, H. R., Carro, A., Ravaorinorotsihoarana, L., Glass, L., Teoh, M., Benson, L., Cripps, G., Giri, C., Zafindrasiivonona, B., Raherindray, R., Andriamahenina, Z., & Andriamahefazafy, M. (2016). The mangroves of Ambanja and Ambaro Bays, Northwest Madagascar: Historical dynamics, current status and deforestation mitigation strategy. In S. Diop, P. Scheren, & J. F. Machiwa (Eds.), *Estuaries of the world* (pp. 67–85). Springer International Publishing. https://doi.org/10.1007/978-3-319-25370-1_5
- Kairis, P. A., & Rybczyk, J. M. (2010). Sea level rise and eelgrass (*Zostera marina*) production: A spatially explicit relative elevation model for Padilla Bay, WA. *Ecological Modelling*, 221, 1005–1016. <https://doi.org/10.1016/j.ecolmodel.2009.01.025>
- Kamp-Nielsen, L., Vermaat, J., Wesseling, I., Borum, J., & Geertz-Hansen, O. (2002). Sediment properties along gradients of siltation in South-East Asia. *Estuarine, Coastal and Shelf Science*, 54, 127–137. <https://doi.org/10.1006/ecss.2001.0822>
- Kao, W.-Y., Tsai, H.-C., Shih, C.-N., Tsai, T.-T., & Handley, L. L. (2002). Nutrient contents, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ during leaf senescence in the mangrove, *Kandelia candel* (L.) Druce. *Botanical Bulletin of Academia Sinica*, 43, 277–282.
- Kauffman, J., Giovannoni, L., Kelly, J., Dunstan, N., Borde, A., Diefenderfer, H., Cornu, C., Janousek, C., Apple, J., & Brophy, L. (2020a). Total ecosystem carbon stocks at the marine-terrestrial interface: Blue carbon of the pacific northwest coast, United States. *Global Change Biology*, 26, 5679–5692. <https://doi.org/10.1111/gcb.15248>
- Kauffman, J., Giovannoni, L., Kelly, J., Dunstan, N., Borde, A., Diefenderfer, H., Cornu, C., Janousek, C., Apple, J., & Brophy, L. (2020b). Dataset: Carbon stocks in seagrass meadows, emergent marshes, and forested tidal swamps of the pacific northwest. https://smithsonian.figshare.com/articles/dataset/Dataset_Carbon_stocks_in_seagrass_meadows_emergent_marshes_and_forested_tidal_swamps_of_the_Pacific_Northwest/12640172
- Kearney, M. S., & Ward, L. G. (1986). Accretion rates in brackish marshes of a Chesapeake Bay Estuarine tributary. *Geo-Marine Letters*, 6, 41–49.
- Kemp, A. C., Sommerfield, C. K., Vane, C. H., Horton, B. P., Chenery, S., Anisfeld, S., & Nikitina, D. (2012). Use of lead isotopes for developing chronologies in recent salt-marsh sediments. *Quaternary Geochronology*, 12, 40–49. <https://doi.org/10.1016/j.quageo.2012.05.004>
- Kemp, A. C., Sommerfield, C. K., Vane, C. H., Horton, B. P., Chenery, S., Anisfeld, S., & Nikitina, D. (2020). Dataset: Use of lead isotopes for developing chronologies in recent salt-marsh sediments.
- Kenig, F., Huc, A. Y., Purser, B. H., & Oudin, J.-L. (1990). Sedimentation, distribution and diagenesis of organic matter in a recent carbonate environment, Abu Dhabi, U.A.E. *Organic Geochemistry*, 16, 735–747. [https://doi.org/10.1016/0146-6380\(90\)90113-e](https://doi.org/10.1016/0146-6380(90)90113-e)
- Keshta, A. E., Yarwood, S. A., & Baldwin, A. H. (2017). Hydrology, soil redox, and pore-water iron regulate carbon cycling in natural and restored tidal freshwater wetlands in the Chesapeake Bay, Maryland, USA. Ph.D. thesis, University of Maryland. <https://drum.lib.umd.edu/handle/1903/20335>
- Keshta, A. E., Yarwood, S. A., & Baldwin, A. H. (2020). Dataset: Soil redox and hydropattern control soil carbon stocks across different habitats in tidal freshwater wetlands in a sub-estuary of the Chesapeake Bay. https://smithsonian.figshare.com/articles/dataset/Dataset_Soil_Redox_and_Hydropattern_control_Soil_Carbon_Stocks_across_different_habitats_in_Tidal_Freshwater_Wetlands_in_a_Sub-estuary_of_the_Chesapeake_Bay/13187549
- Keshta, A. E., Yarwood, S. A., & Baldwin, A. H. (2021). A new in situ method showed greater persistence of added soil organic matter in natural than restored wetlands. *Restoration Ecology*, 29, e13437. <https://doi.org/10.1111/rec.13437>
- Keuskamp, J. A., Schmitt, H., Laanbroek, H. J., Verhoeven, J. T., & Hefting, M. M. (2013). Nutrient amendment does not increase mineralisation of sequestered carbon during incubation of a nitrogen limited mangrove soil. *Soil Biology and Biochemistry*, 57, 822–829. <https://doi.org/10.1016/j.soilbio.2012.08.007>
- Koch, M., & Madden, C. (2001). Patterns of primary production and nutrient availability in a Bahamas lagoon with fringing mangroves. *Marine Ecology Progress Series*, 219, 109–119. <https://doi.org/10.3354/meps219109>
- Koepfler, E. T., Benner, R., & Montagna, P. A. (1993). Variability of dissolved organic carbon in sediments of a seagrass bed and an unvegetated area within an estuary in southern Texas. *Estuaries*, 16, 391. <https://doi.org/10.2307/1352587>
- Krause-Jensen, D., Carstensen, J., Nielsen, S., Dalsgaard, T., Christensen, P., Fossing, H., & Rasmussen, M. (2011). Sea bottom characteristics affect depth limits of eelgrass *Zostera marina*. *Marine Ecology Progress Series*, 425, 91–102. <https://doi.org/10.3354/meps09026>
- Krauss, K. W., Noe, G. B., Duberstein, J. A., Conner, W. H., Stagg, C. L., Cormier, N., Jones, M. C., Bernhardt, C. E., Lockaby, B. G., From, A. S., Doyle, T. W., Day, R. H., Ensign, S. H., Pierfelice, K. N., Hupp, C. R., Chow, A. T., & Whitbeck, J. L. (2018). The role of the upper tidal estuary in wetland blue carbon storage and flux. *Global Biogeochemical Cycles*, 32, 817–839. <https://doi.org/10.1029/2018gb005897>
- Kristensen, E., Andersen, F., Holmboe, N., Holmer, M., & Thongtham, N. (2000). Carbon and nitrogen mineralization in sediments of the Bangrong mangrove area, Phuket, Thailand. *Aquatic Microbial Ecology*, 22, 199–213. <https://www.int-res.com/abstracts/ame/v22/n2/p199-213/>
- Kristensen, E., Devol, H. A., Ahmed, S. I., & Saleem, M. (1992). Preliminary study of benthic metabolism and sulfate reduction in a mangrove swamp of the Indus delta, Pakistan. *Marine Ecology Progress Series*. <https://www.jstor.org/stable/24832497>, 90, 287–297.
- Kroeger, K. D., Crooks, S., Moseman-Valtierra, S., & Tang, J. (2017). Restoring tides to reduce methane emissions in impounded wetlands: A new and potent blue carbon climate change intervention. *Scientific Reports*, 7, 1–12.
- Kulawardhana, R. W., Feagin, R. A., Popescu, S. C., Boutton, T. W., Yeager, K. M., & Bianchi, T. S. (2015). The role of elevation, relative sea-level history and vegetation transition in determining carbon distribution in *Spartina alterniflora* dominated salt marshes. *Estuarine, Coastal and Shelf Science*, 154, 48–57. <https://doi.org/10.1016/j.ecss.2014.12.032>
- Lacerda, L., Ittekkot, V., & Patchineelam, S. (1995). Biogeochemistry of mangrove soil organic matter: A comparison between Rhizophora and Avicennia soils in South-Eastern Brazil. *Estuarine, Coastal and Shelf Science*, 40, 713–720. <https://doi.org/10.1006/ecss.1995.0048>
- Lagomasino, D., Corbett, D. R., & Walsh, J. (2013). Influence of wind-driven inundation and coastal geomorphology on sedimentation in

- two microtidal marshes, Pamlico River Estuary, NC. *Estuaries and Coasts*, 36, 1165–1180.
- Lagomasino, D., Corbett, D. R., & Walsh, J. (2020). Dataset: Influence of wind-driven inundation and coastal geomorphology on sedimentation in two microtidal marshes, pamlico river estuary, NC. https://smithsonian.figshare.com/articles/Dataset_Influence_of_Wind-Driven_Inundation_and_Coastal_Geomorphology_on_Sedimentation_in_Two_Microtidal_Marshes_Pamlico_River_Estuary_NC/1204335
- Lallier-Verges, E., Perrussel, B. P., Disnar, J.-R., & Baltzer, F. (1998). Relationships between environmental conditions and the diagenetic evolution of organic matter derived from higher plants in a modern mangrove swamp system (Guadeloupe, French West Indies). *Organic Geochemistry*, 29, 1663–1686. [https://doi.org/10.1016/s0146-6380\(98\)00179-x](https://doi.org/10.1016/s0146-6380(98)00179-x)
- Lang, J. K. S., Kairo, J. G., Mencuccini, M., Bouillon, S., Skov, M. W., Waldron, S., & Huxham, M. (2014). Rapid losses of surface elevation following tree girdling and cutting in tropical mangroves. *PLoS One*, 9, e107868. <https://doi.org/10.1371/journal.pone.0107868>
- Larkum, A. W., Orth, R. J., & Duarte, C. (2006). *Seagrasses: Biology, ecology and conservation*. Springer Science & Business Media.
- Larned, S. (2003). Effects of the invasive, nonindigenous seagrass *Zostera japonica* on nutrient fluxes between the water column and benthos in a NE Pacific Estuary. *Marine Ecology Progress Series*, 254, 69–80. <https://doi.org/10.3354/meps254069>
- Leduc, D., & Probert, P. K. (2010). Small-scale effect of intertidal seagrass (*Zostera muelleri*) on meiofaunal abundance, biomass, and nematode community structure. *Journal of the Marine Biological Association of the United Kingdom*, 91, 579–591. <https://doi.org/10.1017/s0025315410001645>
- Lee, K. H., & Baker, G. E. (1972). An ecological study of the soil microfungi in a hawaiian mangrove swamp. *Pacific Science*, 26, 1–10.
- Lee, S. Y., Oh, J. H., Choi, C. I., Suh, Y., & Mukai, H. (2005). Leaf growth and population dynamics of intertidal *Zostera japonica* on the western coast of Korea. *Aquatic Botany*, 83, 263–280. <https://doi.org/10.1016/j.aquabot.2005.07.004>
- Leopold, A., Marchand, C., Deborde, J., Chaduteau, C., & Allenbach, M. (2013). Influence of mangrove zonation on CO₂ fluxes at the sediment-air interface (New Caledonia). *Geoderma*, 202–203, 62–70. <https://doi.org/10.1016/j.geoderma.2013.03.008>
- Lewis, M. A., Dantin, D. D., Chancy, C. A., Abel, K. C., & Lewis, C. G. (2007). Florida seagrass habitat evaluation: A comparative survey for chemical quality. *Environmental Pollution*, 146, 206–218. <https://doi.org/10.1016/j.envpol.2006.04.041>
- Lillebø, A. I., Flindt, M. R., Pardal, M. A., & Marques, J. C. (2006). The effect of *Zostera noltii*, *Spartina maritima* and *Scirpus maritimus* on sediment pore-water profiles in a temperate intertidal estuary. *Hydrobiologia*, 555, 175–183. <https://doi.org/10.1007/s10750-005-1114-4>
- Liu, H., Ren, H., Hui, D., Wang, W., Liao, B., & Cao, Q. (2014). Carbon stocks and potential carbon storage in the Mangrove forests of China. *Journal of Environmental Management*, 133, 86–93. <https://doi.org/10.1016/j.jenvman.2013.11.037>
- Liu, M., Ye, Y., Cao, C., & Tang, F. (2008). Fractal characteristics and related affecting factors of particle size distribution in mangrove soils in Dongzhai Harbor of Hainan. *Chinese Journal of Ecology*, 9, 1557–1561.
- Livesley, S. J., & Andrusiak, S. M. (2012). Temperate mangrove and salt marsh sediments are a small methane and nitrous oxide source but important carbon store. *Estuarine, Coastal and Shelf Science*, 97, 19–27. <https://doi.org/10.1016/j.ecss.2011.11.002>
- Loi, L. T., & Ni, L. H. (2015). The effects of land surface elevation on belowground carbon accumulation of mangrove ecosystem at the Ong Trang Sand bar, NGOC Hien District, Ca Mau province. *Unknown*.
- Lovelock, C. E., & Duarte, C. M. (2019). Dimensions of blue carbon and emerging perspectives. *Biology Letters*, 15, 20180781.
- Lovelock, C. E., Sorrell, B. K., Hancock, N., Hua, Q., & Swales, A. (2010). Mangrove forest and soil development on a rapidly accreting shore in New Zealand. *Ecosystems*, 13, 437–451. <https://doi.org/10.1007/s10021-010-9329-2>
- Luk, S., Spivak, A., Eagle, M. J., & O'Keefe Suttles, J. A. (2020). Collection, analysis, and age-dating of sediment cores from a salt marsh platform and ponds, Rowley, Massachusetts, 2014–15. <https://www.sciencebase.gov/catalog/item/5fb41153d34eb413d5e0af37>
- Luk, S. Y., Todd-Brown, K., Eagle, M., McNichol, A. P., Sanderman, J., Gosselin, K., & Spivak, A. C. (2021). Soil organic carbon development and turnover in natural and disturbed salt marsh environments. *Geophysical Research Letters*, 48, e2020GL090287. <https://doi.org/10.1029/2020GL090287>
- MacKenzie, R. A., Foulk, P. B., Klump, J. V., Weckerly, K., Purbospito, J., Murdiyarsa, D., Donato, D. C., & Nam, V. N. (2016). Sedimentation and belowground carbon accumulation rates in mangrove forests that differ in diversity and land use: A tale of two mangroves. *Wetlands Ecology and Management*, 24, 245–261. <https://doi.org/10.1007/s11273-016-9481-3>
- Maher, D. T., & Eyre, B. D. (2010). Benthic fluxes of dissolved organic carbon in three temperate Australian Estuaries: Implications for global estimates of benthic DOC fluxes. *Journal of Geophysical Research*, 115, G04039. <https://doi.org/10.1029/2010jg001433>
- Malhotra, A., Todd-Brown, K., Nave, L. E., Batjes, N. H., Holmquist, J. R., Hoyt, A. M., Iversen, C. M., Jackson, R. B., Lajtha, K., Lawrence, C., Vindušková, O., Wieder, W., Williams, M., Hugelius, G., & Harden, J. (2019). The landscape of soil carbon data: Emerging questions, synergies and databases. *Progress in Physical Geography: Earth and Environment*, 43, 707–719.
- Mao, Z. L., Lai, H. D., Zhao, Z. Y., & Yang, X. M. (2011). Effect of invasion plants (*Mikania micrantha* H.B.K.) on carbon stock of mangrove ecosystem in Shenzhen Bay. *Ecology and Environment*, 20, 1813–1818.
- Mao, Z. L., Yang, X. M., Zhao, Z. Y., Lai, M. D., Yang, D. Y., Wu, C. L., & Xu, H. L. (2012). Preliminary study on mangrove ecosystem carbon cycle of *Kandelia candel* in Futian Nature Reserve, Shenzhen, China. *Ecology and Environmental Sciences*, 21, 1189.
- Marchand, C., Baltzer, F., Lallier-Vergès, E., & Albéric, P. (2004). Pore-water chemistry in mangrove sediments: Relationship with species composition and developmental stages (French Guiana). *Marine Geology*, 208, 361–381. <https://doi.org/10.1016/j.margeo.2004.04.015>
- Marchand, C., Lallier-Vergès, E., & Baltzer, F. (2003). The composition of sedimentary organic matter in relation to the dynamic features of a mangrove-fringed coast in French Guiana. *Estuarine, Coastal and Shelf Science*, 56, 119–130. [https://doi.org/10.1016/s0272-7714\(02\)00134-8](https://doi.org/10.1016/s0272-7714(02)00134-8)
- Marchio, D., Savarese, M., Bovard, B., & Mitsch, W. (2016). Carbon sequestration and sedimentation in mangrove swamps influenced by hydrogeomorphic conditions and urbanization in Southwest Florida. *Forests*, 7, 116. <https://doi.org/10.3390/f7060116>
- Mateo, M., & Romero, J. (1997). Detritus dynamics in the seagrass *Posidonia oceanica*: Elements for an ecosystem carbon and nutrient budget. *Marine Ecology Progress Series*, 151, 43–53. <https://doi.org/10.3354/meps151043>
- Mateo, M., Romero, J., Pérez, M., Littler, M., & Littler, D. (1997). Dynamics of millenary organic deposits resulting from the growth of the Mediterranean seagrass *Posidonia oceanica*. *Estuarine, Coastal and Shelf Science*, 44, 103–110. <https://doi.org/10.1006/ecss.1996.0116>
- Matsui and Naohiro. (1998). Estimated stocks of organic carbon in mangrove roots and sediments in Hinchinbrook Channel, Australia. *Mangroves and Salt Marshes*, 4, 199–204. <https://doi.org/10.1023/A:1009959909208>
- McClellan, S. A. (2021). Data for: Root-zone carbon and nitrogen pools across two chronosequences of coastal marshes formed using

- different restoration techniques: Dredge sediment versus river sediment diversion. <https://data.mendeley.com/datasets/5zbv2mb5zp/1>
- McClellan, S. A., Elsey-Quirk, T., Laws, E. A., & DeLaune, R. D. (2021). Root-zone carbon and nitrogen pools across two chronosequences of coastal marshes formed using different restoration techniques: Dredge sediment versus river sediment diversion. *Ecological Engineering*, 169, 106326. <https://www.sciencedirect.com/science/article/pii/S0925857421001816>
- McFadden, T. N., Kauffman, J. B., & Bhomia, R. K. (2016). Effects of nesting waterbirds on nutrient levels in mangroves, Gulf of Fonseca, Honduras. *Wetlands Ecology and Management*, 24, 217–229. <https://doi.org/10.1007/s11273-016-9480-4>
- McGlathery, K., Reynolds, L., Cole, L., Orth, R., Marion, S., & Schwarzschild, A. (2012). Recovery trajectories during state change from bare sediment to eelgrass dominance. *Marine Ecology Progress Series*, 448, 209–221. <https://doi.org/10.3354/meps09574>
- McKenzie, L. J., Nordlund, L. M., Jones, B. L., Cullen-Unsworth, L. C., Roelfsema, C., & Unsworth, R. K. (2020). The global distribution of seagrass meadows. *Environmental Research Letters*, 15, 074041.
- McLeod, E., Chmura, G. L., Bouillon, S., Salm, R., Björk, M., Duarte, C. M., Lovelock, C. E., Schlesinger, W. H., & Silliman, B. R. (2011). A blueprint for blue carbon: Toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Frontiers in Ecology and the Environment*, 9, 552–560.
- McTigue, N., Davis, J., Rodriguez, A., McKee, B., Atencio, A., & Currin, C. (2020). Dataset: Carbon accumulation rates in a salt marsh over the past two millennia.
- McTigue, N., Davis, J., Rodriguez, T., McKee, B., Atencio, A., & Currin, C. (2019). Sea-level rise explains changing carbon accumulation rates in a salt marsh over the past two millennia. *Journal of Geophysical Research: Biogeosciences*, 124, 2945–2957.
- Mellors, J., Marsh, H., Carruthers, T. J. B., & Waycott, M. (2002). Testing the sediment-trapping paradigm of seagrass: Do seagrasses influence nutrient status and sediment structure in tropical intertidal environments? *Bulletin of Marine Science*, 71, 1215–1226. <https://www.ingentaconnect.com/content/umrsmas/bullmar/2002/00000071/00000003/art00008>
- Merrill, J. Z. (1999). *Tidal freshwater marshes as nutrient sinks: Particulate nutrient burial and denitrification*. Ph.D. thesis, University of Maryland, College Park. <https://elibrary.ru/item.asp?id=5305392>
- Messerschmidt, T. C., & Kirwan, M. L. (2020). Dataset: Soil properties and accretion rates of c3 and c4 marshes at the global change research wetland, Edgewater, Maryland. https://smithsonian.figshare.com/articles/Dataset_Soil_properties_and_accretion_rates_of_C3_and_C4_marshes_at_the_Global_Change_Research_Wetland_Edgewater_Maryland/11914140
- Mfilinge, P., Atta, N., & Tsuchiya, M. (2002). Nutrient dynamics and leaf litter decomposition in a subtropical mangrove forest at Oura Bay, Okinawa, Japan. *Trees*, 16, 172–180. <https://doi.org/10.1007/s00468-001-0156-0>
- Middelburg, J. J., Nieuwenhuize, J., Slim, F. J., & Ohowa, B. (1996). Sediment biogeochemistry in an East African mangrove forest (Gazi Bay, Kenya). *Biogeochemistry*, 34, 133–155.
- Miller, C. B., Rodriguez, A. B., Bost, M. C., McKee, B. A., & McTigue, N. D. (2022). Carbon accumulation rates are highest at young and expanding salt marsh edges. *Communications Earth & Environment*, 3, 173. <https://doi.org/10.1038/s43247-022-00501-x>
- Miyajima, T., Koike, I., Yamano, H., & Iizumi, H. (1998). Accumulation and transport of seagrass-derived organic matter in reef flat sediment of Green Island, Great Barrier Reef. *Marine Ecology Progress Series*, 175, 251–259. <https://doi.org/10.3354/meps175251>
- Morris, J. T., Barber, D. C., Callaway, J. C., Chambers, R., Hagen, S. C., Hopkinson, C. S., Johnson, B. J., Megonigal, P., Neubauer, S. C., Troxler, T., & Wigand, C. (2016). Contributions of organic and inorganic matter to sediment volume and accretion in tidal wetlands at steady state. *Earth's Future*, 4, 110–121.
- Morris, J. T., Sundareshwar, P., Nietch, C. T., Kjerfve, B., & Cahoon, D. R. (2002). Responses of coastal wetlands to rising sea level. *Ecology*, 83, 2869–2877.
- Murdiyarso, D., Purbopuspito, J., Kauffman, J. B., Warren, M. W., Sasmito, S. D., Donato, D. C., Manuri, S., Krisnawati, H., Taberima, S., & Kurnianto, S. (2015). The potential of Indonesian mangrove forests for global climate change mitigation. *Nature Climate Change*, 5, 1089–1092. <https://doi.org/10.1038/nclimate2734>
- Murray, N. J., Worthington, T. A., Bunting, P., Duce, S., Hagger, V., Lovelock, C. E., Lucas, R., Saunders, M. I., Sheaves, M., Spalding, M., Waltham, N. J., & Lyons, M. B. (2022). High-resolution mapping of losses and gains of earth's tidal wetlands. *Science*, 376, 744–749.
- Nahlik, A. M., & Fennelly, M. S. (2016). Carbon storage in US wetlands. *Nature Communications*, 7, 13835. <https://doi.org/10.1038/ncomm13835>
- Naidoo, G. (1980). Mangrove soils of the Beachwood area, Durban. *Journal of South African Botany*, 46, 293–304.
- Naidoo, G., & Raiman, F. (1982). Some physical and chemical properties of mangrove soils at Sipingo and Mgeni, Natal. *South African Journal of Botany*, 1, 85–90.
- Nam, V. N., Sasmito, S. D., Murdiyarso, D., Purbopuspito, J., & MacKenzie, R. A. (2016). Carbon stocks in artificially and naturally regenerated mangrove ecosystems in the Mekong delta. *Wetlands Ecology and Management*, 24, 231–244. <https://doi.org/10.1007/s11273-015-9479-2>
- Needelman, B. A., Emmer, I. M., Emmett-Mattox, S., Crooks, S., Megonigal, J. P., Myers, D., Oreska, M. P., & McGlathery, K. (2018). The science and policy of the verified carbon standard methodology for tidal wetland and seagrass restoration. *Estuaries and Coasts*, 41, 2159–2171.
- Neubauer, S., Anderson, I., Constantine, J., & Kuehl, S. (2002). Sediment deposition and accretion in a mid-Atlantic (USA) tidal freshwater marsh. *Estuarine, Coastal and Shelf Science*, 54, 713–727. <https://doi.org/10.1006/ecss.2001.0854>
- Noe, G. B., Hupp, C. R., Bernhardt, C. E., & Krauss, K. W. (2016). Contemporary deposition and long-term accumulation of sediment and nutrients by tidal freshwater forested wetlands impacted by sea level rise. *Estuaries and Coasts*, 39, 1006–1019. <https://doi.org/10.1007/s12237-016-0066-4>
- Noe, G. B., Krauss, K. W., Lockaby, B. G., Conner, W. H., & Hupp, C. R. (2012). The effect of increasing salinity and forest mortality on soil nitrogen and phosphorus mineralization in tidal freshwater forested wetlands. *Biogeochemistry*, 114, 225–244. <https://doi.org/10.1007/s10533-012-9805-1>
- Nolte, S. (2020). Dataset: Does livestock grazing affect sediment deposition and accretion rates in salt marshes? https://smithsonian.figshare.com/articles/Dataset_Does_livestock_grazing_affect_sediment_deposition_and_accretion_rates_in_salt_marshes/_11958996
- Nolte, S., Müller, F., Schuerch, M., Wanner, A., Esselink, P., Bakker, J. P., & Jensen, K. (2013). Does livestock grazing affect sediment deposition and accretion rates in salt marshes? *Estuarine, Coastal and Shelf Science*, 135, 296–305. <https://doi.org/10.1016/j.ecss.2013.10.026>
- Nsomblo, E. N., Bengono, F. A., Etame, J., Ndongo, D., Ajonina, G., & Bilong, P. (2016). Effects of vegetation degradation on carbon stock, morphological, physical and chemical characteristics of soils within the mangrove forest of the Rio Del Rey Estuary: Case study Bamusso (South-West Cameroon). *African Journal of Environmental Science and Technology*, 10, 58–66. <https://doi.org/10.5897/ajest2015.1956>
- Nuttle, W. (1996). *Tidal freshwater marshes as nutrient sinks: Particulate nutrient burial and denitrification*. Environmental Data Initiative.
- Nyman, J., DeLaune, R., Roberts, H., & Patrick, W. (1993). Relationship between vegetation and soil formation in a rapidly submerging coastal marsh. *Marine Ecology Progress Series*, 96, 269–279. <https://doi.org/10.3354/meps09629>

- Oakes, J. M., & Connolly, R. M. (2004). Causes of sulfur isotope variability in the seagrass, *Zostera capricorni*. *Journal of Experimental Marine Biology and Ecology*, 302, 153–164. <https://doi.org/10.1016/j.jembe.2003.10.011>
- O'keefe Suttles, J. A., Eagle, M. J., Mann, A. C., & Kroeger, K. D. (2021). *Collection, analysis, and age-dating of sediment cores from mangrove and salt marsh ecosystems in Tampa Bay, Florida, 2015*. <https://www.sciencebase.gov/catalog/item/60bfb8a4d34e86b938916d6f>
- O'keefe Suttles, J. A., Eagle, M. J., Mann, A. C., Moseman-Valtierre, S., Pratt, S., & Kroeger, K. D. (2021). *Collection, analysis, and age-dating of sediment cores from salt marshes, Rhode Island, 2016*. <https://www.sciencebase.gov/catalog/item/60bfb7c2d34e86b938916d1e>
- O'keefe Suttles, J. A., Eagle, M. J., Mann, A. C., Spivak, A., Sanks, K., Daniel, R., & Kroeger, K. D. (2021). *Collection, analysis, and age-dating of sediment cores from natural and restored salt marshes on Cape Cod, Massachusetts, 2015–16*. <https://www.sciencebase.gov/catalog/item/60bfb987d34e86b938916dc9>
- O'keefe Suttles, J. A., Eagle, M. J., Mann, A. C., Wang, F., Tang, J., Daniel, R., Sanks, K., Smith, T., & Kroeger, K. D. (2021). *Collection, analysis, and age-dating of sediment cores from Herring River wetlands and other nearby wetlands in Wellfleet, Massachusetts, 2015–17*. <https://www.sciencebase.gov/catalog/item/60bfb916d34e86b938916da1>
- O'keefe Suttles, J. A., Wigand, C., Eagle, M. J., Branoff, B., Balogh, S., Miller, K., Martin, R., Hanson, A., Oczkowski, A., Huertas, E., Watson, E., & Loffredo, J. (2021). *Collection, analysis, and age-dating of sediment cores from mangrove wetlands in San Juan Bay estuary, Puerto Rico, 2016*. <https://www.sciencebase.gov/catalog/item/60902e3fd34e93746a710491>
- Ooi, J. L. S., Kendrick, G. A., Van Niel, K. P., & Affendi, Y. A. (2011). Knowledge gaps in tropical Southeast Asian seagrass systems. *Estuarine, Coastal and Shelf Science*, 92, 118–131. <https://doi.org/10.1016/j.ecss.2010.12.021>
- Orem, W. H., Holmes, C. W., Kendall, C., Lerch, H. E., Bates, A. L., Silva, S. R., Boylan, A., Corum, M., Marot, M., & Hedgeman, C. (1999). Geochemistry of Florida Bay sediments: Nutrient history at five sites in Eastern and Central Florida Bay. *Journal of Coastal Research*, 15, 1055–1071. <https://www.jstor.org/stable/4299024>
- Orson, R. A., & Simpson, R. L. (1990). Rates of sediment accumulation in a tidal freshwater marsh. *Journal of Sedimentary Research*, 60, 859–869. <https://doi.org/10.1306/d4267631-2b26-11d7-864800102c1865d>
- Osland, M. J. (2017). *Vegetation, soil, and landscape data*. <https://www.sciencebase.gov/catalog/item/5731fbace4b0dae0d5dc1e59>
- Osland, M. J., Spivak, A. C., Nestlerode, J. A., Lessmann, J. M., Almario, A. E., Heitmuller, P. T., Russell, M. J., Krauss, K. W., Alvarez, F., Dantin, D. D., Harvey, J. E., From, A. S., Cormier, N., & Stagg, C. L. (2012). Ecosystem development after mangrove wetland creation: Plant-soil change across a 20-year chronosequence. *Ecosystems*, 15, 848–866. <https://doi.org/10.1007/s10021-012-9551-1>
- Otero, X., Ferreira, T., Huerta-D, M., Partiti, C., Souza, V., Vidal-Torrado, P., & Mací, F. (2009). Geochemistry of iron and manganese in soils and sediments of a mangrove system, Island of Pai Matos (Cananeia—SP, Brazil). *Geoderma*, 148, 318–335. <https://doi.org/10.1016/j.geoderma.2008.10.016>
- Ouyang, X., & Lee, S. (2014). Updated estimates of carbon accumulation rates in coastal marsh sediments. *Biogeosciences*, 11, 5057–5071.
- Pastore, M. A., Megonigal, J. P., & Langley, J. A. (2017). Elevated CO₂ and nitrogen addition accelerate net carbon gain in a brackish marsh. *Biogeochemistry*, 133, 73–87. <https://doi.org/10.1007/s10533-017-0312-2>
- Paula, J., Ecosta, P. F., Martins, A., & Gove, D. (2001). Patterns of abundance of seagrasses and associated infaunal communities at Inhaca Island, Mozambique. *Estuarine, Coastal and Shelf Science*, 53, 307–318. <https://doi.org/10.1006/ecss.2001.0809>
- Pazi, A. M. M., Gandaseca, S., Rosli, N., Hamzah, A. H., Tindit, A. E., & Nyangon, L. (2016). Soil pH and carbon at different depth in three zones of mangrove forest in Sarawak, Malaysia. *The Malaysian Forester*, 79, 164–173.
- Pebesma, E. (2018). Simple features for R: Standardized support for spatial vector data. *The R Journal*, 10, 439–446. <https://doi.org/10.32614/RJ-2018-009>
- Peck, E. K., Wheatcroft, R. A., & Brophy, L. S. (2020a). Dataset: Controls on sediment accretion and blue carbon burial in tidal saline wetlands: Insights from the Oregon Coast, USA. https://smithsonian.figshare.com/articles/Dataset_Controls_on_sediment_accretion_and_blue_carbon_burial_in_tidal_saline_wetlands_Insights_from_the_Oregon_coast_U_S_A_11317820/2
- Peck, E. K., Wheatcroft, R. A., & Brophy, L. S. (2020b). Controls on sediment accretion and blue carbon burial in tidal saline wetlands: Insights from the Oregon Coast, USA. *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005464. <https://doi.org/10.1029/2019jg005464>
- Pedersen, M., Duarte, C., & Cebrián, J. (1997). Rate of changes in organic matter and nutrient stocks during seagrass *Cymodocea nodosa* colonization and stand development. *Marine Ecology Progress Series*, 159, 29–36. <https://doi.org/10.3354/meps159029>
- Piazza, S. C., Steyer, G. D., Cretini, K. F., Sasser, C. E., Visser, J. M., Holm, G. O., Sharp, L. A., Evers, D. E., & Meriwether, J. R. (2011). Geomorphic and ecological effects of Hurricanes Katrina and Rita on coastal Louisiana marsh communities. <https://doi.org/10.3133/ofr20111094>
- Poppe, K. (2015). *An Ecogeomorphic model to assess the response of Padilla Bay's eelgrass habitat to sea level rise*. Master's thesis, Western Washington University. <https://cedar.wwu.edu/wwuet/458/>
- Poppe, K., & Rybczyk, J. (2018). Carbon sequestration in a pacific northwest eelgrass (*Zostera marina*) meadow. *Northwest Science*, 92, 80–91. <https://doi.org/10.3955/046.092.0202>
- Poppe, K., & Rybczyk, J. (2021). Tidal marsh restoration enhances sediment accretion and carbon accumulation in the Stillaguamish River Estuary, Washington. *PLoS One*, 16, e0257244. <https://doi.org/10.1371/journal.pone.0257244>
- Poppe, K., & Rybczyk, J. (2022). Assessing the future of an intertidal seagrass meadow in response to sea level rise with a hybrid eco-geomorphic model of elevation change. *Ecological Modeling*, 469, 109975. <https://doi.org/10.1016/j.ecolmodel.2022.109975>
- Poppe, K. L., & Rybczyk, J. M. (2019). Dataset: Sediment carbon stocks and sequestration rates in the pacific northwest region of Washington, USA. https://smithsonian.figshare.com/articles/Dataset_Sediment_carbon_stocks_and_sequestration_rates_in_the_Pacific_Northwest_region_of_Washington_USA/10005248
- Poulter, B., Fluet-Chouinard, E., Hugelius, G., Koven, C., Fatoyinbo, L., Page, S. E., Rosentreter, J. A., Smart, L. S., Taillie, P. J., Thomas, N., Zhang, Z., & Wijedasa, L. S. (2021). A review of global wetland carbon stocks and management challenges. In K. W. Krauss, Z. Zhu, & C. L. Stagg (Eds.), *Wetland carbon and environmental management* (pp. 1–20). Wiley Online Library.
- Prentice, C., Poppe, K., Lutz, M., Murray, E., Stephens, T., Spooner, A., Hessing-Lewis, M., Sanders-Smith, R., Rybczyk, J., Apple, J., & Short, F. (2020). A synthesis of blue carbon stocks, sources, and accumulation rates in eelgrass (*Zostera marina*) meadows in the Northeast Pacific. *Global Biogeochemical Cycles*, 34, e2019GB006345. <https://doi.org/10.1029/2019GB006345>
- Pulich, W. M., Jr. (1987). Subtropical seagrasses and trace metals cycling. In *Florida Marine Research Publications* (Vol. 42). Florida Department of Natural Resources: Bureau of Marine Research. [https://scholar.google.com/scholar?hl=en&as_sdt=0%2C9&q=Pulich%2C+W.+M.%2C+Jr.+%2821987%29.+Subtropical+seagrasses+and+trace+metals+cycling.+In+%27Proceedings+of+a+symposium+on+subtropical-tropical+seagrasses+of+the+southeastern+United+States.+Florida+Marine+Research+Publication+No.+42%27.+%](https://scholar.google.com/scholar?hl=en&as_sdt=0%2C9&q=Pulich%2C+W.+M.%2C+Jr.+%2821987%29.+Subtropical+seagrasses+and+trace+metals+cycling.+In+%27Proceedings+of+a+symposium+on+subtropical-tropical+seagrasses+of+the+southeastern+United+States.+Florida+Marine+Research+Publication+No.+42%27.+%27)

- 28Eds+M.+J.+Durako%2C+R.+C.+Phillips+and+R.+R.+Lewis%2C+III.%29+pp.+39-52.%28Florida+Department+of+Natural+Resources%2C+Bureau+of+Marine+Research%3A+St.+Peterburgh.%29&btnG
- Qu, W., Morrison, R., West, R., & Su, C. (2006). Organic matter and benthic metabolism in Lake Illawarra, Australia. *Continental Shelf Research*, 26, 1756–1774. <https://doi.org/10.1016/j.csr.2006.05.007>
- R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Radabaugh, K. R., Moyer, R. P., Chappel, A. R., Powell, C. E., Bociu, I., Clark, B. C., & Smoak, J. M. (2017). Coastal blue carbon assessment of mangroves, salt marshes, and salt barrens in Tampa Bay, Florida, USA. *Estuaries and Coasts*, 41, 1496–1510. <https://doi.org/10.1007/s12237-017-0362-7>
- Redfield, A. C. (1972). Development of a New England salt marsh. *Ecological Monographs*, 42, 201–237.
- Rigollet, V. (2004). Seasonal evolution of heavy metal concentrations in the surface sediments of two Mediterranean *Zostera Marina* L. beds at Thau Lagoon (France) and Venice lagoon (Italy). *Bioresource Technology*, 95, 159–167. <https://doi.org/10.1016/j.biortech.2003.12.018>
- Rodríguez, M. A., & Molly, B. C. (2022). Salt marsh radiocarbon and loss on ignition data. <https://doi.org/10.6084/m9.figshare.20137649.v2>
- Rogers, K., Kelleway, J. J., Saintilan, N., Megonigal, J. P., Adams, J. B., Holmquist, J. R., Lu, M., Schile-Beers, L., Zawadzki, A., Mazumder, D., & Woodroffe, C. D. (2019). Wetland carbon storage controlled by millennial-scale variation in relative sea-level rise. *Nature*, 567, 91–95.
- Rosenfeld, J. K. (1979). Interstitial water and sediment chemistry of two cores from Florida Bay. *SEPM Journal of Sedimentary Research*, 49, 989–994. <https://doi.org/10.1306/212f7897-2b24-11d7-864800102c1865d>
- Sahu, S. C., Kumar, M., & Ravindranath, N. H. (2016). Carbon stocks in natural and planted mangrove forests of Mahanadi mangrove wetland, east coast of India. *Current Science*, 110, 2253. <https://doi.org/10.18520/cs/v110/i12/2253-2260>
- Saintilan, N., Rogers, K., Mazumder, D., & Woodroffe, C. (2013). Allochthonous and autochthonous contributions to carbon accumulation and carbon store in southeastern Australian coastal wetlands. *Estuarine, Coastal and Shelf Science*, 128, 84–92. <https://doi.org/10.1016/j.ecss.2013.05.010>
- Sanbor, P., & Darwyn, C. (2020). Dataset: Carbon data for intertidal soils and sediments, skeena river estuary, British Columbia. https://smithsonian.figshare.com/articles/Dataset_Carbon_data_for_intertidal_soils_and_sediments_Skeena_River_estuary_British_Columbia/12252005
- Sanderman, J. (2017). *Global mangrove soil carbon: Dataset and spatial maps*. <https://dataVERSE.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/OCYUIT>
- Sanderman, J., Hengl, T., Fiske, G., Solvik, K., Adame, M. F., Benson, L., Bukoski, J. J., Carnell, P., Cifuentes-Jara, M., Donato, D., Duncan, C., zu Ermgassen, P., Ewers Lewis, C. J., Macreadie, P. I., Glass, L., Gress, S., Jardine, S. L., Jones, T. G., Nsombo, E. N., ... Landis, E. (2018). A global map of mangrove forest soil carbon at 30 m spatial resolution. *Environmental Research Letters*, 13(5), 055002.
- Sanders, C. J., Eyre, B. D., Santos, I. R., Machado, W., Luiz-Silva, W., Smoak, J. M., Breithaupt, J. L., Ketterer, M. E., Sanders, L., Marotta, H., & Silva-Filho, E. (2014). Elevated rates of organic carbon, nitrogen, and phosphorus accumulation in a highly impacted mangrove wetland. *Geophysical Research Letters*, 41, 2475–2480. <https://doi.org/10.1002/2014gl059789>
- Sanders, C. J., Smoak, J. M., Naidu, A. S., & Patchineelam, S. R. (2008). Recent sediment accumulation in a mangrove forest and its relevance to local sea-level rise (Ilha Grande, Brazil). *Journal of Coastal Research*, 242, 533–536. <https://doi.org/10.2112/07-0872.1>
- Schile, L. M., Callaway, J. C., Morris, J. T., Stralberg, D., Parker, V. T., & Kelly, M. (2014). Modeling tidal marsh distribution with sea-level rise: Evaluating the role of vegetation, sediment, and upland habitat in marsh resiliency. *PLoS One*, 9, e88760.
- Schile, L. M., Callaway, J. C., Suding, K. N., & Kelly, N. M. (2017). Can community structure track sea-level rise? Stress and competitive controls in tidal wetlands. *Ecology and Evolution*, 7, 1276–1285.
- Schile, L. M., & Megonigal, J. P. (2017). *Abu Dhabi Blue Carbon Demonstration Project*. <https://repository.si.edu/handle/10088/31949>
- Serwadda, D., Ndebele, P., Grabowski, M. K., Bajunirwe, F., & Wanyenze, R. K. (2018). Open data sharing and the global south—Who benefits? *Science*, 359, 642–643.
- Sfriso, A., & Marcomini, A. (1999). Macrophyte production in a shallow coastal lagoon. Part II: Coupling with sediment, SPM and tissue carbon, nitrogen and phosphorus concentrations. *Marine Environmental Research*, 47, 285–309. [https://doi.org/10.1016/s0141-1136\(98\)00122-6](https://doi.org/10.1016/s0141-1136(98)00122-6)
- Smeaton, C., Barlow, N. L., & Austin, W. E. (2020). Coring and compaction: Best practice in blue carbon stock and burial estimations. *Geoderma*, 364, 114180.
- Smith, K. E. (2012). *Paleoecological study of coastal Marsh in the Chenier plain, Louisiana: Investigating the diatom composition of hurricane-deposited sediments and a diatom-based quantitative reconstruction of sea-level characteristics*. Ph.D. thesis, University of Florida. <https://ufdc.ufl.edu/UFE0044969/00001/pdf>
- Smith, K. E. L., Flocks, J. G., Steyer, G. D., & Piazza, S. C. (2015). *Wetland paleoecological study of southwest coastal Louisiana: Sediment cores and diatom calibration dataset*. <https://doi.org/10.3133/ds877>
- Smoak, J. M., Breithaupt, J. L., Smith, T. J., & Sanders, C. J. (2013). Sediment accretion and organic carbon burial relative to sea-level rise and storm events in two mangrove forests in Everglades National Park. *Catena*, 104, 58–66. <https://doi.org/10.1016/j.catena.2012.10.009>
- Spivak, A. (2020). *Bulk soil and elemental properties of marsh and infilled pond soils collected in 2014–2015 within plum Island ecosystems Iter*. <https://hdl.handle.net/1912/26379>
- Spivak, A., Canuel, E., Duffy, J., Douglass, J., & Richardson, J. (2009). Epifaunal community composition and nutrient addition alter sediment organic matter composition in a natural eelgrass *Zostera marina* bed: A field experiment. *Marine Ecology Progress Series*, 376, 55–67. <https://doi.org/10.3354/meps07813>
- Spruzen, F. L., Richardson, A. M. M., & Woehler, E. J. (2007). Spatial variation of intertidal macroinvertebrates and environmental variables in Robbins passage wetlands, NW Tasmania. *Hydrobiologia*, 598, 325–342. <https://doi.org/10.1007/s10750-007-9166-2>
- St. Laurent, K. A., Hribar, D. J., Carlson, A. J., Crawford, C. M., & Siok, D. (2020a). Assessing coastal carbon variability in two Delaware tidal marshes. *Journal of Coastal Conservation*, 24, 1–16. <https://doi.org/10.1007/s11852-020-00783-3>
- St. Laurent, K. A., Hribar, D. J., Carlson, A. J., Crawford, C. M., & Siok, D. (2020b). Dataset: Assessing coastal carbon variability in two Delaware tidal marshes. https://smithsonian.figshare.com/articles/dataset/Dataset_Assessing_coastal_carbon_variability_in_two_Delaware_tidal_marshes/13315472
- Stoner, A. W., Ray-Culp, M., & O'Connell, S. M. (1998). Settlement and recruitment of queen conch, *strombus gigas*, in seagrass meadows: Associations with habitat and micropredators. *Fishery Bulletin*, 96, 885–899. https://scholar.google.com/scholar?hl=en&as_sdt=0%2C9&q=Stoner%2C+A.+W%2C+Ray-Culp%2C+M.%2C+and+O%27Connell%2C+S.+M.+%281998%29.+Settlement+and+recruitment+of+queen+conch%2C+Strombus+gigas%2C+in+seagrass+meadows%3A+Associations+with+habitat+and+micropredators.+Fishery+Bulletin+96%2C+885-899.&btnG=
- Stringer, C. E., Trettin, C. C., Zarnoch, S. J., & Tang, W. (2015). Carbon stocks of mangroves within the Zambezi River delta, Mozambique.

- Forest Ecology and Management*, 354, 139–148. <https://doi.org/10.1016/j.foreco.2015.06.027>
- Tam, N. F. Y., & Wong, Y. S. (1998). Variations of soil nutrient and organic matter content in a subtropical mangrove ecosystem. *Water, Air, and Soil Pollution*, 103, 245–261.
- Tedersoo, L., Küngas, R., Oras, E., Köster, K., Eenmaa, H., Leijen, Ä., Pedaste, M., Raju, M., Astapova, A., Lukner, H., Kogermann, K., & Sepp, T. (2021). Data sharing practices and data availability upon request differ across scientific disciplines. *Scientific Data*, 8, 1–11.
- Thimdee, W., Deedin, G., Sangrungruang, C., Nishioka, J., & Matsunaga, K. (2003). Sources and fate of organic matter in Khung Krabaen Bay (Thailand) as traced by $\delta^{13}\text{C}$ and C/N atomic ratios and C/N atomic ratios. *Wetlands*, 23, 729–738.
- Thom, R. M. (1992). Accretion rates of low intertidal salt marshes in the Pacific Northwest. *Wetlands*, 12, 147–156. <https://doi.org/10.1007/bf03160603>
- Thom, R. M. (2019). Dataset: Accretion rates of low intertidal salt marshes in the pacific northwest. https://smithsonian.figshare.com/articles/Dataset_Accretion_rates_of_low_intertidal_salt_marshes_in_the_Pacific_Northwest/10046189
- Thompson, B. S., Clubbe, C. P., Primavera, J. H., Curnick, D., & Koldewey, H. J. (2014). Locally assessing the economic viability of blue carbon: A case study from Panay Island, The Philippines. *Ecosystem Services*, 8, 128–140. <https://doi.org/10.1016/j.ecoser.2014.03.004>
- Thorne, K., MacDonald, G., Guntenspergen, G., Ambrose, R., Buffington, K., Dugger, B., Freeman, C., Janousek, C., Brown, L., Rosencranz, J., Holmquist, J., Smol, J., Hargan, K., & Takekawa, J. (2018). U.S. pacific coastal wetland resilience and vulnerability to sea-level rise. *Science Advances*, 4, eaao3270. <https://doi.org/10.1126/sciadv.aao3270>
- Todd-Brown, K. E., Abramoff, R. Z., Beem-Miller, J., Blair, H. K., Earl, S., Frederick, K. J., Fuka, D. R., Guevara Santamaría, M., Harden, J. W., Heckman, K., Heran, L. J., Holmquist, J. R., Hoyt, A. M., Klinges, D. H., LeBauer, D. S., Malhotra, A., McClelland, S. C., Nave, L. E., Rocci, K. S., ... Younger, M. L. (2022). Reviews and syntheses: The promise of big diverse soil data, moving current practices towards future potential. *Biogeosciences*, 19, 3505–3522.
- Tomlinson, P. B. (2016). *The botany of mangroves*. Cambridge University Press.
- Townsend, E., & Fonseca, M. (1998). Bioturbation as a potential mechanism influencing spatial heterogeneity of North Carolina seagrass beds. *Marine Ecology Progress Series*, 169, 123–132. <https://doi.org/10.3354/meps169123>
- Trettin, C. C., Stringer, C. E., Zarnoch, S. J., Tang, W., & Dai, Z. (2017). Mangrove carbon stocks in Zambezi River delta, Mozambique. <https://doi.org/10.2737/rds-2017-0053>
- Tue, N. T., Dung, L. V., Nhuan, M. T., & Omori, K. (2014). Carbon storage of a tropical mangrove forest in Mui Ca Mau National Park, Vietnam. *Catena*, 121, 119–126. <https://doi.org/10.1016/j.catena.2014.05.008>
- Turner, R. E., Milan, C. S., & Swenson, E. M. (2006). Recent volumetric changes in salt marsh soils. *Estuarine, Coastal and Shelf Science*, 69, 352–359.
- Twilley, R. R., Pozo, M., Garcia, V. H., Rivera-Monroy, V. H., Zambrano, R., & Bodero, A. (1997). Litter dynamics in riverine mangrove forests in the Guayas River Estuary, Ecuador. *Oecologia*, 111, 109–122. <https://doi.org/10.1007/s004420050214>
- U.S. Geological Survey, & Thorne, K. (2015). *Marshes to mudflats: Climate change effects along a latitudinal gradient in the Pacific Northwest*. <https://www.sciencebase.gov/catalog/item/5006e99ee4b0abf7ce733f58>
- Unger, V., Elsey-Quirk, T., Sommerfield, C., & Velinsky, D. (2016). Stability of organic carbon accumulating in spartina alterniflora-dominated salt marshes of the mid-Atlantic U.S. *Estuarine, Coastal and Shelf Science*, 182, 179–189. <https://doi.org/10.1016/j.ecss.2016.10.001>
- USDA, NRCS. (2022). *The plants database*. <http://plants.usda.gov>
- Van Engeland, T. (2010). *Dissolved organic nitrogen dynamics in coastal ecosystems*. Ph.D. thesis, Utrecht University. <https://dspace.library.uu.nl/handle/1874/41801>
- Vasconcelos, R. W., dos Santos Gomes, V., de Lucena, D. R., da Silva, O. A., Sousa, A. C., & D'Andrea, A. F. (2014). Soil organic matter and soil acidity in mangrove areas in the River Paraíba Estuary, Cabedelo, Paraíba, Brazil. *Eurasian Journal of Soil Science*, 3, 157. <https://doi.org/10.18393/ejss.76445>
- Vaughn, D., Bianchi, T., Shields, M., Kenney, W., & Osborne, T. (2021). Dataset: Increased organic carbon burial in northern Florida mangrove-salt Marsh transition zones. https://smithsonian.figshare.com/articles/dataset/Dataset_Increased_Organic_Carbon_Burial_in_Northern_Florida_Mangrove-Salt_Marsh_Transition_Zones/10552004
- Vaughn, D. R., Bianchi, T. S., Shields, M. R., Kenney, W. F., & Osborne, T. Z. (2020). Increased organic carbon burial in northern Florida mangrove-salt marsh transition zones. *Global Biogeochemical Cycles*, 34, e2019GB006334. <https://doi.org/10.1029/2019GB006334>
- Vichkovitten, T., & Holmer, M. (2005). Dissolved and particulate organic matter in contrasting *Zostera marina* (eelgrass) sediments. *Journal of Experimental Marine Biology and Ecology*, 316, 183–201. <https://doi.org/10.1016/j.jembe.2004.11.002>
- Volkman, J. K., Revill, A. T., Holdsworth, D. G., & Fredericks, D. (2008). Organic matter sources in an enclosed coastal inlet assessed using lipid biomarkers and stable isotopes. *Organic Geochemistry*, 39, 689–710. <https://doi.org/10.1016/j.orggeochem.2008.02.014>
- Wang, G., Guan, D., Peart, M., Chen, Y., & Peng, Y. (2013). Ecosystem carbon stocks of mangrove forest in Yingluo Bay, Guangdong province of South China. *Forest Ecology and Management*, 310, 539–546. <https://doi.org/10.1016/j.foreco.2013.08.045>
- Ward, M. (2021). Data from: Organic carbon, grain size, elemental/isotopic composition. <https://doi.org/10.5061/dryad.m0cfxpp31>
- Ward, M. A., Hill, T. M., Souza, C., Filipczyk, T., Ricart, A. M., Merolla, S., Capece, L. R., O'Donnell, B. C., Elsmore, K., Oechel, W. C., & Beheshti, K. M. (2021). Blue carbon stocks and exchanges along the California coast. *Biogeosciences*, 18, 4717–4732. <https://bg.copernicus.org/articles/18/4717/2021/>
- Watson, E. B., & Byrne, R. (2013). Late holocene marsh expansion in southern San Francisco Bay, California. *Estuaries and Coasts*, 36, 643–653. <https://doi.org/10.1007/s12237-013-9598-z>
- Weis, D. A., & Anthony, D. (1999). Vertical accretion rates and heavy metal chronologies in wetland sediments of Tijuana estuary. Master's thesis, San Diego State University.
- Weis, D. A., Callaway, J. C., & Gersberg, R. M. (2001). Vertical accretion rates and heavy metal chronologies in wetland sediments of the Tijuana Estuary. *Estuaries*, 24, 840. <https://doi.org/10.2307/1353175>
- Weston, N. B., Rodriguez, E., Donnelly, B., Solohin, E., Jezycki, K., Demberger, S., Sutter, L., Morris, J. T., Neubauer, S. C., & Craft, C. B. (2022). Dataset: Recent acceleration of coastal wetland accretion along the U.S. East Coast. https://smithsonian.figshare.com/articles/dataset/Dataset_Recent_Acceleration_of_Coastal_Wetland_Accretion_Along_the_U_S_East_Coast/13043054
- White, J. R., Sapkota, Y., Chambers, L. G., Cook, R. L., & Xue, Z. (2020). Biogeochemical properties of sediment cores from Barataria Basin, Louisiana, 2018 and 2019. <https://hdl.handle.net/1912/26484>
- Wickham, H. (2014). Tidy data. *Journal of Statistical Software*, 59, 1–23.
- Wigand, C., Eagle, M., Branoff, B. L., Balogh, S., Miller, K. M., Martin, R. M., Hanson, A., Oczkowski, A. J., Huertas, E., Loffredo, J., & Watson, E. B. (2021). Recent carbon storage and burial exceed historic rates in the San Juan Bay estuary peri-urban mangrove forests (Puerto Rico, United States). *Frontiers in Forests and Global Change*, 4, 67. <https://www.frontiersin.org/article/10.3389/ffgc.2021.676691>
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B.,

- Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., ... Mons, B. (2016). The fair guiding principles for scientific data management and stewardship. *Scientific Data*, 3, 1–9.
- Wilson, G., Bryan, J., Cranston, K., Kitzes, J., Nederbragt, L., & Teal, T. K. (2017). Good enough practices in scientific computing. *PLoS Computational Biology*, 13, e1005510.
- Windham-Myers, L., Marvin-DiPasquale, M. C., Agee, J. L., Kieu, L. H., Kakourous, E., Erikson, L. H., & Ward, K. (2010). Biogeochemical processes in an urban, restored wetland of San Francisco Bay, California, 2007–2009 methods and data for plant, sediment and water parameters. <https://doi.org/10.3133/ofr20101299>
- Woodroffe, D. C. (1985). Studies of a mangrove basin, Tuff Crater, New Zealand: I. Mangrove biomass and production of detritus. *Estuarine, Coastal and Shelf Science*, 20, 265–280.
- Wooller, M., Smallwood, B., Jacobson, M., & Marilyn, F. (2003). Carbon and nitrogen stable isotopic variation in *Laguncularia racemosa* (L.) (White mangrove) from Florida and Belize: Implications for trophic level studies. *Hydrobiologia*, 499, 13–23. <https://doi.org/10.1023/A:1026339517242>
- Wylie, L., Sutton-Grier, A. E., & Moore, A. (2016). Keys to successful blue carbon projects: Lessons learned from global case studies. *Marine Policy*, 65, 76–84.
- Xia, P., Meng, X., Li, Z., Feng, A., Yin, P., & Zhang, Y. (2015). Mangrove development and its response to environmental change in Yingluo Bay (SW China) during the last 150 years: Stable carbon isotopes and mangrove pollen. *Organic Geochemistry*, 85, 32–41. <https://doi.org/10.1016/j.orggeochem.2015.04.003>
- Xu, H. F., Zhang, P. J., Zhang, M. Q., Lin, Y. K., & Lin, X. G. (2012). Carbon storage of three natural mangrove forests in Gaoqiao, Zhanjiang. *Value Engineering*, 15, e00575.
- Yamamoto, M., Koike, I., & Izumi, H. (1993). Partitioning of the nitrogen stock in the vicinity of a Fijian seagrass bed dominated by *Syringodium isoetifolium* (Ascherson) Dandy. *Marine and Freshwater Research*, 44, 101. <https://doi.org/10.1071/mf9930101>
- Yando, E. S., Osland, M. J., Willis, J. M., Day, R. H., Krauss, K. W., & Hester, M. W. (2016). Salt marsh-mangrove ecotones: Using structural gradients to investigate the effects of woody plant encroachment on plant-soil interactions and ecosystem carbon pools. *Journal of Ecology*, 104, 1020–1031. <https://doi.org/10.1111/1365-2745.12571>
- Yang, J., Gao, J., Cheung, A., Liu, B., Schwendenmann, L., & Costello, M. J. (2013). Vegetation and sediment characteristics in an expanding mangrove forest in New Zealand. *Estuarine, Coastal and Shelf Science*, 134, 11–18. <https://doi.org/10.1016/j.ecss.2013.09.017>
- Yarbro, L. A., & Carlson, P. R. (2008). Community oxygen and nutrient fluxes in seagrass beds of Florida Bay, USA. *Estuaries and Coasts*, 31, 877–897. <https://doi.org/10.1007/s12237-008-9071-6>
- Zhang, J.-P., Shen, C.-D., Ren, H., Wang, J., & Han, W.-D. (2012). Estimating change in sedimentary organic carbon content during mangrove restoration in southern China using carbon isotopic measurements. *Pedosphere*, 22, 58–66. [https://doi.org/10.1016/s1000-0160\(11\)60191-4](https://doi.org/10.1016/s1000-0160(11)60191-4)

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Holmquist, J. R., Klinges, D., Lonneman, M., Wolfe, J., Boyd, B., Eagle, M., Sanderman, J., Todd-Brown, K., Belshe, E. F., Brown, L. N., Chapman, S., Corstanje, R., Janousek, C., Morris, J. T., Noe, G., Rovai, A., Spivak, A., Vahsen, M., Windham-Myers, L. ... Megonigal, J. P. (2024). The Coastal Carbon Library and Atlas: Open source soil data and tools supporting blue carbon research and policy. *Global Change Biology*, 30, e17098. <https://doi.org/10.1111/gcb.17098>