

KOLMOGOROV–SMIRNOV STATISTICS AND THE
ANALYTIC PROPERTIES OF DIRICHLET SERIES
ASSOCIATED TO ELLIPTIC CURVES

A Dissertation

Presented to the Faculty of the Graduate School

of Cornell University

in Partial Fulfillment of the Requirements for the Degree of

Doctor of Philosophy

by

Daniel Miller

May 2017

© 2017 Daniel Miller
ALL RIGHTS RESERVED

KOLMOGOROV-SMIRNOV STATISTICS AND THE ANALYTIC PROPERTIES OF
DIRICHLET SERIES ASSOCIATED TO ELLIPTIC CURVES

Daniel Miller, Ph.D.
Cornell University 2017

Abstract here.

BIOGRAPHICAL SKETCH

Brief biographical sketch.

Dedication here.

ACKNOWLEDGEMENTS

Some acknowledgments.

TABLE OF CONTENTS

1	Introduction	1
2	Discrepancy	2
2.1	Definitions and first results	2
2.2	Comparing sequences	2
2.3	Combining sequences	2
3	Strange Dirichlet series	3
4	Irrationality exponents	4
5	Deformation theory	5
5.1	Category of test objects	5
5.2	Quotients in the flat topology	6
5.3	Groupoids and quotient stacks	8
5.4	Deformations of group representations	8
5.5	Tangent spaces and obstruction theory	9
6	Constructing Galois representations	11
7	First counterexample	12
8	Second counterexample	13
9	Computational evidence for the Akiyama–Tanigawa conjecture	14
10	Concluding remarks and future directions	15

CHAPTER 1
INTRODUCTION

CHAPTER 2

DISCREPANCY

2.1 Definitions and first results

regular discrepancy
star discrepancy
Euclidean space vs. torus

2.2 Comparing sequences

If $\{x_n\} \subset [0, \pi/2)$ has some discrepancy with respect to some measure, then the “flipped” sequence $\{\pi/2 - x_n\}$ has the same discrepancy with respect to the “flipped” measure.

2.3 Combining sequences

If $\{x_n\}$ and $\{y_n\}$ are sequences supported on $[0, \pi/2)$ and $[\pi/2, \pi)$ respectively, and both are equidistributed with respect to measures supported on their respective intervals, then the “interleaved” sequence $(x_1, y_1, x_2, y_2, \dots)$ also has equidistribution (with respect to the combined measure) and discrepancy which decays no faster than the slower of the two.

CHAPTER 3
STRANGE DIRICHLET SERIES

CHAPTER 4
IRRATIONALITY EXPONENTS

CHAPTER 5 DEFORMATION THEORY

5.1 Category of test objects

The following is an exposition and explication of the theory outlined in [SGA 3₁, VII_B, §0–1]. In particular, we will heavily use the notions of a pseudocompact ring, pseudocompact modules, etc. Let Λ be a pseudocompact ring. Write \mathbf{C}_Λ for the opposite of the category of Λ -algebras which have finite length as Λ -modules. Given such a Λ -algebra A , write $X = \mathrm{Spf}(A)$ for the corresponding object of \mathbf{C}_Λ , and we put $A = \mathcal{O}(X)$.

Lemma 5.1.1. *Let Λ be a pseudocompact ring, \mathbf{C}_Λ as above. Then \mathbf{C}_Λ is closed under finite limits and colimits.*

Lemma 5.1.2. *Let Λ be a pseudocompact local ring. Then Λ is henselian, in any of the following senses:*

1. *d*

Proof. [EGA 4₄, 18.5.?] □

Following Grothendieck, if \mathcal{C} is an arbitrary category, we write $\widehat{\mathcal{C}} = \mathrm{hom}(\mathcal{C}^\circ, \mathbf{Set})$ for the category of contravariant functors $\mathcal{C} \rightarrow \mathbf{Set}$. We regard \mathcal{C} as a full subcategory of $\widehat{\mathcal{C}}$ via the Yoneda embedding, so for $X, Y \in \mathcal{C}$, we write $X(Y) = \mathrm{hom}_{\mathcal{C}}(Y, X)$. With this notation, the Yoneda Lemma states that $\mathrm{hom}_{\widehat{\mathcal{C}}}(X, P) = P(X)$ for all $X \in \mathcal{C}$.

Lemma 5.1.3. *Let $\mathcal{X} \in \widehat{\mathbf{C}_\Lambda}$. Then \mathcal{X} is left exact if and only if there exists a filtered system $\{X_i\}_{i \in I}$ in \mathbf{C}_Λ together with a natural isomorphism $\mathcal{X}(\cdot) \simeq \varinjlim X_i(\cdot)$. Write $\mathrm{Ind}(\mathbf{C}_\Lambda)$ for the category of such functors. Then $\mathrm{Ind}(\mathbf{C}_\Lambda)$ is closed under colimits, and the Yoneda embedding $\mathbf{C}_\Lambda \hookrightarrow \mathrm{Ind}(\mathbf{C}_\Lambda)$ preserves filtered colimits.*

Proof. This follows from the results of [KS06, 6.1]. □

If R is a pseudocompact Λ -algebra, write $\mathrm{Spf}(R)$ for the object of $\widehat{\mathbf{C}_\Lambda}$ defined by $\mathrm{Spf}(R)(A) = \mathrm{hom}_{\mathrm{cts}/\Lambda}(R, A)$, the set of continuous Λ -algebra homomorphisms.

Lemma 5.1.4. *The functor Spf induces an (anti-)equivalence between the category of pseudocompact Λ -algebras and $\mathrm{Ind}(\mathbf{C}_\Lambda)$.*

Proof. This is [SGA 3₁, VII_B 0.4.2 Prop.]. □

So $\mathrm{Ind}(\mathbf{C}_\Lambda)$ is the category of pro-representable functors on finite length Λ -algebras. *Warning:* in many papers, for example the foundational [Maz97], one reserves the term *pro-representable* for functors of the form $\mathrm{Spf}(R)$, where R is *noetherian*. We do not make this restriction.

Lemma 5.1.5. *The category $\mathrm{Ind}(\mathbf{C}_\Lambda)$ is an exponential ideal in $\widehat{\mathbf{C}_\Lambda}$.*

Proof. By this we mean the following. Let $\mathcal{X} \in \mathrm{Ind}(\mathbf{C}_\Lambda)$, $P \in \widehat{\mathbf{C}_\Lambda}$. Then the functor \mathcal{X}^P defined by

$$\mathcal{X}^P(S) = \mathrm{hom}_{\widehat{\mathbf{C}_\Lambda/S}}(P/S, \mathcal{X}/S)$$

is also in $\mathrm{Ind}(\mathbf{C}_\Lambda)$. Given the characterization of $\mathrm{Ind}(\mathbf{C}_\Lambda)$ as left exact functors, this is easy to prove, see e.g. [Joh02, 4.2.3]. □

If \mathcal{C} is a category, we write $\mathbf{Gp}(\mathcal{C})$ for the category of group objects in \mathcal{C} .

Corollary 5.1.6. *Let $\Gamma \in \mathbf{Gp}(\widehat{\mathbf{C}}_\Lambda)$ and $\mathcal{G} \in \mathbf{Gp}(\mathbf{Ind}(\mathbf{C}_\Lambda))$, then the functor $[\Gamma, \mathcal{G}]$ defined by*

$$[\Gamma, \mathcal{G}](S) = \mathrm{hom}_{\mathbf{Gp}/S}(\Gamma/S, \mathcal{G}/S)$$

is in $\mathbf{Ind}(\mathbf{C}_\Lambda)$. In particular, if Γ is a profinite group, then the functor

$$[\Gamma, \mathcal{G}](S) = \mathrm{hom}_{\mathbf{cts}/\mathbf{Gp}}(\Gamma, \mathcal{G}(S))$$

is in $\mathbf{Ind}(\mathbf{C}_\Lambda)$.

Proof. The first claim follows easily from 5.1.5. Just note that $[\Gamma, \mathcal{G}]$ is the equalizer:

$$[\Gamma, \mathcal{G}] \longrightarrow \mathcal{G}^\Gamma \xrightarrow[m_{\mathcal{G}*}]{m_\Gamma^*} \mathcal{G}^{\Gamma \times \Gamma},$$

that is, those $f: \Gamma \rightarrow \mathcal{G}$ such that $f \circ m_\Gamma = m_{\mathcal{G}} \circ (f \times f)$. The latter claim is just a special case. \square

5.2 Quotients in the flat topology

If Λ is a pseudocompact ring, the category $\mathbf{Ind}(\mathbf{C}_\Lambda)$ has nice “geometric” properties. However, for operations like taking quotients, we will embed it into the larger category $\mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$ of flat sheaves. We call a collection $\{U_i \rightarrow X\}$ of morphisms in \mathbf{C}_Λ a *flat cover* if each ring map $\mathcal{O}(X) \rightarrow \mathcal{O}(U_i)$ is flat, and moreover $\mathcal{O}(X) \rightarrow \prod \mathcal{O}(U_i)$ is faithfully flat. By [SGA 3₁, IV 6.3.1], this is a subcanonical Grothendieck topology on \mathbf{C}_Λ . We call it the *flat topology*, even though finite presentation comes for free because all the rings are finite length.

Lemma 5.2.1. *Let $\mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$ be the category of sheaves (of sets) on \mathbf{C}_Λ with respect to the flat topology. Then a presheaf $P \in \widehat{\mathbf{C}}_\Lambda$ lies in $\mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$ if and only if $P(\coprod U_i) = \prod P(U_i)$ and moreover, whenever $U \rightarrow X$ is a flat cover where $\mathcal{O}(U)$ and $\mathcal{O}(X)$ are local rings, the sequence*

$$P(X) \longrightarrow P(U) \rightrightarrows P(U \times_X U).$$

is exact. Moreover, $\mathbf{Ind}(\mathbf{C}_\Lambda) \subset \mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$.

Proof. The first claim is the content of [SGA 3₁, IV 6.3.1(ii)]. For the second, note that any $\mathcal{X} \in \mathbf{Ind}(\mathbf{C}_\Lambda)$ will, by 5.1.3, convert (arbitrary) colimits into limits. Thus $\mathcal{X}(\coprod U_i) = \prod \mathcal{X}(U_i)$. If $U \rightarrow X$ is a flat cover, then by (loc. cit.), $U \times_X U \rightrightarrows U \rightarrow X$ is a coequalizer diagram in \mathbf{C}_Λ , hence $\mathcal{X}(X) \rightarrow \mathcal{X}(U) \rightrightarrows \mathcal{X}(U \times_X U)$ is an equalizer. \square

Our main reason for introducing the category $\mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$ is that, as a (Grothendieck) topos, it is closed under arbitrary colimits. Recall that in an *equivalence relation* in $\widehat{\mathbf{C}}_\Lambda$ is a morphism $R \rightarrow X \times X$ such that, for all S , the map $R(S) \rightarrow X(S) \times X(S)$ is an injection whose image is an equivalence relation on $X(S)$. We define the quotient X/R to be the coequalizer

$$R \rightrightarrows X \longrightarrow X/R.$$

By Giraud’s Theorem [MLM94, App.], for any $S \in \mathbf{C}_\Lambda$, the natural map $X(S)/R(S) \rightarrow (X/R)(S)$ is injective. It will not be surjective in general.

We let $\mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$ inherit definitions from \mathbf{C}_Λ as follows. If P is a property of maps in \mathbf{C}_Λ (for example, “flat,” or “smooth,”) and $f: X \rightarrow Y$ is a morphism in $\mathbf{Sh}_\mathfrak{fl}(\mathbf{C}_\Lambda)$, we say that f has P if for all $S \in \mathbf{C}_\Lambda$ and $y \in Y(S)$, the pullback $X_S = X \times_Y S$ lies in \mathbf{C}_Λ , and the pullback map $X_S \rightarrow S$ has property P . For example, if $X = \mathrm{Spf}(R')$ and $Y = \mathrm{Spf}(R)$, then $X \rightarrow Y$ has property P if and only if for all finite length A and continuous Λ -algebra maps $R \rightarrow A$, the induced map $A \rightarrow R' \otimes_R A$ has P .

Theorem 5.2.2. *Let $\mathcal{R} \rightarrow \mathcal{X} \times \mathcal{X}$ be an equivalence relation in $\mathrm{Ind}(\mathbf{C}_\Lambda)$ such that one of the maps $\mathcal{R} \rightarrow \mathcal{X}$ is flat. Then the quotient \mathcal{X}/\mathcal{R} lies in $\mathrm{Ind}(\mathbf{C}_\Lambda)$, and $\mathcal{X} \rightarrow \mathcal{X}/\mathcal{R}$ is a flat cover.*

Proof. This is [SGA 3_I, VII_B 1.4]. \square

By [Mat89, 29.7], if k is a field and R is a complete regular local k -algebra, then $R \simeq k[[t_1, \dots, t_n]]$. In particular, R admits an augmentation $\epsilon: R \rightarrow k$. There is a general analogue of this result, but first we need a definition.

Definition 5.2.3. *A map $f: \mathcal{X} \rightarrow \mathcal{Y}$ in $\mathrm{Ind}(\mathbf{C}_\Lambda)$ is a residual isomorphism if for all $S = \mathrm{Spf}(k) \in \mathbf{C}_\Lambda$ where k is a field, the map $f: \mathcal{X}(S) \rightarrow \mathcal{Y}(S)$ is a bijection.*

Lemma 5.2.4. *Let $f: \mathcal{X} \rightarrow \mathcal{Y}$ be a smooth map in $\mathrm{Ind}(\mathbf{C}_\Lambda)$ that is a residual isomorphism. Then f admits a section.*

Proof. By [SGA 3_I, VII_B 0.1.1], it suffices to prove the result when $\mathcal{X} = \mathrm{Spf}(R')$, $\mathcal{Y} = \mathrm{Spf}(R)$, for local Λ -algebras $R \rightarrow R'$ with the same residue field. Let $k = R/\mathfrak{m}_R \xrightarrow{\sim} R'/\mathfrak{m}_{R'}$ be their common residue field. From the diagram

$$\begin{array}{ccc} R' & \cdots \cdots \rightarrow & R \\ \uparrow & \searrow & \downarrow \\ R & \longrightarrow & k, \end{array}$$

the definition of (formal) smoothness, and a limiting argument involving the finite length quotients R/\mathfrak{a} , we obtain the result. \square

Corollary 5.2.5. *Let $\mathcal{R} \rightarrow \mathcal{X} \times \mathcal{X}$ be an equivalence relation satisfying the hypotheses of 5.2.2. Suppose further that*

1. *One of the maps $\mathcal{R} \rightarrow \mathcal{X}$ is smooth, and*
2. *The projection $\mathcal{X} \rightarrow \mathcal{X}/\mathcal{R}$ is a residual isomorphism.*

Then $\mathcal{X} \rightarrow \mathcal{X}/\mathcal{R}$ admits a section, so $\mathcal{X}(S)/\mathcal{R}(S) \xrightarrow{\sim} (\mathcal{X}/\mathcal{R})(S)$ for all $S \in \mathbf{C}_\Lambda$.

Proof. By 5.2.4, it suffices to prove that $\mathcal{X} \rightarrow \mathcal{X}/\mathcal{R}$ is smooth. By [EGA 4₄, 17.7.3(ii)], smoothness can be detected after flat descent. So base-change with respect to the projection $\mathcal{X} \rightarrow \mathcal{X}/\mathcal{R}$. In the following commutative diagram

$$\begin{array}{ccc} \mathcal{R} & \xrightarrow{\quad} & \mathcal{X} \\ \parallel & \searrow & \downarrow \\ \mathcal{X} \times_{\mathcal{X}/\mathcal{R}} \mathcal{X} & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \mathcal{X} & \longrightarrow & \mathcal{X}/\mathcal{R} \end{array}$$

we can ensure the smoothness of $\mathcal{R} \rightarrow \mathcal{X}$ by our hypotheses. Since $\mathcal{X} \rightarrow \mathcal{X}/\mathcal{R}$ is smooth after flat base-change, the original map is smooth. \square

Example 5.2.6. The hypothesis on residue fields in 5.2.5 is necessary. To see this, let $\Lambda = k$ be a field, $k \hookrightarrow K$ a finite Galois extension with Galois group G . Then $G \times \mathrm{Spf}(K) \rightrightarrows \mathrm{Spf}(K)$ has quotient $\mathrm{Spf}(k)$, but the map $\mathrm{Spf}(K)(S) \rightarrow \mathrm{Spf}(k)(S)$ is *not* surjective for all $S \in \mathbf{C}_k$, e.g. it is not for $S = \mathrm{Spf}(k)$.

Example 5.2.7. The hypothesis of smoothness in 5.2.5 is necessary. To see this, let k be a field of characteristic $p > 0$. Then the formal additive group $\widehat{\mathbf{G}}_a = \mathrm{Spf}(k[[t]])$ has a subgroup α_p defined by

$$\alpha_p(S) = \{s \in \mathcal{O}(S) : s^p = 0\}.$$

The quotient $\widehat{\mathbf{G}}_a/\alpha_p$ has as affine coordinate ring $k[[t^p]]$. In particular, the following sequence is exact in the flat topology:

$$0 \longrightarrow \alpha_p \longrightarrow \widehat{\mathbf{G}}_a \xrightarrow{(\cdot)^p} \widehat{\mathbf{G}}_a \longrightarrow 0.$$

It follows that $\alpha_p \times \widehat{\mathbf{G}}_a \rightrightarrows \widehat{\mathbf{G}}_a \xrightarrow{(\cdot)^p} \widehat{\mathbf{G}}_a$ is a coequalizer in $\mathrm{Sh}_{\mathrm{fl}}(\mathbf{C}_k)$ satisfying all the hypotheses of 5.2.5 except smoothness. And indeed, as one sees by letting $S = \mathrm{Spf}(A)$ for any non-perfect k -algebra A , the map $(\cdot)^p : \widehat{\mathbf{G}}_a(S) \rightarrow \widehat{\mathbf{G}}_a(S)$ is *not* surjective for all S .

5.3 Groupoids and quotient stacks

Lemma 5.3.1. *Let $\mathcal{G} \in \mathrm{Ind}(\mathbf{C}_\Lambda)$ be a smooth connected group. Then every \mathcal{G} -torsor is trivial.*

Proof. Let $\mathcal{P} \rightarrow \mathcal{B}$ be a \mathcal{G} -torsor in $\mathrm{Ind}(\mathbf{C}_\Lambda)$. That is, \mathcal{P} has an action of \mathcal{G}_S for which $\mathcal{P} \times_{\mathcal{B}} \mathcal{P} \simeq \mathcal{G} \times \mathcal{P}$ as \mathcal{G} -spaces. [...not done...] \square

Theorem 5.3.2. *Let \mathcal{G} be a smooth connected group in $\mathrm{Ind}(\mathbf{C}_\Lambda)$, and $\mathcal{X} \in \mathrm{Ind}(\mathbf{C}_\Lambda)$ a \mathcal{G} -object. Then the quotient stack $[\mathcal{X}/\mathcal{G}](S)$ has as objects $\mathcal{X}(S)/\mathcal{G}(S)$, but with extra automorphisms?*

Proof. Use triviality of torsors. \square

5.4 Deformations of group representations

Let $\Gamma \in \mathrm{Gp}(\widehat{\mathbf{C}}_\Lambda)$ and $\mathcal{G} \in \mathrm{Ind}(\mathbf{C}_\Lambda)$. By 5.1.6, the functor

$$\mathrm{Rep}^\square(\Gamma, \mathcal{G})(S) = \mathrm{hom}_{\mathrm{Gp}/S}(\Gamma_S, \mathcal{G}_S)$$

is in $\mathrm{Ind}(\mathbf{C}_\Lambda)$. We would like to define an ind-scheme $\mathrm{Rep}(\Gamma, \mathcal{G})$ as “ $\mathrm{Rep}^\square(\Gamma, \mathcal{G})$ modulo conjugation,” but this requires some care. The conjugation action of \mathcal{G} on $\mathrm{Rep}^\square(\Gamma, \mathcal{G})$ will have fixed points, so the quotient will be badly behaved. We loosely follow [Til96].

Assume Λ is local, with maximal ideal \mathfrak{m} and residue field \mathbf{k} . Fix $\bar{\rho} \in \mathrm{Rep}^\square(\Gamma, \mathcal{G})(\mathbf{k})$, i.e. a residual representation $\bar{\rho} : \Gamma \rightarrow \mathcal{G}(\mathbf{k})$. Let $\mathrm{Rep}^\square(\Gamma, \mathcal{G})_{\bar{\rho}}$ be the connected component of $\bar{\rho}$ in $\mathrm{Rep}^\square(\Gamma, \mathcal{G})$. Assume that \mathcal{G} and $Z(\mathcal{G})$ are smooth; then the quotient $\mathcal{G}^{\mathrm{ad}} = \mathcal{G}/Z(\mathcal{G})$ is also smooth. Let $\mathcal{G}^{\mathrm{ad}, \circ}$ be the connected component of 1 in $\mathcal{G}^{\mathrm{ad}}$.

Theorem 5.4.1. *Suppose $(\Lambda, \mathfrak{m}, \mathbf{k})$ is local. If $\mathcal{X}, \mathcal{Y} \in \mathrm{Ind}(\mathbf{C}_\Lambda)$ are connected and $\mathcal{X}(\mathbf{k}) \neq \emptyset$, then $\mathcal{X} \times_\Lambda \mathcal{Y}$ is connected.*

Proof. We are reduced to proving the following result from commutative algebra: if R, S are local pro-artinian Λ -algebras and R has residue field \mathbf{k} , then $R \widehat{\otimes}_\Lambda S$ is local. Since $R \widehat{\otimes}_\Lambda S = \varprojlim (R/\mathfrak{r}) \otimes_\Lambda (S/\mathfrak{s})$, \mathfrak{r} (resp. \mathfrak{s}) ranges over all open ideals in R (resp. S), we may assume that both R and S are artinian. The rings R and S are henselian, so $R \otimes S$ is local if and only if $(R/\mathfrak{m}_R) \otimes (S/\mathfrak{m}_S) = S/\mathfrak{m}_S$ is local, which it is. \square

We conclude that the action of $\mathcal{G}^{\mathrm{ad}, \circ}$ on $\mathrm{Rep}^\square(\Gamma, \mathcal{G})$ preserves $\mathrm{Rep}^\square(\Gamma, \mathcal{G})_{\bar{\rho}}$. Thus we may put

$$\mathrm{Rep}(\Gamma, \mathcal{G})_{\bar{\rho}} = [\mathrm{Rep}^\square(\Gamma, \mathcal{G})_{\bar{\rho}}/\mathcal{G}^{\mathrm{ad}, \circ}].$$

If $\mathcal{G}^{\mathrm{ad}, \circ}$ acts faithfully on $\mathrm{Rep}^\square(\Gamma, \mathcal{G})_{\bar{\rho}}$, then we recover the classical notion of the deformation functor.

Theorem 5.4.2. *Let Γ be a profinite group, $\bar{\rho}: \Gamma \rightarrow \mathcal{G}(\mathbf{k})$ a representation with $H^0(\Gamma, \text{Ad } \bar{\rho}) = 0$. Then $\text{Rep}(\Gamma, \mathcal{G})_{\bar{\rho}}$ exists and is what you expect.*

Proof. Need assumptions on $Z(\mathcal{G})$, \mathcal{G} should be smooth.

Need $Z(\mathcal{G}) = \ker(\mathcal{G} \rightarrow \text{GL}(\mathfrak{g}))$ in connected case. This should use $\mathfrak{g} = \text{Lie}(\text{Aut } \mathcal{G})$, via deviations in [SGA 3₁]. \square

[... local conditions]

5.5 Tangent spaces and obstruction theory

For $S_0 \in \mathbf{C}_\Lambda$, let Ex_{S_0} be the category of square-zero thickenings of S_0 . An object of Ex_{S_0} is a closed embedding $S_0 \hookrightarrow S$ whose ideal of definition has square zero. Should be “exponential exact sequence”

$$0 \longrightarrow \mathfrak{g}(I) \longrightarrow \mathcal{G}(S) \longrightarrow \mathcal{G}(S_0) \longrightarrow 1$$

This gives us a class $\text{exp} \in H^2(\mathcal{G}(S_0), \mathfrak{g}(I))$. For $\rho_0: \Gamma \rightarrow \mathcal{G}(S_0)$, the obstruction class is $o(\rho_0, I) = \rho_0^*(\text{exp}) \in H^2(\Gamma, \mathfrak{g}(I))$. It's easy to check that $o(\rho_0, I) = 0$ if and only if ρ_0 lifts to ρ . So obstruction theory naturally for $\text{Rep}^\square(\Gamma, \mathcal{G})$.

[Use [Wei94, 6.6.4]. Given setting as above, $\rho_0^*(\text{exp})$ is the pullback by ρ_0 :

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{g}(I) & \longrightarrow & \mathcal{G}(S) \times_{\mathcal{G}(S_0)} \Gamma & \longrightarrow & \Gamma \longrightarrow 1 \\ & & \parallel & & \downarrow & & \downarrow \rho_0 \\ 0 & \longrightarrow & \mathfrak{g}(I) & \longrightarrow & \mathcal{G}(S) & \longrightarrow & \mathcal{G}(S_0) \longrightarrow 1 \end{array}$$

Computing explicitly, we see the result.]

Proposition 5.5.1. *Let $f: G \rightarrow H$ be a morphism of profinite groups. Suppose M is a discrete H -module and $c \in H^2(H, M)$ corresponds to the extension*

$$0 \longrightarrow M \longrightarrow \tilde{H} \longrightarrow H \longrightarrow 1.$$

*Then $f^*c = 0$ in $H^2(G, M)$ if and only if there is a map $\tilde{f}: G \rightarrow \tilde{H}$ making the following diagram commute:*

$$\begin{array}{ccc} & & \tilde{H} \\ & \nearrow \tilde{f} & \downarrow \\ G & & H \\ & \searrow f & \end{array}$$

Proof. By [Wei94, 6.6.4], the class f^*c corresponds to the pullback diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M & \longrightarrow & G \times_H \tilde{H} & \longrightarrow & G \longrightarrow 1 \\ & & \parallel & & \downarrow & & \downarrow f \\ 0 & \longrightarrow & M & \longrightarrow & \tilde{H} & \longrightarrow & H \longrightarrow 1. \end{array}$$

Writing explicitly what it means for $G \times_H \tilde{H} \rightarrow G$ to split yields the result. \square

Let $\mathcal{X} \in \text{Ind}(\mathbf{C}_\Lambda)$ be smooth, and $L_{\mathcal{X}/\Lambda} \simeq \Omega_{\mathcal{X}/\Lambda}^1[0]$ be its cotangent complex. Fix $x_0 \in \mathcal{X}(S_0)$. From the chain $S_0 \xrightarrow{x_0} \mathcal{X} \rightarrow \text{Spf}(\Lambda)$, we get a distinguished triangle [Ill71, II 2.1.5.6]

$$x_0^* L_{\mathcal{X}/\Lambda} \longrightarrow L_{S_0/\Lambda} \longrightarrow L_{S_0/\mathcal{X}} \longrightarrow .$$

If I is a coherent sheaf on S_0 , we get a long exact sequence:

$$\mathrm{Ext}^0(\mathrm{L}_{S_0/\Lambda}, M) \rightarrow \mathrm{Ext}^0(x_0^* \mathrm{L}_{\mathcal{X}/\Lambda}, M) \rightarrow \mathrm{Ext}^1(\mathrm{L}_{S_0/\mathcal{X}}, M) \rightarrow \mathrm{Ext}^1(\mathrm{L}_{S_0/\Lambda}, M) \rightarrow \mathrm{Ext}^1(x_0^* \mathrm{L}_{\mathcal{X}/\Lambda}, M)$$

If \mathcal{X}/Λ is smooth, then $\mathrm{Ext}^1(x_0^* \mathrm{L}_{\mathcal{X}/\Lambda}, M) = 0$ and $\mathrm{L}_{\mathcal{X}/\Lambda} = \Omega_{\mathcal{X}/\Lambda}^1$. This gives us an exact sequence

$$\mathrm{Ext}^0(\mathrm{L}_{S_0/\Lambda}, M) \rightarrow \mathrm{hom}(\Omega_{\mathcal{X}/\Lambda}^1, M) \rightarrow \mathrm{Ext}^1(\mathrm{L}_{S_0/\mathcal{X}}, M) \rightarrow \mathrm{Ext}^1(\mathrm{L}_{S_0/\Lambda}, M) \rightarrow 0.$$

The result [Ill71, III 2.1.7] tells us that the choice of $S \in \mathrm{Ex}_{S_0}(M)$ gives us an element of $\mathrm{Ext}^1(\mathrm{L}_{S_0/\Lambda}, M)$. Its fiber admits an action of $\mathrm{hom}(\Omega_{\mathcal{X}/\Lambda}^1, M)$. The only thing remaining is: we need $\mathrm{Ext}^0(\mathrm{L}_{S_0/\Lambda}, M) = 0$, which doesn't hold in complete generality.

CHAPTER 6
CONSTRUCTING GALOIS REPRESENTATIONS

CHAPTER 7
FIRST COUNTEREXAMPLE

CHAPTER 8
SECOND COUNTEREXAMPLE

CHAPTER 9
COMPUTATIONAL EVIDENCE FOR THE AKIYAMA–TANIGAWA
CONJECTURE

CHAPTER 10
CONCLUDING REMARKS AND FUTURE DIRECTIONS

BIBLIOGRAPHY

- [EGA 4₄] Alexandre Grothendieck. *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV*. 32. 1967.
- [SGA 3₁] Alexandre Grothendieck and Michel Demazure, eds. *Schémas en groupes (SGA 3). Tome I. Propriétés générales des schémas en groupes*. Vol. 151. Lecture Notes in Mathematics. Springer-Verlag, 1970.
- [Ill71] Luc Illusie. *Complexe cotangent et déformations. I*. Vol. 239. Lecture Notes in Mathematics. Springer-Verlag, 1971.
- [Joh02] Peter Johnstone. *Sketches of an elephant: a topos theory compendium*. Vol. 44, 45. Oxford Logic Guides. Oxford University Press, 2002.
- [KS06] Massaki Kashiwara and Pierre Schapira. *Categories and sheaves*. Vol. 332. Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, 2006.
- [MLM94] Saunders Mac Lane and Ieke Moerdijk. *Sheaves in geometry and logic*. Second. Universitext. A first introduction to topos theory. Springer-Verlag, 1994.
- [Mat89] Hideyuki Matsumura. *Commutative ring theory*. Second. Vol. 8. Cambridge Studies in Advanced Mathematics. Translated from the Japanese by M. Reid. Cambridge University Press, 1989.
- [Maz97] Barry Mazur. “An introduction to the deformation theory of Galois representations”. In: *Modular forms and Fermat’s last theorem (Boston, MA, 1995)*. New York: Springer, 1997, pp. 243–311.
- [Til96] Jacques Tilouine. *Deformations of Galois representations and Hecke algebras*. Mehta Research Institute of Mathematics, 1996.
- [Wei94] Charles Weibel. *An introduction to homological algebra*. Vol. 38. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1994.