

Counterexamples related to the Sato–Tate conjecture

Daniel Miller

25 April 2017

Cornell University

Motivation and background

Discrepancy and Dirichlet series

Main theorem

Idea of the proof

Motivation and background

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$,

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$.

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Satake parameter: $\theta_p = \cos^{-1} \left(\frac{a_p}{2\sqrt{p}} \right)$.

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Satake parameter: $\theta_p = \cos^{-1} \left(\frac{a_p}{2\sqrt{p}} \right)$.

Sato–Tate measure: $\mathrm{ST} = \frac{2}{\pi} \sin^2 \theta \, \mathrm{d}\theta$

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Satake parameter: $\theta_p = \cos^{-1} \left(\frac{a_p}{2\sqrt{p}} \right)$.

Sato–Tate measure: $\mathrm{ST} = \frac{2}{\pi} \sin^2 \theta \, \mathrm{d}\theta$ (Haar measure on $\mathrm{SU}(2)^{\natural}$).

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Satake parameter: $\theta_p = \cos^{-1} \left(\frac{a_p}{2\sqrt{p}} \right)$.

Sato–Tate measure: $\mathrm{ST} = \frac{2}{\pi} \sin^2 \theta \, \mathrm{d}\theta$ (Haar measure on $\mathrm{SU}(2)^{\natural}$).

Theorem (Taylor et. al.)

$\{\theta_p\}$ is equidistributed with respect to ST .

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Satake parameter: $\theta_p = \cos^{-1} \left(\frac{a_p}{2\sqrt{p}} \right)$.

Sato–Tate measure: $\mathrm{ST} = \frac{2}{\pi} \sin^2 \theta \, \mathrm{d}\theta$ (Haar measure on $\mathrm{SU}(2)^{\natural}$).

Theorem (Taylor et. al.)

$\{\theta_p\}$ is equidistributed with respect to ST .

Quantify rate of convergence of $\frac{1}{\pi(N)} \sum_{p \leq N} \delta_{\theta_p}$ to ST .

Sato–Tate Conjecture

E/\mathbf{Q} non-CM elliptic curve, l prime, $\rho_l: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$.

Fact: $a_p = \mathrm{tr} \rho_l(\mathrm{fr}_p) = p + 1 - \#E(\mathbf{F}_p)$, $|a_p| \leq 2\sqrt{p}$. (Hasse)

Satake parameter: $\theta_p = \cos^{-1} \left(\frac{a_p}{2\sqrt{p}} \right)$.

Sato–Tate measure: $\mathrm{ST} = \frac{2}{\pi} \sin^2 \theta \, d\theta$ (Haar measure on $\mathrm{SU}(2)^{\natural}$).

Theorem (Taylor et. al.)

$\{\theta_p\}$ is equidistributed with respect to ST .

Quantify rate of convergence of $\frac{1}{\pi(N)} \sum_{p \leq N} \delta_{\theta_p}$ to ST .

Use discrepancy (Kolmogorov–Smirnov statistic).

Akiyama–Tanigawa Conjecture

$$D_N = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int 1_{[0, x)}(\theta) \, dST(\theta) \right|.$$

Akiyama–Tanigawa Conjecture

$$D_N = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int 1_{[0, x)}(\theta) \, dST(\theta) \right|.$$

Conjecture (Akiyama–Tanigawa)

$$D_N \ll N^{-\frac{1}{2} + \epsilon}.$$

Akiyama–Tanigawa Conjecture

$$D_N = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int 1_{[0, x)}(\theta) \, dST(\theta) \right|.$$

Conjecture (Akiyama–Tanigawa)

$$D_N \ll N^{-\frac{1}{2} + \epsilon}.$$

There is a variant of this conjecture for CM elliptic curves.

Akiyama–Tanigawa Conjecture

$$D_N = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int 1_{[0, x)}(\theta) \, dST(\theta) \right|.$$

Conjecture (Akiyama–Tanigawa)

$$D_N \ll N^{-\frac{1}{2} + \epsilon}.$$

There is a variant of this conjecture for CM elliptic curves.

Theorem (Akiyama–Tanigawa)

Akiyama–Tanigawa conjecture \Rightarrow Riemann hypothesis for E .

Akiyama–Tanigawa Conjecture

$$D_N = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int 1_{[0, x)}(\theta) \, dST(\theta) \right|.$$

Conjecture (Akiyama–Tanigawa)

$$D_N \ll N^{-\frac{1}{2} + \epsilon}.$$

There is a variant of this conjecture for CM elliptic curves.

Theorem (Akiyama–Tanigawa)

Akiyama–Tanigawa conjecture \Rightarrow Riemann hypothesis for E .

Theorem (Mazur)

Akiyama–Tanigawa conjecture \Rightarrow Riemann hypothesis for $\text{sym}^k E$

Theorem (Bucar–Kedlaya). Assume analytic continuation of $L(\mathrm{sym}^k E, s)$, GRH, and functional equation for all $k \geq 1$. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Related results

Theorem (Bucar–Kedlaya). Assume analytic continuation of $L(\mathrm{sym}^k E, s)$, GRH, and functional equation for all $k \geq 1$. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Theorem (Thorner). Same result (with explicit constants) for a modular form of arbitrary weight .

Related results

Theorem (Bucar–Kedlaya). Assume analytic continuation of $L(\mathrm{sym}^k E, s)$, GRH, and functional equation for all $k \geq 1$. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Theorem (Thorner). Same result (with explicit constants) for a modular form of arbitrary weight .

Theorem (Niederreiter). Let E/F , where F is a function field. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Related results

Theorem (Bucar–Kedlaya). Assume analytic continuation of $L(\mathrm{sym}^k E, s)$, GRH, and functional equation for all $k \geq 1$. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Theorem (Thorner). Same result (with explicit constants) for a modular form of arbitrary weight .

Theorem (Niederreiter). Let E/F , where F is a function field. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Theorem (Rosengarten). For $G = \mathrm{ST}(M)$ over a function field, $D_N \ll N^{-\alpha_G+\epsilon}$, where $\alpha \rightarrow 0$ as $\mathrm{rk} G \rightarrow \infty$.

Related results

Theorem (Bucar–Kedlaya). Assume analytic continuation of $L(\mathrm{sym}^k E, s)$, GRH, and functional equation for all $k \geq 1$. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Theorem (Thorner). Same result (with explicit constants) for a modular form of arbitrary weight .

Theorem (Niederreiter). Let E/F , where F is a function field. Then $D_N \ll N^{-\frac{1}{4}+\epsilon}$.

Theorem (Rosengarten). For $G = \mathrm{ST}(M)$ over a function field, $D_N \ll N^{-\alpha_G+\epsilon}$, where $\alpha \rightarrow 0$ as $\mathrm{rk} G \rightarrow \infty$.

Common ingredient. Erdős–Turán–Koksma inequality: from a bound on $\left| \sum_{p \leq N} \mathrm{tr} \rho(x_p) \right|$ to a bound on D_N .

A. Pande: is the Sato–Tate conjecture a Galois-theoretic result?

A. Pande: is the Sato–Tate conjecture a Galois-theoretic result?

Theorem (Pande)

Let $\epsilon > 0$. Then there exists an infinitely ramified representation $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that $\theta_p \in B_{\epsilon}(\pi/2)$ for a density one set of primes.

A. Pande: is the Sato–Tate conjecture a Galois-theoretic result?

Theorem (Pande)

Let $\epsilon > 0$. Then there exists an infinitely ramified representation $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that $\theta_p \in B_\epsilon(\pi/2)$ for a density one set of primes.

Theorem (Khare–Rajan)

Any $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ is ramified at a density zero set of primes.

A. Pande: is the Sato–Tate conjecture a Galois-theoretic result?

Theorem (Pande)

Let $\epsilon > 0$. Then there exists an infinitely ramified representation $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that $\theta_p \in B_\epsilon(\pi/2)$ for a density one set of primes.

Theorem (Khare–Rajan)

Any $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ is ramified at a density zero set of primes.

Question (Serre): can you *a priori* control the ratio $\frac{\pi_{\mathrm{ram}(\rho)}(x)}{\pi(x)}$?

A. Pande: is the Sato–Tate conjecture a Galois-theoretic result?

Theorem (Pande)

Let $\epsilon > 0$. Then there exists an infinitely ramified representation $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that $\theta_p \in B_\epsilon(\pi/2)$ for a density one set of primes.

Theorem (Khare–Rajan)

Any $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ is ramified at a density zero set of primes.

Question (Serre): can you *a priori* control the ratio $\frac{\pi_{\mathrm{ram}(\rho)}(x)}{\pi(x)}$?

Answer (Khare–Larsen–Ramakrishna): no!

Q1. Can Pande's results be strengthened to yield equidistribution?

Questions

- Q1.** Can Pande's results be strengthened to yield equidistribution?
- Q2.** If so, can the measure be specified?

Questions

- Q1.** Can Pande's results be strengthened to yield equidistribution?
- Q2.** If so, can the measure be specified?
- Q3.** Can the rate of convergence of empirical measures to the true measure be specified?
- Q4.** Can the growth of $\pi_{\text{ram}(\rho)}(x)$ be controlled?

Questions

- Q1.** Can Pande's results be strengthened to yield equidistribution?
- Q2.** If so, can the measure be specified?
- Q3.** Can the rate of convergence of empirical measures to the true measure be specified?
- Q4.** Can the growth of $\pi_{\text{ram}(\rho)}(x)$ be controlled?
- Q5.** Can anything be said about the L -functions associated with ρ ?

Questions

Q1. Can Pande's results be strengthened to yield equidistribution?

Q2. If so, can the measure be specified?

Q3. Can the rate of convergence of empirical measures to the true measure be specified?

Q4. Can the growth of $\pi_{\text{ram}(\rho)}(x)$ be controlled?

Q5. Can anything be said about the L -functions associated with ρ ?

Answer: Yes! to Q1–Q5.

Discrepancy and Dirichlet series

Definition

Let $\{\theta_p\}$ be a sequence in $[0, \pi]$, μ a measure on $[0, \pi]$. The *discrepancy* is

$$D_N(\{\theta_p\}, \mu) = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int_0^x d\mu \right|.$$

Definition

Let $\{\theta_p\}$ be a sequence in $[0, \pi]$, μ a measure on $[0, \pi]$. The *discrepancy* is

$$D_N(\{\theta_p\}, \mu) = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int_0^x d\mu \right|.$$

Fact: $\{\theta_p\}$ are μ -equidistributed if and only if $D_N \rightarrow 0$.

Definition

Let $\{\theta_p\}$ be a sequence in $[0, \pi]$, μ a measure on $[0, \pi]$. The *discrepancy* is

$$D_N(\{\theta_p\}, \mu) = \sup_{x \in [0, \pi]} \left| \frac{1}{\pi(N)} \sum_{p \leq N} 1_{[0, x)}(\theta_p) - \int_0^x d\mu \right|.$$

Fact: $\{\theta_p\}$ are μ -equidistributed if and only if $D_N \rightarrow 0$.

Fact: $\frac{\log N}{N} \ll D_N$. The *van der Corput sequence* achieves this.

Definition

For $k \geq 1$,

$$L(\mathrm{sym}^k \rho, s) = \prod_p \det \left(1 - \mathrm{sym}^k \begin{pmatrix} e^{i\theta_p} & 0 \\ 0 & e^{-i\theta_p} \end{pmatrix} p^{-s} \right)^{-1}$$

Definition

For $k \geq 1$,

$$L(\text{sym}^k \rho, s) = \prod_p \det \left(1 - \text{sym}^k \begin{pmatrix} e^{i\theta_p} & 0 \\ 0 & e^{-i\theta_p} \end{pmatrix} p^{-s} \right)^{-1}$$

Definition

For $f: [0, \pi] \rightarrow \mathbf{C}$ of bounded variation with $\mu(f) = 0$,

$$L_f(s) = \prod_p (1 - f(\theta_p) p^{-s})^{-1}$$

Definition

For $k \geq 1$,

$$L(\text{sym}^k \rho, s) = \prod_p \det \left(1 - \text{sym}^k \begin{pmatrix} e^{i\theta_p} & 0 \\ 0 & e^{-i\theta_p} \end{pmatrix} p^{-s} \right)^{-1}$$

Definition

For $f: [0, \pi] \rightarrow \mathbf{C}$ of bounded variation with $\mu(f) = 0$,

$$L_f(s) = \prod_p (1 - f(\theta_p) p^{-s})^{-1}$$

Example (Ramakrishna): $L_{\text{sgn}}(s) = \prod_p (1 - \text{sgn}(a_p) p^{-s})^{-1}$.

Theorem

If $\left| \sum_{p \leq N} f(\theta_p) \right| \ll N^{\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Dirichlet series—basic facts

Theorem

If $\left| \sum_{p \leq N} f(\theta_p) \right| \ll N^{\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Corollary

If $D_N \ll N^{-\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Dirichlet series—basic facts

Theorem

If $\left| \sum_{p \leq N} f(\theta_p) \right| \ll N^{\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Corollary

If $D_N \ll N^{-\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Definition

$$U_k(\theta) = \frac{\sin((k+1)\theta)}{\sin \theta} = \text{tr sym}^k \begin{pmatrix} e^{i\theta_p} & 0 \\ 0 & e^{-i\theta_p} \end{pmatrix}.$$

Dirichlet series—basic facts

Theorem

If $\left| \sum_{p \leq N} f(\theta_p) \right| \ll N^{\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Corollary

If $D_N \ll N^{-\alpha+\epsilon}$, then $L_f(s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Definition

$$U_k(\theta) = \frac{\sin((k+1)\theta)}{\sin \theta} = \text{tr sym}^k \begin{pmatrix} e^{i\theta_p} & 0 \\ 0 & e^{-i\theta_p} \end{pmatrix}.$$

Theorem

If $\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\alpha+\epsilon}$, then $L(\text{sym}^k \rho, s)$ admits a nonvanishing analytic continuation to $\Re > \alpha$.

Main theorem

Ingredients

1. Fix a rational prime $l \geq 7$.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.
3. Fix a function $h: \mathbf{R}^+ \rightarrow \mathbf{R}_{\geq 1}$ which increases slowly to infinity.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.
3. Fix a function $h: \mathbf{R}^+ \rightarrow \mathbf{R}_{\geq 1}$ which increases slowly to infinity.
We will have $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.
3. Fix a function $h: \mathbf{R}^+ \rightarrow \mathbf{R}_{\geq 1}$ which increases slowly to infinity. We will have $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$.
4. Fix an absolutely continuous probability measure μ on $[0, \pi]$, with bounded probability density function.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.
3. Fix a function $h: \mathbf{R}^+ \rightarrow \mathbf{R}_{\geq 1}$ which increases slowly to infinity. We will have $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$.
4. Fix an absolutely continuous probability measure μ on $[0, \pi]$, with bounded probability density function. The angles $\{\theta_p\}$ will be μ -equidistributed.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.
3. Fix a function $h: \mathbf{R}^+ \rightarrow \mathbf{R}_{\geq 1}$ which increases slowly to infinity. We will have $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$.
4. Fix an absolutely continuous probability measure μ on $[0, \pi]$, with bounded probability density function. The angles $\{\theta_p\}$ will be μ -equidistributed.
5. Fix $\alpha \in (0, \frac{1}{3})$.

Ingredients

1. Fix a rational prime $l \geq 7$.
2. Fix an odd, absolutely irreducible, weight 2 representation $\bar{\rho}: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{F}_l)$. ρ will be a lift of $\bar{\rho}$.
3. Fix a function $h: \mathbf{R}^+ \rightarrow \mathbf{R}_{\geq 1}$ which increases slowly to infinity. We will have $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$.
4. Fix an absolutely continuous probability measure μ on $[0, \pi]$, with bounded probability density function. The angles $\{\theta_p\}$ will be μ -equidistributed.
5. Fix $\alpha \in (0, \frac{1}{3})$. The discrepancy D_N will decay like $\pi(N)^{-\alpha}$.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$,

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$,

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$, $A^{-1}(x)$)

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$, $A^{-1}(x)$)
3. For each unramified p , $a_p = \mathrm{tr} \rho(\mathrm{fr}_p) \in \mathbf{Z}$ and satisfies the Hasse bound.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$, $A^{-1}(x)$)
3. For each unramified p , $a_p = \mathrm{tr} \rho(\mathrm{fr}_p) \in \mathbf{Z}$ and satisfies the Hasse bound.
4. $D_N(\{\theta_p\}, \mu) = \Theta(\pi(x)^{-\alpha})$.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$, $A^{-1}(x)$)
3. For each unramified p , $a_p = \mathrm{tr} \rho(\mathrm{fr}_p) \in \mathbf{Z}$ and satisfies the Hasse bound.
4. $D_N(\{\theta_p\}, \mu) = \Theta(\pi(x)^{-\alpha})$. (Yes to Q1–Q3.)

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$, $A^{-1}(x)$)
3. For each unramified p , $a_p = \mathrm{tr} \rho(\mathrm{fr}_p) \in \mathbf{Z}$ and satisfies the Hasse bound.
4. $D_N(\{\theta_p\}, \mu) = \Theta(\pi(x)^{-\alpha})$. (Yes to Q1–Q3.)
5. If $(\theta \mapsto \pi - \theta)_* \mu = \mu$, then for each odd k , $L(\mathrm{sym}^k \rho, s)$ satisfies the Riemann hypothesis.

Theorem

Let l , $\bar{\rho}$, h , μ , and α be as above. Then there exists $\rho: G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{Z}_l)$ such that

1. $\rho \equiv \bar{\rho} \pmod{l}$.
2. $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$. (Yes to Q4. $\log x$, $\log^{10^{10}} x$, $A^{-1}(x)$)
3. For each unramified p , $a_p = \mathrm{tr} \rho(\mathrm{fr}_p) \in \mathbf{Z}$ and satisfies the Hasse bound.
4. $D_N(\{\theta_p\}, \mu) = \Theta(\pi(x)^{-\alpha})$. (Yes to Q1–Q3.)
5. If $(\theta \mapsto \pi - \theta)_* \mu = \mu$, then for each odd k , $L(\mathrm{sym}^k \rho, s)$ satisfies the Riemann hypothesis. (Yes to Q5.)

Idea of the proof

Theorem

If $\alpha \in (0, \frac{1}{3})$, there exists a sequence (x_2, x_3, x_5, \dots) in $[-1, 1]$ such that $|D_N - \pi(N)^{-\alpha}| \ll \pi(N)^{-1}$.

Theorem

If $\alpha \in (0, \frac{1}{3})$, there exists a sequence (x_2, x_3, x_5, \dots) in $[-1, 1]$ such that $|D_N - \pi(N)^{-\alpha}| \ll \pi(N)^{-1}$.

(Can have $x_p = \frac{a_p}{2\sqrt{p}}$ for $a_p \in \mathbf{Z}$ satisfying the Hasse bound.)

Theorem

If $\alpha \in (0, \frac{1}{3})$, there exists a sequence (x_2, x_3, x_5, \dots) in $[-1, 1]$ such that $|D_N - \pi(N)^{-\alpha}| \ll \pi(N)^{-1}$.

(Can have $x_p = \frac{a_p}{2\sqrt{p}}$ for $a_p \in \mathbf{Z}$ satisfying the Hasse bound.)

Fact: Discrepancy is invariant under pushforward by \cos and \cos^{-1} .

Theorem

If $\alpha \in (0, \frac{1}{3})$, there exists a sequence (x_2, x_3, x_5, \dots) in $[-1, 1]$ such that $|D_N - \pi(N)^{-\alpha}| \ll \pi(N)^{-1}$.

(Can have $x_p = \frac{a_p}{2\sqrt{p}}$ for $a_p \in \mathbf{Z}$ satisfying the Hasse bound.)

Fact: Discrepancy is invariant under pushforward by \cos and \cos^{-1} .

Idea: Construct ρ so that $\frac{a_p}{2\sqrt{p}} \approx x_p$.

Prescribing discrepancy decay

Theorem

If $\alpha \in (0, \frac{1}{3})$, there exists a sequence (x_2, x_3, x_5, \dots) in $[-1, 1]$ such that $|D_N - \pi(N)^{-\alpha}| \ll \pi(N)^{-1}$.

(Can have $x_p = \frac{a_p}{2\sqrt{p}}$ for $a_p \in \mathbf{Z}$ satisfying the Hasse bound.)

Fact: Discrepancy is invariant under pushforward by \cos and \cos^{-1} .

Idea: Construct ρ so that $\frac{a_p}{2\sqrt{p}} \approx x_p$.

Fact: If $(x_p^{(1)})$ is a sequence with $|x_p - x_p^{(1)}| \ll p^{-1/2+\epsilon}$, then $D_N^{(1)} = \Theta(\pi(N)^{-\alpha})$.

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

For all n , require $\det \rho_n \equiv \kappa \pmod{l^n}$

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

For all n , require $\det \rho_n \equiv \kappa \pmod{l^n}$ (l -adic cyclotomic character)

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

For all n , require $\det \rho_n \equiv \kappa \pmod{l^n}$ (l -adic cyclotomic character)

Passage from ρ_n to ρ_{n+1} is governed by $H^i(G_{\mathbf{Q}, R_n}, \mathrm{Ad}^0 \bar{\rho})$, $i = 1, 2$

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

For all n , require $\det \rho_n \equiv \kappa \pmod{l^n}$ (l -adic cyclotomic character)

Passage from ρ_n to ρ_{n+1} is governed by $H^i(G_{\mathbf{Q}, R_n}, \mathrm{Ad}^0 \bar{\rho})$, $i = 1, 2$

Theorem (K–L–R). Fix a finite set U of primes. Then there exists a finite set N of primes such that

$$H^1(G_{\mathbf{Q}, R \cup N}, \mathrm{Ad}^0 \bar{\rho}) \xrightarrow{\sim} \prod_{p \in R} H^1(G_{\mathbf{Q}_p}, \mathrm{Ad}^0 \bar{\rho}) \times \prod_{p \in U} H^1_{\mathrm{nr}}(G_{\mathbf{Q}_p}, \mathrm{Ad}^0 \bar{\rho})$$

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

For all n , require $\det \rho_n \equiv \kappa \pmod{l^n}$ (l -adic cyclotomic character)

Passage from ρ_n to ρ_{n+1} is governed by $H^i(G_{\mathbf{Q}, R_n}, \mathrm{Ad}^0 \bar{\rho})$, $i = 1, 2$

Theorem (K–L–R). Fix a finite set U of primes. Then there exists a finite set N of primes such that

$$H^1(G_{\mathbf{Q}, R \cup N}, \mathrm{Ad}^0 \bar{\rho}) \xrightarrow{\sim} \prod_{p \in R} H^1(G_{\mathbf{Q}_p}, \mathrm{Ad}^0 \bar{\rho}) \times \prod_{p \in U} H^1_{\mathrm{nr}}(G_{\mathbf{Q}_p}, \mathrm{Ad}^0 \bar{\rho})$$

Corollary: Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$, can choose $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p)$ for all p in a finite set.

Lifting Galois representations

Construct ρ as $\varprojlim \rho_n$, where $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$.

For all n , require $\det \rho_n \equiv \kappa \pmod{l^n}$ (l -adic cyclotomic character)

Passage from ρ_n to ρ_{n+1} is governed by $H^i(G_{\mathbf{Q}, R_n}, \mathrm{Ad}^0 \bar{\rho})$, $i = 1, 2$

Theorem (K–L–R). Fix a finite set U of primes. Then there exists a finite set N of primes such that

$$H^1(G_{\mathbf{Q}, R \cup N}, \mathrm{Ad}^0 \bar{\rho}) \xrightarrow{\sim} \prod_{p \in R} H^1(G_{\mathbf{Q}_p}, \mathrm{Ad}^0 \bar{\rho}) \times \prod_{p \in U} H^1_{\mathrm{nr}}(G_{\mathbf{Q}_p}, \mathrm{Ad}^0 \bar{\rho})$$

Corollary: Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$, can choose $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p)$ for all p in a finite set. (Finitely many more ramified primes.)

Controlling ramified primes

Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/I^n)$ and choices of $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p)$ $(\mathrm{mod} \ I^{n+1})$, need to add finite set N to R_n .

Controlling ramified primes

Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/I^n)$ and choices of $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p) \pmod{I^{n+1}}$, need to add finite set N to R_n .

Each $p \in N$ is chosen from a positive-density set of primes.

Controlling ramified primes

Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/I^n)$ and choices of $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p) \pmod{I^{n+1}}$, need to add finite set N to R_n .

Each $p \in N$ is chosen from a positive-density set of primes.

So p can be arbitrarily large!

Controlling ramified primes

Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$ and choices of $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p) \pmod{l^{n+1}}$, need to add finite set N to R_n .

Each $p \in N$ is chosen from a positive-density set of primes.

So p can be arbitrarily large!

If $\pi_R(x) \leq h(x) \ (\forall x)$, can force this for $\pi_{R \cup N}(x)$.

Controlling ramified primes

Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$ and choices of $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p) \pmod{l^{n+1}}$, need to add finite set N to R_n .

Each $p \in N$ is chosen from a positive-density set of primes.

So p can be arbitrarily large!

If $\pi_R(x) \leq h(x) \ (\forall x)$, can force this for $\pi_{R \cup N}(x)$.

$\pi_{\mathrm{ram}(\bar{\rho})}(x) \leq h(x)$ may not hold—scale h to make this true!

Controlling ramified primes

Given $\rho_n: G_{\mathbf{Q}, R_n} \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$ and choices of $\mathrm{tr} \rho_{n+1}(\mathrm{fr}_p) \pmod{l^{n+1}}$, need to add finite set N to R_n .

Each $p \in N$ is chosen from a positive-density set of primes.

So p can be arbitrarily large!

If $\pi_R(x) \leq h(x) \ (\forall x)$, can force this for $\pi_{R \cup N}(x)$.

$\pi_{\mathrm{ram}(\bar{\rho})}(x) \leq h(x)$ may not hold—scale h to make this true!

Fact: constant in $\pi_{\mathrm{ram}(\rho)}(x) \ll h(x)$ only depends on $\bar{\rho}$.

Lifting Galois representations—first stage

Lift from \mathbf{Z}/l to \mathbf{Z}/l^2 .

Lifting Galois representations—first stage

Lift from \mathbf{Z}/l to \mathbf{Z}/l^2 .

Fix a large finite set U_1 of primes.

Lifting Galois representations—first stage

Lift from \mathbf{Z}/l to \mathbf{Z}/l^2 .

Fix a large finite set U_1 of primes.

For $p \in U_1$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_1(\text{fr}_p) \pmod{l}$.

Lifting Galois representations—first stage

Lift from \mathbf{Z}/l to \mathbf{Z}/l^2 .

Fix a **large** finite set U_1 of primes.

For $p \in U_1$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_1(\text{fr}_p) \pmod{l}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l}{2\sqrt{p}}$.

Lifting Galois representations—first stage

Lift from \mathbf{Z}/l to \mathbf{Z}/l^2 .

Fix a **large** finite set U_1 of primes.

For $p \in U_1$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_1(\text{fr}_p) \pmod{l}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l}{2\sqrt{p}}$.

For $N \leq \max U_1$, $D_N(\{\theta_p\}, \mu) = \Theta(\pi(N)^{-\alpha})$.

Lifting Galois representations—first stage

Lift from \mathbf{Z}/l to \mathbf{Z}/l^2 .

Fix a **large** finite set U_1 of primes.

For $p \in U_1$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_1(\text{fr}_p) \pmod{l}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l}{2\sqrt{p}}$.

For $N \leq \max U_1$, $D_N(\{\theta_p\}, \mu) = \Theta(\pi(N)^{-\alpha})$.

Make U_1 so large that for $p > \max U_1$, $l^2 < \log p$.

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$.

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Fix a really huge $U_{n+1} \supset U_n$.

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Fix a **really huge** $U_{n+1} \supset U_n$.

For $p \in U_{n+1} \setminus U_n$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_n(\text{fr}_p) \pmod{l^n}$.

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Fix a **really huge** $U_{n+1} \supset U_n$.

For $p \in U_{n+1} \setminus U_n$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_n(\text{fr}_p) \pmod{l^n}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l^n}{2\sqrt{p}}$.

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Fix a **really huge** $U_{n+1} \supset U_n$.

For $p \in U_{n+1} \setminus U_n$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_n(\text{fr}_p) \pmod{l^n}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l^n}{2\sqrt{p}}$. ($l^n \ll \log p$).

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Fix a **really huge** $U_{n+1} \supset U_n$.

For $p \in U_{n+1} \setminus U_n$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_n(\text{fr}_p) \pmod{l^n}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l^n}{2\sqrt{p}}$. ($l^n \ll \log p$).

For $N \leq \max U_{n+1}$, $D_N(\{\theta_p\}, \mu) = \Theta(\pi(N)^{-\alpha})$.

Lifting Galois representations—inductive step

Lift from \mathbf{Z}/l^n to \mathbf{Z}/l^{n+1} .

Have already chosen a_p for $p \in U_n$. (1–5 hold)

Fix a **really huge** $U_{n+1} \supset U_n$.

For $p \in U_{n+1} \setminus U_n$, can choose $a_p \in \mathbf{Z}$ subject only to $|a_p| \leq 2\sqrt{p}$ and $a_p \equiv \text{tr } \rho_n(\text{fr}_p) \pmod{l^n}$.

We can ensure $\left| \frac{a_p}{2\sqrt{p}} - x_p \right| \leq \frac{l^n}{2\sqrt{p}}$. ($l^n \ll \log p$).

For $N \leq \max U_{n+1}$, $D_N(\{\theta_p\}, \mu) = \Theta(\pi(N)^{-\alpha})$.

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

Riemann hypothesis

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

When k is odd, $U_k(\pi - \theta) = -U_k(\theta)$.

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

When k is odd, $U_k(\pi - \theta) = -U_k(\theta)$.

Enumerate the primes $p_1 = 2, q_1 = 3, p_2 = 5, q_2 = 7, \dots$

Riemann hypothesis

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

When k is odd, $U_k(\pi - \theta) = -U_k(\theta)$.

Enumerate the primes $p_1 = 2, q_1 = 3, p_2 = 5, q_2 = 7, \dots$

If $\theta_{q_i} \approx \pi - \theta_{p_i}$, then $U_k(\theta_{q_i}) \approx -U_k(\theta_{p_i})$

Riemann hypothesis

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

When k is odd, $U_k(\pi - \theta) = -U_k(\theta)$.

Enumerate the primes $p_1 = 2, q_1 = 3, p_2 = 5, q_2 = 7, \dots$

If $\theta_{q_i} \approx \pi - \theta_{p_i}$, then $U_k(\theta_{q_i}) \approx -U_k(\theta_{p_i})$ (within $p_i^{-1/2}$).

Riemann hypothesis

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

When k is odd, $U_k(\pi - \theta) = -U_k(\theta)$.

Enumerate the primes $p_1 = 2, q_1 = 3, p_2 = 5, q_2 = 7, \dots$

If $\theta_{q_i} \approx \pi - \theta_{p_i}$, then $U_k(\theta_{q_i}) \approx -U_k(\theta_{p_i})$ (within $p_i^{-1/2}$).

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| = \left| \sum_{p_i, q_i \leq N} (U_k(\theta_{p_i}) + U_k(\theta_{q_i})) \right| \ll \sum_{n \leq N} n^{-1/2}$$

Riemann hypothesis

How can we make $L(\text{sym}^k \rho, s)$, k odd, satisfy the Riemann hypothesis?

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| \ll N^{\frac{1}{2} + \epsilon} \text{ implies RH for } L(\text{sym}^k \rho, s).$$

When k is odd, $U_k(\pi - \theta) = -U_k(\theta)$.

Enumerate the primes $p_1 = 2, q_1 = 3, p_2 = 5, q_2 = 7, \dots$

If $\theta_{q_i} \approx \pi - \theta_{p_i}$, then $U_k(\theta_{q_i}) \approx -U_k(\theta_{p_i})$ (within $p_i^{-1/2}$).

$$\left| \sum_{p \leq N} U_k(\theta_p) \right| = \left| \sum_{p_i, q_i \leq N} (U_k(\theta_{p_i}) + U_k(\theta_{q_i})) \right| \ll \sum_{n \leq N} n^{-1/2} \\ \ll N^{\frac{1}{2}}.$$

Consequences

If $f \in C([0, \pi])$, $f \circ \cos^{-1}: [-1, 1] \rightarrow \mathbf{C}$ is Lipschitz, and $f(\pi - \theta) = -f(\theta)$, then $L_f(\rho, s)$ has a nonvanishing analytic continuation to $\Re > \frac{1}{2}$ (Riemann hypothesis).

Consequences

If $f \in C([0, \pi])$, $f \circ \cos^{-1}: [-1, 1] \rightarrow \mathbf{C}$ is Lipschitz, and $f(\pi - \theta) = -f(\theta)$, then $L_f(\rho, s)$ has a nonvanishing analytic continuation to $\Re > \frac{1}{2}$ (Riemann hypothesis).

For μ any “bump measure,” there exists ρ with $\{\theta_\rho\}$ μ -equidistributed.

Consequences

If $f \in C([0, \pi])$, $f \circ \cos^{-1}: [-1, 1] \rightarrow \mathbf{C}$ is Lipschitz, and $f(\pi - \theta) = -f(\theta)$, then $L_f(\rho, s)$ has a nonvanishing analytic continuation to $\Re > \frac{1}{2}$ (Riemann hypothesis).

For μ any “bump measure,” there exists ρ with $\{\theta_\rho\}$ μ -equidistributed.

Can get equidistribution with respect to μ with non-continuous probability distribution functions.

Questions?