

Counterexamples related to the Sato–Tate conjecture for CM abelian varieties*

Daniel Miller

31 March 2017

1 Introduction and motivation

Let K/\mathbf{Q} be a finite Galois extension, A/K a g -dimensional abelian variety. Assume that A has CM defined over K ; then there is a CM field F together with an isomorphism $F \simeq \text{End}_K(A)_{\mathbf{Q}}$. Let $\mathfrak{a} = \text{Lie}(A)$. Then the determinant of the action of K on \mathfrak{a} (viewed as an F -vector space) gives a map $\det_{\mathfrak{a}}: \text{R}_{K/\mathbf{Q}} \mathbf{G}_m \rightarrow \text{R}_{F/\mathbf{Q}} \mathbf{G}_m$. Put $G_A = \text{im}(\det_{\mathfrak{a}})$ and $G_A^1 = \text{im}(\det_{\mathfrak{a}})^{\text{N}_{F/\mathbf{Q}}=1}$. The group G_A is the motivic Galois group of A , and $\text{ST}(A)$, the Sato–Tate group of A , is a maximal compact subgroup of $G_A^1(\mathbf{C})$. It is obvious that the l -adic Galois representation associated with A , $\rho_l: G_{\mathbf{Q}} \rightarrow \text{GL}_{2g}(\mathbf{Q}_l)$, has image in $(\text{R}_{F/\mathbf{Q}} \mathbf{G}_m)(\mathbf{Q}_l)$; it actually is a map $\rho_l: G_{\mathbf{Q}} \rightarrow G_A(\mathbf{Q}_l)$.

Unitary representations of $\text{ST}(A)$ are just characters, and basic representation theory tells us that all such representations are induced by an (algebraic) character of G_A defined over $\overline{\mathbf{Q}}$. For $r \in X^*(G_A)$, there is an L -function $L(r_*\rho_l, s)$ coming from the composite Galois representation $r \circ \rho_l: G_{\mathbf{Q}} \rightarrow G_A(\mathbf{Q}_l) \rightarrow \overline{\mathbf{Q}_l}^{\times}$. The Sato–Tate conjecture for A says that all $L(r_*\rho_l, s)$ have non-vanishing analytic continuation past $\Re = 1$, and the Generalized Riemann hypothesis for A says that all $L(r_*\rho_l, s)$ satisfy the Riemann hypothesis.

Choose an isomorphism $(\mathbf{R}/\mathbf{Z})^d \simeq \text{ST}(A)$, and put

$$D_x(A) = \sup_{t \in [0,1]^d} \left| \frac{1}{\pi_K(x)} \sum_{\text{N}(\mathfrak{p}) \leq x} 1_{[0,t)}(\theta_{\mathfrak{p}}) - \int 1_{[0,t)} \right|.$$

Akiyama and Tanigawa conjectured that for non-CM elliptic curves, $D_x(E) \ll x^{-\frac{1}{2}+\epsilon}$. We call the “Akiyama–Tanigawa conjecture” for A the discrepancy de-

*Notes for a talk given in Cornell’s Number Theory Seminar.

cays like $D_x(A) \ll x^{-\frac{1}{2}+\epsilon}$. Via the Koksma–Hlawka inequality, the Akiyama–Tanigawa conjecture implies that for all bounded-variation functions f on $\text{ST}(A)$, the estimate

$$\left| \sum_{N(\mathfrak{p}) \leq x} f(\theta_{\mathfrak{p}}) \right| \ll \text{Var}(f) x^{\frac{1}{2}+\epsilon}.$$

For $r \in X^*(G_A)$, this estimate implies the Riemann hypothesis for the L -function $L(r_*\rho_l, s)$.

Analogy with Artin L -functions (go into detail here!) seems to suggest that if all $L(r_*\rho_l, s)$ satisfy the Riemann Hypothesis, then the Akiyama–Tanigawa conjecture for A holds. We’ll show: this converse is false, in a limited sense.

2 Diophantine approximation

d

3 Fake Satake parameters

d