JAMA | Original Investigation

Frequency and Clinical Outcomes Associated With Tau Positron Emission Tomography Positivity

Alexis Moscoso, PhD; Fiona Heeman, PhD; Sheelakumari Raghavan, PhD; Alejandro Costoya-Sánchez, PhD; Martijn van Essen, MD, PhD; Ismini Mainta, MD; Valle Camacho, MD, PhD; Omar Rodríguez-Fonseca, MD, PhD; Jesús Silva-Rodríguez, PhD; Andrés Perissinotti, MD, PhD; Yuna Gu, MSc; Jihwan Yun, MSc; Debora Peretti, PhD; Federica Ribaldi, PhD; Emma M. Coomans, PhD; Wagner S. Brum; Michel J. Grothe, PhD; Pablo Aguiar, PhD; Gérard N. Bischof, PhD; Alexander Drzezga, MD; Sang Won Seo, MD, PhD; Sylvia Villeneuve, PhD; Maura Malpetti, PhD; John T. O'Brien, DM; James B. Rowe, MD, PhD; Elsmarieke M. van de Giessen, MD, PhD; Rik Ossenkoppele, PhD; William J. Jagust, MD; Ruben Smith, MD, PhD; Oskar Hansson, MD, PhD; Giovanni B. Frisoni, MD; Valentina Garibotto, MD; David N. Soleimani-Meigooni, MD; Maria Carrillo, PhD; Bradford C. Dickerson, MD; Renaud La Joie, PhD; Gil D. Rabinovici, MD; Liana G. Apostolova, MD; Pamela J. LaMontagne, PhD; Michael J. Pontecorvo, PhD; Keith A. Johnson, MD; Reisa A. Sperling, MD; Michael W. Weiner, MD; Ronald C. Petersen, MD, PhD; Clifford R. Jack Jr, MD; Prashanthi Vemuri, PhD; Michael Schöll, PhD; for the PREVENT-AD Research Group, the Harvard Aging Brain Study, the LEADS Consortium, and the Alzheimer's Disease Neuroimaging Initiative

IMPORTANCE Tau positron emission tomography (PET) allows in vivo detection of neurofibrillary tangles, a core neuropathologic feature of Alzheimer disease (AD).

OBJECTIVE To provide estimates of the frequency of tau PET positivity and its associated risk of clinical outcomes.

DESIGN, SETTING, AND PARTICIPANTS Longitudinal study using data pooled from 21 cohorts, comprising a convenience sample of 6514 participants from 13 countries, collected between January 2013 and June 2024. Cognitively unimpaired individuals and patients with a clinical diagnosis of mild cognitive impairment (MCI), AD dementia, or other neurodegenerative disorders were included.

EXPOSURES Tau PET with flortaucipir F 18, amyloid- β (A β) PET, and clinical examinations. Tau PET scans were visually rated as positive according to a US Food and Drug Administrationand European Medicines Agency-approved method, designed to indicate the presence of advanced neurofibrillary tangle pathology (Braak stages V-VI).

MAIN OUTCOMES AND MEASURES Frequency of tau PET positivity and absolute risk of clinical progression (eg., progression to MCI or dementia).

RESULTS Among the 6514 participants (mean age, 69.5 years; 50.5% female), median follow-up time ranged from 1.5 to 4.0 years. Of 3487 cognitively unimpaired participants, 349 (9.8%) were tau PET positive; the estimated frequency of tau PET positivity was less than 1% in those aged younger than 50 years, and increased from 3% (95% CI, 2%-4%) at 60 years to 19% (95% CI, 16%-24%) at 90 years. Tau PET positivity frequency estimates increased across MCI and AD dementia clinical diagnoses (43% [95% CI, 41%-46%] and 79% [95% CI, 77%-82%] at 75 years, respectively). Most tau PET-positive individuals (92%) were also A β PET positive. Cognitively unimpaired participants who were positive for both A β PET and tau PET had a higher absolute risk of progression to MCI or dementia over the following '5 years (57% [95% CI, 45%-71%]) compared with both A β PET-positive/tau PET-negative (17% [95% CI, 13%-22%]) and A β PET-negative/tau PET-negative (6% [95% CI, 5%-8%]) individuals. Among participants with MCI at the time of the tau PET scan, an A β PET-positive/tau PET-positive profile was associated with a 5-year absolute risk of progression to dementia of 70% (95% CI, 59%-81%).

CONCLUSIONS AND RELEVANCE In a large convenience sample, a positive tau PET scan occurred at a nonnegligible rate among cognitively unimpaired individuals, and the combination of A β PET positivity and tau PET positivity was associated with a high risk of clinical progression in both preclinical and symptomatic stages of AD. These findings underscore the potential of tau PET as a biomarker for staging AD pathology.

JAMA. doi:10.1001/jama.2025.7817 Published online June 16, 2025. Supplemental content

Author Affiliations: Author affiliations are listed at the end of this article.

Group Information: The PREVENT-AD Research Group, Harvard Aging Brain Study, LEADS Consortium, and Alzheimer's Disease Neuroimaging Initiative members are listed in Supplement 2.

Corresponding Authors: Michael Schöll, PhD (michael.scholl@neuro.gu.se), and Alexis Moscoso, PhD (alexis.moscoso@gu.se), Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Wallinsgatan 6, 431 41 Mölndal, Sweden.

he development of imaging and fluid biomarkers of amyloid- β (A β) plaques, ^{1,2} a core neuropathologic feature of Alzheimer disease (AD), ³ has had a major impact on AD research and drug development. ⁴ Positron emission tomography (PET) imaging of A β has also demonstrated clinical utility in patients with mild cognitive impairment (MCI) or dementia of uncertain etiology. ⁵ However, the precise clinical consequences of A β positivity in earlier stages of the disease remain uncertain because A β positivity can occur in asymptomatic older individuals ⁶ who remain symptom free over their lifetime. ⁷ This lack of strong correspondence between A β positivity and relevant outcomes has led some experts in the AD field to question the clinical utility of biological definitions of AD independent of symptoms, ⁸ although this opinion is not universally accepted. ⁹

Contrary to Aβ, autopsy studies have found that the presence of widespread neocortical tau neurofibrillary tangles, another neuropathologic hallmark of AD,3 was strongly associated with neurodegeneration and clinical symptoms. 10,11 This close relationship underscores the importance of detecting tau pathology in vivo, complementing the information provided by Aβ biomarkers. To date, the only biomarker of neurofibrillary tangles approved for clinical use by the US Food and Drug Administration (FDA) and the European Medicines Agency (EMA) is tau PET imaging with flortaucipir F 18 (formerly known as [18F]T807 or [18F]AV1451).12 When assessed using an approved visual interpretation method, a positive flortaucipir F 18 PET scan reflects the presence of advanced neurofibrillary tangle pathology, corresponding to Braak stages V and VI at autopsy. 13 Braak staging 14 is a neuropathological framework that describes the spatial progression of tau neurofibrillary tangle pathology in AD. In this system, stages I and II involve the transentorhinal region, III and IV extend to limbic areas, and V and VI represent the most advanced stages with widespread neocortical involvement. Clinically, Braak stages V and VI represent the neuropathological stage most closely associated with significant cognitive impairment and dementia. 3,10,11,15

The investigation of tau PET imaging using an FDA- and EMA-approved method to establish positivity could provide valuable insights directly relevant for its use in clinical settings. However, previous studies on tau PET have relied on the use of varying, nonclinically applicable definitions of tau PET positivity, leading to conflicting results on crucial metrics that inform its potential utility. These include inconsistent frequency estimates of tau PET positivity (eg, ranging from $2\%-3\%^{16,17}$ to $20\%-50\%^{18-20}$ in Aβ-negative cognitively unimpaired individuals), as well as highly variable estimates of the risk of clinical progression associated with tau PET positivity, with some studies suggesting limited relevance¹⁸ while others reported a high risk. 16,17 To clarify the potential of tau PET in clinical settings, it is essential to investigate the frequency of and outcomes associated with tau PET positivity when established using clinically applicable methods.

The purpose of this study was to contribute to a better understanding of the potential of tau PET by providing estimates of 2 key metrics: the frequency of tau PET positivity and its associated risk of clinical progression. For this, data were

Key Points

Question What is the frequency of tau positron emission tomography (PET) positivity, defined using a clinically applicable visual interpretation method, and its associated risk of cognitive decline?

Findings In this longitudinal cohort study that included a convenience sample of 6514 participants from 21 cohorts, 9.8% of cognitively unimpaired individuals had positive tau PET scans, and the frequency of positivity increased with age and across increasingly symptomatic stages of Alzheimer disease. Participants who were positive for both amyloid- β (Aβ) PET and tau PET had a higher risk of clinically relevant cognitive decline over the following 5 years compared with both Aβ PET–positive/tau PET–negative and Aβ PET–negative/tau PET–negative individuals.

Meaning These findings underscore the potential role of tau PET as a biomarker for staging Alzheimer disease neuropathologic changes across the spectrum of the disease.

analyzed from a large convenience sample of participants who underwent tau PET imaging with an approved radiotracer, flortaucipir F 18, in whom tau PET positivity was determined using a clinically applicable visual interpretation method.

Methods

Local institutional review boards for human research at each participating institution approved use of data for this study. All participants provided written informed consent. The study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

Participants

A convenience sample of 6514 participants pooled from 21 different cohorts was included (eMethods 1.1 and eTable 1 in Supplement 1). All participants underwent a tau PET scan with flortaucipir F 18 between January 2013 and June 2024 and had a concurrent clinical evaluation (Table 1). Participants were either cognitively unimpaired (defined as not MCI, dementia, or any other major neurological disorder; see eMethods 1.1 in Supplement 1) or met established clinical criteria for a syndromic diagnosis of MCI,²¹ AD dementia (including posterior and logopenic variants),22 dementia with Lewy bodies,23 behavioral variant frontotemporal dementia,24 semantic or nonfluent variant primary progressive aphasia,25 Parkinson disease²⁶ with cognitive impairment, progressive supranuclear palsy,²⁷ or corticobasal syndrome.²⁸ This diagnostic categorization was based solely on the aforementioned clinical criteria, independent of biomarker results or dementia severity rating instruments.

In addition, to assess whether the frequency of visual tau PET positivity (a proxy of Braak stages V-VI) was consistent with the frequency of autopsy-confirmed Braak stages V and VI across different ages, we conducted an indirect comparison using an independent sample of participants from the National Alzheimer's Coordinating Center dataset. These participants were either cognitively unimpaired or had a clinical

E2 JAMA Published online June 16, 2025

Table 1. Characteristics of the Study Participants

Baseline Characteristics	Cognitively unimpaired (n = 3487)	Mild cognitive impairment (n = 1326)	Alzheimer disease dementia (n = 1332)	Dementia with Lewy bodies (n = 81)	Behavioral- variant FTD (n = 69)	Semantic variant PPA (n = 29)	Nonfluent variant PPA (n = 29)	Parkinson disease (n = 31)	Progressive supra- nuclear palsy (n = 73)	Corticobasal syndrome (n = 57)
Age, median (IQR), y	70 (65 to 77)	72 (65 to 78)	69 (60 to 76)	70 (66 to 75)	65 (58 to 69)	67 (63 to 70)	70 (58 to 75)	71 (65 to 75)	71 (66 to 74)	70 (64 to 74)
Sex, No. (%)										
Female	1875 (53.8)	565 (42.6)	695 (52.2)	17 (21.0)	33 (47.8)	9 (31.0)	19 (65.5)	11 (35.5)	36 (49.3)	29 (50.9)
Male	1612 (46.2)	761 (57.4)	637 (47.8)	64 (79.0)	36 (52.2)	20 (69.0)	10 (34.5)	20 (64.5)	37 (50.7)	28 (49.1)
APOE ε4 carrier, No (%)	1069 (34.2)	453 (44.4)	617 (63.3)	24 (42.1)	6 (27.3)	2 (25.0)	1 (33.3)	NA	NA	0
Not assessed	363	305	358	24	47	21	26			56
CDR, median (range)										
Global ^a	0 (0 to 0.5)	0.5 (0 to 2)	1 (0 to 3)	1 (0 to 3)	1 (0 to 2)	0.5 (0 to 2)	0.5 (0 to 3)	0.5 (0 to 2)	0.5 (0 to 3)	0.5 (0 to 2)
Sum of Boxes ^b	0 (0 to 4)	1.5 (0 to 12)	4.5 (0 to 18)	4.8 (1.5 to 17)	5.5 (0.5 to 12)	3 (0 to 7.5)	1.5 (0 to 14)	4 (0 to 12)	4 (0 to 15)	1.5 (0 to 10)
Not assessed	534	148	211	32	15	10	9	0	40	22
Αβ ΡΕΤ										
Centiloid, mean (range) ^c	26 (-37 to 212)	49 (-39 to 248)	88 (-31 to 196)	50 (-6 to 157)	10 (-32 to 122)	25 (-7 to 113)	16 (-14 to 141)	4 (-5 to 44)	9 (-11 to 69)	8 (-34 to 127)
Aβ positive (>24.4 Centiloids or positive visual read), No. (%)	1074 (34.1)	695 (57.9)	1050 (90.8)	35 (59.3)	8 (18.2)	5 (29.4)	3 (16.7)	1 (10.0)	5 (19.2)	9 (17.0)
Not assessed	342	126	175	22	25	12	11	21	47	4
Tau PET										
Positive, No. (%) ^d	343 (9.8)	605 (45.6)	1165 (87.5)	29 (35.8)	23 (33.3)	20 (69.0)	9 (31.0)	9 (29.0)	3 (4.1)	15 (26.3)
Follow-up characteristics ^e										
Participants with follow-up data, No.										
Clinical diagnosis	1977	490	NA	NA	NA	NA	NA	NA	NA	NA
CDR	2253	647	560	NA	NA	NA	NA	NA	NA	NA
Follow-up time, median (maximum), y ^f										
Clinical diagnosis	3.8 (7.1)	2.9 (6.8)	NA	NA	NA	NA	NA	NA	NA	NA
CDR	4.0 (8.1)	1.7 (7.0)	1.5 (6.4)	NA	NA	NA	NA	NA	NA	NA

Abbreviations: A β , amyloid- β ; APOE ϵ 4, apolipoprotein ϵ 4 allele; CDR, Clinical Dementia Rating; FTD, frontotemporal dementia; NA, not assessed/available; PET, positron emission tomography; PPA, primary progressive aphasia.

diagnosis of AD dementia within 1 year of death, and were assessed neuropathologically post mortem (neuropathology cohort, n = 3178). None of these participants had available tau PET scans.

PET Imaging

All tau PET scans were performed with flortaucipir F 18 and classified as either positive or negative using its FDA- and EMA-approved visual interpretation method. ¹² This method de-

fines a scan as positive if it exhibits increased signal in posterolateral temporal, occipital, and/or parietal/precuneus regions, with or without elevated signal in frontal areas. A visually positive flortaucipir F 18 PET scan reflects Braak stages V and VI.¹³ Additional details can be found in eMethods 1.2 in Supplement 1.

To visually assess the large number of tau PET scans in our sample, we gathered a team of 18 readers, including nuclear medicine physicians and neuroimaging researchers,

E3

jama.com JAMA Published online June 16, 2025

^a CDR-Global scores range from O (best) to 3 (worst).

^b CDR-Sum of Boxes scores range from 0 (best) to 18 (worst).

 $^{^{\}rm c}$ Assessed with [18 F] florbetapir, [18 F] florbetaben, [18 F] NAV4694, or [11 C] PiB.

 $^{^{\}rm d}$ Assessed using the approved visual interpretation method for flortaucipir F18 PET.

 $^{^{\}rm e}$ Longitudinal analyses included only participants with A β status available at baseline. For those with mild cognitive impairment or Alzheimer disease dementia, a baseline CDR-G score of 0.5 or 1 was also required.

 $^{^{\}rm f}$ Median follow-up times were estimated using the reverse Kaplan-Meier method. $^{\rm 42}$

who were trained using materials provided by Avid Radio-pharmaceuticals to interpret flortaucipir F 18 PET scans (eMethods 1.3-1.4 in Supplement 1). The robustness of our visual interpretation procedure was validated in interrater agreement analyses (Cohen κ across readers; mean, 0.89; range, 0.78-0.97; eResults 2.1 and eFigure 1 in Supplement 1).

A β PET scans were available for most of the study participants (n = 5729, 87.9%). A β PET positivity was defined as a Centiloid³⁰ higher than 24.4³¹ or with visual reads.

Outcomes

The outcomes were (1) frequency of visual tau PET positivity as a function of clinical diagnosis, age, AB PET status, sex, and apolipoprotein E (APOE) &4 carriership; and (2) clinically relevant progression among tau PET-positive individuals. For the latter, we studied 3 different definitions of relevant progression as perceived by patients, caregivers, and clinicians^{32,33}: (1) progression to a clinical diagnosis of MCI or dementia among initially cognitively unimpaired individuals or progression to dementia among individuals with MCI; (2) progression on the Clinical Dementia Rating (CDR) Global score (CDR-G), defined as any increase in the score³⁴; and (3) group-level mean CDR Sum of Boxes (CDR-SB) score after 5 years (baseline age, 70 years) higher than 1 for cognitively unimpaired, 5 for MCI (baseline CDR-G, 0.5-1), and 10 for mild AD dementia (baseline CDR-G, 0.5-1), which correspond to scores consistent with MCI, mild AD dementia, and moderate AD dementia, 35 respectively. The CDR-G score ranges from 0 to 3, with higher values indicating more severe cognitive and functional impairment. The CDR-SB score ranges from 0 to 18, based on the sum of ratings across 6 domains; higher scores reflect greater overall impairment.

Clinical progressions to MCI or dementia at follow-up visits were established by the respective study clinicians and committees based on published criteria. ^{21,22,36} The CDR was administered through a semistructured interview with both the participant and their study partner. While outcome assessors were not fully blinded to tau PET scans, clinical assessments were conducted independently of biomarker and CDR results, and tau PET visual reads were performed only after outcome ascertainment.

Statistical Analysis

We used generalized additive models to derive frequency estimates of tau PET positivity conditional on covariates (age, sex, APOE $\varepsilon 4$ carriership, and A β PET status) and diagnostic categories. Educational attainment was not included as a covariate due to its limited association with clinical decline, ^{37,38} age-cohort differences, and the lack of a standardized education metric across the included studies. Of note, the models account for the impact of studies that overrepresent A β -positive individuals, such as the A4 study, on the frequency of tau PET positivity. Ninety-five percent confidence intervals were calculated using either the Wald approximation when estimates were derived directly from model predictions or using the bootstrap method (n = 1000 repetitions) when estimates involved combining predictions from multiple models. Additional details can be found in eMethods 1.5 in Supplement 1.

Generalized additive mixed models were used to estimate mean temporal trajectories of the CDR-SB score (eMethods 1.6 in Supplement 1). Ninety-five percent confidence intervals were calculated using the Wald approximation.

The probability of clinical progression over the next 5 years (for cognitively unimpaired and MCI) or 3 years (for AD dementia) across different Aβ PET/tau PET profiles was estimated using illness-death models, which is the simplest form of a multistate Markov model.³⁹ This approach combines the naive hazard functions for clinical progression and mortality to account for the competing risk of death during follow-up precluding clinical progression, allowing for the estimation of unbiased probabilities of progression (eMethods 1.7 in Supplement 1). The naive hazard functions describing clinical progression were estimated using either Kaplan-Meier estimators (for global absolute risk estimates) or Cox proportional hazards regression (for age- and sexspecific absolute risk estimates). Because death events were not consistently recorded across all cohorts included in this study, we followed a previously published approach⁴⁰ and estimated naive hazard functions for mortality based on publicly available life tables from the 2019 US general population (https://mortality.org/). These estimates were further adjusted to account for the increased mortality risk associated with MCI or dementia⁴¹ (eMethods 1.7 in Supplement 1). Ninetyfive percent confidence intervals of the estimated absolute risks were computed using bootstrap (n = 1000 repetitions). Median follow-up times were estimated using the reverse Kaplan-Meier method.⁴²

Analyses of the clinical outcomes of the A β PET-negative/tau PET-positive groups were regarded as exploratory due to reduced sample size and reported in eFigure 9 in Supplement 1.

Analyses of between-cohort variability in the frequency of positivity and risk estimates were performed by estimating the additional explained variation that can be attributed to cohort membership (eMethods 1.8 in Supplement 1).

All analyses were conducted using the full pooled data-set, with statistical models run separately for cognitively unimpaired and cognitively impaired individuals. A complete-case approach was applied, as the proportion of missing A β PET, APOE ϵ 4, and CDR data was relatively small compared with the overall dataset. Sensitivity analyses using multiple imputation yielded results consistent with the primary findings, indicating that the impact of the missing data was minimal. A 2-sided P value less than .05 was considered statistically significant. R version 4.3.2 (R Foundation for Statistical Computing) and MATLAB 2023b (The MathWorks) were used for all statistical analyses.

Results


A total of 6514 participants with tau PET scans (mean age, 69.5 years; 50.5% female) pooled from 21 different cohorts were included. Baseline and follow-up characteristics of the included participants are reported in Table 1 and eTables 2-9 and eFigure 2 in Supplement 1.


JAMA Published online June 16, 2025


E4


jama.com

Figure 1. Frequency of Tau Positron Emission Tomography (PET) Positivity in Cognitively Unimpaired Individuals, Mild Cognitive Impairment, and Patients With a Clinical Diagnosis of Various Neurodegenerative Disorders









Estimated frequency of tau PET positivity, established using the approved visual interpretation method for flortaucipir F 18, as a function of age in cognitively unimpaired individuals (A), participants with mild cognitive impairment (B), and Alzheimer disease dementia (C). The cognitively unimpaired, mild cognitive impairment, and Alzheimer disease dementia groups were defined based solely on the respective clinical symptoms, independent of amyloid- β status or other biomarker results. Benchmark estimates of the frequency of Braak stages V and VI (orange lines) in cognitively unimpaired individuals and participants with Alzheimer disease dementia were derived using data from an independent sample of participants from the neuropathology cohort (n = 3178).

Panel D shows the overall frequency of tau PET positivity in other clinically diagnosed (ie, independent of amyloid- β status or other biomarker results) neurodegenerative disorders. Whiskers represent 95% confidence intervals. The number of participants with other neurodegenerative disorders were (tau PET positive/total): dementia with Lewy bodies, 29/81; behavioral variant frontotemporal dementia, 23/69; semantic variant primary progressive aphasia, 20/29; nonfluent variant primary progressive aphasia, 9/29; Parkinson disease with cognitive impairment, 9/31; progressive supranuclear palsy, 3/73; and corticobasal syndrome, 15/57. The numbers in the tables in A-C present the number for the decile they lie within.

Frequency of Tau PET Positivity

Of 3487 cognitively unimpaired participants, 349 (9.8%) were tau PET positive. Quantitative neuroimaging analyses revealed significantly elevated tau PET signal in temporoparietal and frontal regions (eResults 2.2 and eFigure 3 in Supplement 1). The estimated frequency of tau PET positivity in cognitively unimpaired individuals younger than 50 years was below 1%, and increased from 3% (95% CI, 2%-4%) at 60 years to 19% (95% CI, 16%-24%) at 90 years (Figure 1A; eTable 10 in


Supplement 1). The estimated frequency of Braak stages V and VI in cognitively unimpaired participants from the neuropathology cohort paralleled that of tau PET positivity until approximately 60 years of age but was consistently lower for older ages (Figure 1A; eTable 10 in Supplement 1).

Tau PET positivity was more frequent among the 1326 patients with MCI and 1332 patients with AD dementia. In patients with MCI, the estimated frequency at 40 years was 26% (95% CI, 20%-33%) and increased with age (from 35%)

E5

jama.com JAMA Published online June 16, 2025

Figure 2. Frequency of Tau Positron Emission Tomography (PET) Positivity Among Amyloid- β (A β) PET-Positive and A β PET-Negative Individuals, Stratified by Clinical Diagnosis



Panels A and B show the estimated frequency of tau PET positivity as a function of age among $A\beta$ PET-positive and $A\beta$ PET-negative individuals, separately for cognitively unimpaired (dark blue), mild cognitive impairment (light blue), and

patients with a clinical diagnosis of Alzheimer disease dementia (orange). The numbers in the tables present the number for the decile they lie within.

[95% CI, 32%-39%] at 60 years to 51% [95% CI, 46%-57%] at 90 years; Figure 1B; eTable 10 in Supplement 1). Among participants with clinically diagnosed AD dementia, tau PET positivity frequency was estimated to be more than 85% for those younger than 65 years of age; however, its estimated frequency decreased with advancing age (from 88% [95% CI, 86%-91%] at 60 years to 65% [95% CI, 58%-72%] at 90 years; Figure 1C; eTable 10 in Supplement 1). This decreasing trend was also observed for the frequency estimates of Braak stages V and VI in the neuropathology cohort, which closely matched the frequency curve of tau PET positivity (Figure 1C; eTable 10 in Supplement 1).

Estimates of tau PET positivity for other neurodegenerative disorders are displayed in Figure 1D. The estimated frequency of positivity was relatively high in patients with frontotemporal dementia, particularly in those with semantic variant primary progressive aphasia (20/29, 69% [95% CI, 49%-85%]), the majority of whom were A β PET negative (eFigure 4D in Supplement 1).

Among cognitively unimpaired participants and those with a clinical diagnosis of MCI and AD dementia, tau PET positivity was more frequently observed in A β PET-positive than in A β PET-negative individuals (A β PET positive vs A β PET negative, cognitively unimpaired: 268/1074, 25% [95% CI, 22%-28%] vs 46/2071, 2% [95% CI, 2%-3%]; MCI: 511/695, 74% [95% CI, 70%-77%] vs 43/505, 9% [95% CI, 6%-11%]; AD dementia: 976/1050, 93% [95% CI, 91%-94%] vs 33/107, 31% [95% CI, 22%-40%]; eFigure 4A-C in Supplement 1). Older age was associated with higher frequency estimates of tau PET positivity among A β PET-positive cognitively unimpaired individuals, but with lower frequencies among A β PET-positive participants with MCI or AD dementia (Figure 2A; eTable 11 in Supplement 1). In A β PET-negative participants, the estimated frequency of positivity increased with age in the cognitively

unimpaired and MCI groups, while it remained approximately constant or decreased slightly for A β PET-negative participants with AD dementia (Figure 2B; eTable 11 in Supplement 1). Higher A β levels (Centiloids) were associated with increased probability of a positive tau PET scan in a dose-dependent manner (eFigure 5 in Supplement 1).

Frequency estimates of tau PET positivity according to age, A β PET status, sex, and APOE $\epsilon 4$ carriership can be seen in eFigures 6 and 7 and eTables 12 to 15 in Supplement 1. The estimated frequency of tau PET positivity was higher in AB PET-positive cognitively impaired females than in males (75year-old with MCI, 74% for females vs 62% for males, difference, 12% [95% CI, 6%-19%]; 75-year-old with AD dementia, 91% for females vs 85% for males, difference, 6% [95% CI, 3%-9%]; eFigure 6 in Supplement 1), but this association was not statistically significant in cognitively unimpaired individuals (eFigure 6C-D and eTables 12-14 in Supplement 1). APOE £4 carriership was associated with increased probability of tau PET positivity in both Aβ PET-negative and Aβ PETpositive cognitively unimpaired individuals (75-year-old Aβ PET negative: 4% for carriers vs 2% for noncarriers, difference, 2% [95% CI, 0%-3%]; 75-year-old Aβ PET positive: 32% for carriers vs 21% for noncarriers, difference, 11% [95% CI, 7%-17%]) and Aβ PET-positive impaired individuals (75year-old with MCI, 71% for carriers vs 59% for noncarriers, difference, 12% [95% CI, 5%-19%]; 75-year-old with AD dementia, 90% for carriers vs 85% for noncarriers, difference, 5% [95% CI, 2%-10%]) (eFigure 6A-B and eTables 12-14 in Supplement 1). These associations were largely attenuated in Aß PET-negative cognitively impaired individuals (eFigure 7 and eTable 15 in Supplement 1).

The percentage deviance attributable to cohort differences was 0.8% for cognitively unimpaired and 11.1% for symptomatic participants, indicating relatively low cohort-related

E6 JAMA Published online June 16, 2025

jama.com

variability in frequency estimates. Visual inspection of study-specific estimates also confirmed low variability (eFigure 8 in Supplement 1).

Clinical Outcomes

The Aβ PET-positive/tau PET-positive profile was associated with a significantly increased estimated risk of experiencing all the evaluated clinical outcomes compared with both the AB PET-negative/tau PET-negative and A β PET-positive/tau PETnegative profiles (Figure 3, Table 2; eTables 16-17 in Supplement 1). For example, among initially cognitively unimpaired individuals with an AB PET-positive/tau PET-positive profile, the cause-specific hazard ratio for progression to MCI or dementia was 8.1 (95% CI, 5.5-12.0; P < .001) compared with Aβ PET-negative/tau PET-negative cognitively unimpaired individuals, and 3.8 (95% CI, 2.6-5.6; *P* < .001) compared with those with an Aβ PET-positive/tau PET-negative profile. Similar findings were observed for AB PET-positive/tau PET-positive individuals with MCI, with a hazard ratio of 6.8 (95% CI, 4.1-11.0; *P* < .001) for progression to dementia compared with the Aβ PET-negative/tau PET-negative MCI group, and a hazard ratio of 4.7 (95% CI, 2.6-8.4; P < .001) compared with the Aβ PET-positive/tau PET-negative MCI group.

The Aβ PET-positive/tau PET-positive profile was associated with a high absolute risk of progression to MCI or dementia among initially cognitively unimpaired individuals (5year absolute risk: 57.4% [95% CI, 45.4%-70.8%]; Figure 3A; Table 2) as well as to progression to dementia among participants with MCI (5-year absolute risk: 69.9% [95% CI, 59.3%-81.2%]; Figure 3B; Table 2). Moreover, the 5-year absolute risk estimates of progression were dependent on baseline age and sex for both AB PET-positive/tau PET-positive cognitively unimpaired participants (60 years: 30% [95% CI, 17%-43%] for females vs 31% [95% CI, 17%-45%] for males; 80 years: 63% [95% CI, 52%-75%] for females vs 62% [95% CI, 50%-74%] for males; eTable 18 in Supplement 1) and Aβ PET-positive/tau PET-positive participants with MCI (60 years: 77% [95% CI, 59%-90%] for females vs 74% [95% CI, 55%-88%] for males; 80 years: 71% [95% CI, 55%-84%] for females vs 66% [95% CI, 55%-76%] for males; eTable 19 in Supplement 1). Similarly, AB PET-positive/tau PET-positive individuals were at high risk of progression on all the CDR-G outcomes (Figure 3C-D, Table 2; eTables 17 and 20-22 in Supplement 1). Additional estimates of the absolute risks of clinical progression, conditional on age and sex, are provided in eTables 18 to 22 in Supplement 1. At the group level, only the Aβ PET-positive/tau PET-positive groups surpassed specified thresholds on the CDR-SB at 5 years

The outcomes of the A β PET-positive/tau PET-negative groups were more similar to those of the corresponding A β PET-negative/tau PET-negative groups, particularly among symptomatic (MCI or AD dementia) participants (Figure 3; eTable 23 in Supplement 1). Analyses of the A β PET-negative/tau PET-positive groups revealed numerically higher rates of clinical decline compared with their A β PET-negative/tau PET-negative counterparts (eFigure 9 in Supplement 1).

Explainable variation in risk of clinical progression attributable to cohort membership ranged from 13.1% to 32.9%

(eTable 24 in Supplement 1). Percentage deviance on CDR-SB trajectories attributable to cohort membership was low (0.2% for cognitively unimpaired and 0.3% for symptomatic participants), and visual inspection of cohort-specific trajectories confirmed limited variability (eFigure 10 in Supplement 1).


Discussion

In this study that applied an FDA- and EMA-approved method for determining tau PET positivity in a large convenience sample from 21 pooled cohorts, the findings suggest that cognitively unimpaired older individuals have visually positive tau PET scans more frequently than previously suggested, 43 exceeding frequencies of 10% in those aged 75 years and older. Tau PET positivity in the cognitively unimpaired group was also systematically higher than the prevalence of Braak stages V and VI in an independent cognitively unimpaired sample from the neuropathology cohort (Figure 1A). This discrepancy likely reflects selection bias in the neuropathology cohort, 44 which overrepresents healthy older individuals who reached old age (median age at death, 88 years) without experiencing cognitive decline. As a result, this population tends to have a lower AD neuropathologic burden, leading to a lower frequency of Braak stages V and VI.²⁹ Moreover, the fact that most of the tau PET-positive cognitively unimpaired individuals were also Aβ positive (85%), exhibited elevated tau PET signal in regions consistent with Braak V and VI stages (eFigure 3 in Supplement 1), and were at high risk of clinical progression (Figure 3) suggest that the FDA- and EMA-approved method for defining tau PET positivity is indeed reflecting neurofibrillary tangle pathology in these individuals.

Previous studies have reported conflicting findings on the clinical outcomes of AB PET-positive/tau PET-positive cognitively unimpaired individuals, 16-18 leaving the implications of this biomarker profile unclear. These discrepancies may stem from substantial differences in how tau PET positivity was defined across studies, including variations in cut points, brain regions, quantification methods, and radiotracers. Consequently, these studies may lack generalizability to clinical practice, where only approved methods for visual tau PET assessment are used. In the present study that used an FDA- and EMAapproved method for determining tau PET positivity, an AB PET-positive/tau PET-positive profile in cognitively unimpaired individuals was associated with higher rates of progression to MCI or dementia over the following 5 years compared with their Aß PET-positive/tau PET-negative and Aß PET-negative/tau PET-negative counterparts. These results highlight the potential clinical relevance of AB PET and tau PET positivity in the absence of symptoms.

In AD dementia, the frequency curve of tau PET positivity closely matched that of Braak stages V and VI derived in the neuropathology cohort (Figure 1C), further supporting that a positive tau PET scan reflects advanced tau pathology in AD.¹³ The decrease in frequency with age is likely attributable to 2 factors: (1) with advancing age, neuropathological changes other than AD pathology are more likely to result in an AD-like clinical syndrome ^{45,46}; and (2) with advancing age, the likelihood

Figure 3. Clinical Outcomes of Tau Positron Emission Tomography (PET)-Positive Individuals



CDR-G score, by group. Crosses in A-D indicate censoring. E and F, Mean the respective Aβ PET-/tau PET- profile as reference.

symptoms) in cognitively unimpaired vs symptomatic individuals, according to

Aβ PET/tau PET profiles. Modeled trajectories are conditional on age (70 years).

iama.com

Panel insets show covariate-adjusted cause-specific hazard ratios (HRs), using

progression to MCI or dementia in cognitively unimpaired vs symptomatic $\,$

C and D, Kaplan-Meier-based estimates of the absolute risk of increasing

individuals according to different Aβ PET/tau PET biomarker profiles.

JAMA Published online June 16, 2025

Table 2. Absolute Risk of Clinical Progression, by Aβ PET/Tau PET Profile^a

	Absolute risk, % (95% CI) ^b				
Outcome	Aβ-/tau-	Aβ+/tau-	Aβ+/tau+		
Progression to MCI or dementia over the next 5 y among cognitively unimpaired	6.4 (4.7 to 8.2)	16.6 (12.5 to 21.8)	57.4 (45.4 to 70.8)		
Progression to dementia over the next 5 y among MCI	15.3 (9.5 to 22.2)	30.4 (15.9 to 47.8)	69.9 (59.3 to 81.2)		
Progressing CDR-G score over the next 5 y among cognitively unimpaired	7.8 (6.0 to 9.6)	20.0 (16.5 to 23.7)	56.1 (48.2 to 64.2)		
Progressing CDR-G score over the next 5 y among MCI	11.1 (5.9 to 17.0)	15.2 (6.6 to 24.9)	60.6 (49.2 to 72.6)		
Progressing CDR-G score over the next 3 y among AD dementia	33.9 (15.5 to 55.7)	32.4 (20.6 to 44.9)	67.2 (60.8 to 73.6)		

Abbreviations: A β , amyloid- β ; AD, Alzheimer disease; CDR-G, Clinical Dementia Rating Global score; MCI, mild cognitive impairment; PET, positron emission tomography.

probability of clinical progression accounting for the competing risk of death during follow-up; see eMethods 1.7 in Supplement 1 for details) to different outcomes among initially cognitively unimpaired individuals, as well as patients with MCI or AD dementia with different A β PET/tau PET biomarker profiles. 95% Confidence intervals were estimated using bootstrap.

of developing comorbid A β pathology increases. ⁶ These 2 phenomena lead to a higher frequency of elderly individuals with an AD-like clinical syndrome primarily caused by a non-AD condition, in whom comorbid A β pathology is present. ^{47,48}

An Aβ PET-positive/tau PET-positive profile was associated with increased incidence of clinically relevant outcomes in both participants with MCI and early AD dementia (Figure 3). These results suggest that tau PET positivity is a reliable indicator of AD pathology being a dominant contributor to clinical symptoms. However, interpretation of tau PET findings must consider that positivity can also occur in A β -negative patients. The reasons behind tau PET positivity in the absence of AB remain unclear, although explanations include flortaucipir F 18 binding to AD-like tau aggregates present in tangle-dominant forms of dementia, 49 binding to targets other than tau, and falsepositive A β PET and/or tau PET scans. It is also important to note that tau PET radiotracers other than flortaucipir F 18 may have different off-target binding profiles, 50 and the frequency of positivity in $A\beta$ -negative conditions with these radiotracers may differ from the estimates derived in this study.

Between 10% and 30% of the participants with a clinical diagnosis of dementia with Lewy bodies, behavioral variant frontotemporal dementia, semantic or nonfluent variant primary progressive aphasia, or corticobasal syndrome exhibited an A β PET-positive/tau PET-positive profile (eFigure 4 in Supplement 1). While A β positivity alone should not be used to infer that AD pathology is a primary contributor to symptoms in non-AD disorders, ⁴³ the relevance of concurrent A β and tau PET positivity in these patients remains uncertain. Future studies are needed to clarify the relevance of this profile in these non-AD conditions.

Tau PET positivity was more frequent in cognitively impaired females, consistent with prior reports of sex-related increases in tau PET signal. ⁵¹ This pattern was not observed in cognitively unimpaired individuals, possibly due to subthreshold or regionally restricted differences. Similarly, higher tau PET positivity rates among APOE ε4 carriers align with previously observed tau PET signal elevations associated with this genotype. ⁵²

Limitations

This study has several limitations. First, pooling data from independent studies may have introduced biases due to differ-

ences in study design. However, complementary analyses showed limited cohort-related variability, with relatively small between-cohort differences (eFigures 8 and 10 and eTable 24 in Supplement 1).

Second, cognitively unimpaired individuals were primarily recruited from research cohorts that are not representative of the general population, which may have introduced bias in the estimated frequency of tau PET positivity. However, comparisons with estimates derived solely from the Mayo Clinic Study of Aging—a population-based cohort—revealed only minimal differences (eFigure 11 in Supplement 1), suggesting that any potential bias is likely limited.

Third, absolute risk estimates were informed by incorporating 2019 US mortality rates into the estimates and may differ in populations with varying life expectancies. Fourth, despite being the largest tau PET study using clinically approved methods, the sample sizes for A β PET-negative/tau PET-positive individuals, other neurodegenerative disorders, and those aged older than 90 years were relatively small and should be interpreted with caution.

Fifth, tau PET quantification or alternative visual read methods might detect early tau deposition (eg, isolated mesial temporal signal) that is not captured by flortaucipir F 18's visual interpretation, which may also be linked to higher clinical progression risk. ¹⁶ Sixth, this study only included flortaucipir F 18 PET scans, as second-generation tau tracers currently lack a standardized, neuropathologically validated, and clinically approved definition of tau PET positivity. Seventh, many cohorts did not collect race or ethnicity data, preventing a robust analysis of the influence of these factors on tau PET positivity and clinical outcomes. Additionally, the majority of participants were White and thus the generalizability of these findings to more diverse populations remains to be confirmed.

Conclusions

Tau PET positivity is detectable in a small but nonnegligible proportion of individuals in the preclinical stages of AD, with increasing frequency in symptomatic stages of the disease. The combination of a positive tau PET scan with a positive $A\beta$ PET

 $[^]a$ Aß± and tau± represent Aß PET and tau PET status (positive/negative), respectively.

^b Shown are the estimated absolute risks of clinical progression (ie, the

scan was associated with an increased risk of clinically relevant outcomes across the entire AD spectrum. These find-

ings highlight the potential role of tau PET for staging ADrelated neuropathologic changes.

ARTICLE INFORMATION

Accepted for Publication: April 29, 2025. Published Online: June 16, 2025. doi:10.1001/jama.2025.7817

Author Affiliations: Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden (Moscoso, Heeman, Brum, Grothe, Schöll); Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg, Sweden (Moscoso, Heeman, Grothe, Schöll); Nuclear Medicine Department and Molecular Imaging Group, Instituto de Investigación Sanitaria de Santiago de Compostela, Travesía da Choupana s/n, Santiago de Compostela, Spain (Moscoso, Costoya-Sánchez, Aguiar); Department of Radiology, Mayo Clinic, Rochester, Minnesota (Raghavan, Jack, Vemuri); Universidade de Santiago de Compostela, Santiago de Compostela, Spain (Costoya-Sánchez); Centro de Investigación Biomédica en Red Sobre Enfermedades Neurodegenerativas, Instituto de Salud Carlos III, Madrid, Spain (Costoya-Sánchez, Silva-Rodríguez, Grothe): Department of Clinical Physiology, Sahlgrenska University Hospital, Gothenburg, Sweden (van Essen); Division of Nuclear Medicine and Molecular Imaging, Geneva University Hospital, Geneva, Switzerland (Mainta, Garibotto); Nuclear Medicine Department, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain (Camacho); Nuclear Medicine Department, Lucus Augusti University Hospital, Lugo, Spain (Rodríguez-Fonseca); Reina Sofía Alzheimer's Centre, CIEN Foundation, Instituto de Salud Carlos III, Madrid, Spain (Silva-Rodríguez, Grothe); Nuclear Medicine Department, Hospital Clinic Barcelona-Institut d'Investigacions Biomèdiques August Pi Sunyer, Barcelona, Spain (Perissinotti): Biomedical Research Networking Centre of Bioengineering, Biomaterials, and Nanomedicine, Instituto de Salud Carlos III. Barcelona, Spain (Perissinotti): Samsung Alzheimer Research Center, Research Institute for Future Medicine, Samsung Medical Center, Seoul. Korea (Gu, Seo); Department of Neurology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea (Yun, Seo); Laboratory of Neuroimaging and Innovative Molecular Tracers, Faculty of Medicine, Geneva University Neurocenter, University of Geneva, Geneva, Switzerland (Peretti, Garibotto); Geneva Memory Center, Department of Rehabilitation and Geriatrics, Geneva University Hospitals, Geneva, Switzerland (Ribaldi, Frisoni): Laboratory of Neuroimaging of Aging, University of Geneva, Geneva, Switzerland (Ribaldi, Frisoni); Radiology and Nuclear Medicine, Vrije Universiteit Amsterdam, VU University Medical Center, Amsterdam, the Netherlands (Coomans, van de Giessen); Amsterdam Neuroscience, Neurodegeneration, Amsterdam, the Netherlands (Coomans, van de Giessen, Ossenkoppele); Alzheimer Center Amsterdam, Neurology, Vrije Universiteit Amsterdam, University Medical Center, Amsterdam, the Netherlands (Coomans, Ossenkoppele); Graduate Program in Biological Sciences: Biochemistry, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil (Brum);

Molecular Imaging and Pharmacokinetic Modelling Group, Center for Research in Molecular Medicine and Chronic Diseases. University of Santiago de Compostela, Spain (Aguiar); Department of Nuclear Medicine, University Hospital Cologne, Cologne, Germany (Bischof, Drzezga); Research Center Juelich, Institute for Neuroscience and Medicine II, Molecular Organization of the Brain, Juelich. Germany (Bischof, Drzezga); German Center for Neurodegenerative Diseases, Bonn-Cologne, Germany (Drzezga); Department of Clinical Research Design & Evaluation, Samsung Advanced Institute for Health Science & Technology, Sungkyunkwan University, Seoul, Korea (Seo); Department of Health Science and Technology, Samsung Advanced Institute for Health Science & Technology, Sungkyunkwan University, Seoul, Korea (Seo); McConnell Brain Imaging Centre, McGill University, Montreal, Quebec, Canada (Villeneuve); Douglas Mental Health University Institute-Research Center, Verdun, Quebec, Canada (Villeneuve); Department of Psychiatry, McGill University, Montreal, Quebec, Canada (Villeneuve); Department of Clinical Neurosciences and Cambridge University Hospitals NHS Trust. University of Cambridge, Cambridge, United Kingdom (Malpetti, Rowe); UK Dementia Research Institute at University of Cambridge, Cambridge, United Kingdom (Malpetti); Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom (O'Brien); Medical Research Council Cognition and Brain Sciences Unit, University of Cambridge, Cambridge, United Kingdom (Rowe); Clinical Memory Research Unit, Department of Clinical Sciences Malmö, Lund University, Lund, Sweden (Ossenkoppele, Smith, Hansson); Helen Wills Neuroscience Institute, University of California Berkeley (Jagust); Memory Clinic, Skåne University Hospital, Lund University, Lund, Sweden (Smith, Hansson): Centre d'Imagerie Biomedicale, Geneva, Switzerland (Garibotto); Department of Neurology, Memory and Aging Center, Weill Institute for Neurosciences, University of California, San Francisco (Soleimani-Meigooni, La Joie, Rabinovici): Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, California (Soleimani-Meigooni); Alzheimer's Association, Chicago, Illinois (Carrillo); Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston (Dickerson, Sperling); Department of Radiology and Biomedical Imaging, University of California San Francisco (Rabinovici); Department of Neurology, Indiana University School of Medicine, Indianapolis (Apostolova); Department of Radiology and Imaging Sciences, Center for Neuroimaging, Indiana University School of Medicine Indianapolis, Indianapolis (Apostolova); Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis (Apostolova); Departments of Radiology, Washington University in St Louis School of Medicine, St Louis, Missouri (LaMontagne); Knight Alzheimer's Disease Research Center, Washington University in St Louis School of Medicine, St Louis, Missouri (LaMontagne); Avid, Eli Lilly and Company, Philadelphia, Pennsylvania (Pontecorvo); Gordon

Radiology, Massachusetts General Hospital, Harvard Medical School, Boston (Johnson); Center for Alzheimer Research and Treatment, Brigham and Women's Hospital, Boston, Massachusetts (Johnson, Sperling); Department of Veterans Affairs Medical Center, Center for Imaging of Neurodegenerative Diseases, San Francisco, California (Weiner); Department of Radiology and Biomedical Imaging, University of California San Francisco (Weiner); Department of Medicine, University of California San Francisco (Weiner); Department of Psychiatry and Behavioral Sciences, University of California San Francisco (Weiner); Department of Neurology, University of California San Francisco (Weiner); Department of Neurology, Mayo Clinic, Rochester, Minnesota (Petersen); Dementia Research Centre, Queen Square Institute of Neurology, University College London, London, United Kingdom (Schöll); Department of Neuropsychiatry, Sahlgrenska University Hospital, Mölndal, Sweden (Schöll).

Author Contributions: Dr Moscoso had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Moscoso, Carrillo, Johnson,

Concept and design: Moscoso, Carrillo, Johnson Sperling, Schöll.

Acquisition, analysis, or interpretation of data: Moscoso, Heeman, Raghavan, Costoya-Sánchez, van Essen, Mainta, Camacho, Rodríguez-Fonseca, Silva-Rodríguez, Perissinotti, Gu, Yun, Peretti, Ribaldi, Coomans, Brum, Grothe, Aguiar, Bischof, Drzezga, Seo, Villeneuve, Malpetti, O'Brien, Rowe, van de Giessen, Ossenkoppele, Jagust, Smith, Hansson, Frisoni, Gariblotto, Soleimani-Meigooni, Carrillo, Dickerson, La Joie, Rabinovici, Apostolova, LaMontagne, Pontecorvo, Weiner, Petersen, Jack, Vemuri, Schöll.

Drafting of the manuscript: Moscoso, Schöll.
Critical review of the manuscript for important
intellectual content: Heeman, Raghavan,
Costoya-Sánchez, van Essen, Mainta, Camacho,
Rodríguez-Fonseca, Silva-Rodríguez, Perissinotti,
Gu, Yun, Peretti, Ribaldi, Coomans, Brum, Grothe,
Aguiar, Bischof, Drzezga, Seo, Villeneuve, Malpetti,
O'Brien, Rowe, van de Giessen, Ossenkoppele,
Jagust, Smith, Hansson, Frisoni, Garibotto,
Soleimani-Meigooni, Carrillo, Dickerson, La Joie,
Rabinovici, Apostolova, LaMontagne, Pontecorvo,
Johnson, Sperling, Weiner, Petersen, Jack, Vemuri,
Schöll.

Statistical analysis: Moscoso.

Obtained funding: Bischof, Villeneuve, O'Brien, Rowe, Ossenkoppele, Jagust, Hansson, Frisoni, Carrillo, Dickerson, La Joie, Rabinovici, Apostolova, Johnson, Sperling, Schöll.

Administrative, technical, or material support:
Moscoso, Rodríguez-Fonseca, Silva-Rodríguez,
Peretti, Ribaldi, Coomans, Aguiar, Bischof, Seo,
Malpetti, Jagust, Smith, Hansson, Garibotto,
Carrillo, Dickerson, La Joie, LaMontagne, Petersen,
Jack. Vemuri.

Supervision: Heeman, Grothe, Drzezga, O'Brien, van de Giessen, Jagust, Garibotto, Carrillo, Apostolova, Johnson, Sperling, Schöll.
Other—visual interpretation of PET studies:
Perissinotti

Other—preprocessing and analysis of PET data: Gu.

E10 JAMA Published online June 16, 2025

jama.com

Center for Medical Imaging, Department of

Conflict of Interest Disclosures: Dr Silva-Rodríguez reported personal fees from Qubiotech Health Intelligence SL outside the submitted work. Dr Bischof reported personal fees from Life Molecular Imaging outside the submitted work. Dr Drzezga reported grants from DFG, German Research Foundation during the conduct of the study and nonfinancial support from Siemens Healthineers, Life Molecular Imaging, GE Healthcare, Avid Radiopharmaceuticals, Sofie, Eisai, Novartis/Advanced Accelerator Applications, and Ariceum Therapeutics; personal fees from Siemens Healthineers, Sanofi, GE Healthcare, Biogen, Novo Nordisk, Invicro, Novartis/AAA, Bayer Vital, Lilly, PeerView Institute for Medical Education, International Atomic Energy Agency; and stock from Siemens Healthineers, Lantheus Holding, and Lilly outside the submitted work. In addition. Dr Drzezga had patent No. EP3765097A1 issued. Dr Malpetti reported grants from Race Against Dementia and Alzheimer's Research UK (grant ARUK-RADF2021A-010) during the conduct of the study and personal fees from Astex Pharmaceuticals (consulting) outside the submitted work. Dr O'Brien reported receiving personal fees from Biogen, Roche, GE Healthcare, and Okwin outside the submitted work; serving as chair on the data and safety monitoring board for TauRx and Novo Nordisk; acting as chair of UK Alzheimer's Society Research Strategy Council; and receiving academic support for research from Avid/Lilly, Merck, UCB, and Alliance Medical. Dr Rowe reported grants from the National Institute for Health and Care Research (NIHR), Wellcome Trust, Medical Research Council, and PSP Association during the conduct of the study; personal fees from Asceneuron, Astex, Astronautx, Alector, Booster Therapeutics, Draig Therapeutics, Cumulus Neuroscience, Eisai, Ferrer, WAVE, Prevail, Clinical Ink, CuraSen, and SV Health outside the submitted work. Dr van de Giessen reported grants from ZonMw, Hersenstichting, Alzheimer Nederland, and Health-Holland paid to institution outside the submitted work and contract research with Heuron Inc. AC Immune, and Roche and consultancy agreements with IXICO and Life Molecular Imaging for reading PET scans, all support was paid to the institution. Dr Ossenkoppele reported research funding/support from Avid Radiopharmaceuticals, Janssen Research & Development, Roche. Quanterix, and Optina Diagnostics; has given lectures in symposia sponsored by GE Healthcare; received speaker fees from Springer; is an advisory board member for Asceneuron; and is a steering committee member for Biogen and Bristol Myers Squibb, all paid to his institutions. Dr Jagust reported grants from the National Institute on Aging (NIA) during the conduct of the study; personal fees from Lilly outside the submitted work: and equity in OptoCeutics and Molecular Medicine. Dr Smith reported grants from the Swedish federal government under the ALF agreement during the conduct of the study and personal fees from Hoffmann La Roche Ltd, Triolab, Novo Nordisk, and Eli Lilly outside the submitted work. Dr Hansson reported nonfinancial support from Roche and Lilly during the conduct of the study and being employed by Lilly. Dr Garibotto reported grants from Siemens Healthineers and GE Healthcare and personal fees from Novartis to institution outside the submitted work. Dr Carrillo reported being a full-time employee of the Alzheimer's Association. Dr Dickerson reported

personal fees from Biogen, Eisai, Lantheus, Lilly, Merck, and Novo Nordisk and grants from Ouanterix outside the submitted work. Dr La Joie reported personal fees from GE Healthcare and grants from the National Institutes of Health (NIH)/ NIA. Alzheimer's Association, and US Department of Defense outside the submitted work. Dr Rabinovici reported grants from the NIH and Rainwater Charitable Foundation during the conduct of the study and grants from Eli Lilly. GE Healthcare, Life Molecular Imaging, and Genentech and personal fees from Alector SAB, Bristol Myers Squibb SAB, Eli Lilly SAB, Johnson & Johnson DSMB, Merck SAB, Roche SAB, Novo Nordisk SAB, PeerView, and Medscape; and serving as associate editor at JAMA. Dr Pontecorvo reported being an employee and minor stockholder of Eli Lilly and Company. Dr Johnson reported grants from the NIH and Eli Lilly during the conduct of the study and consulting for Novartis, Merck, and Janssen outside the submitted work. Dr Sperling reported grants from Eli Lilly and Eisai and consulting for AbbVie, AC Immune, Acumen, Alector, Apellis, Biohaven, Bristol Myers Squibb, Genentech, Ionis, Janssen, NervGen, Oligomerix, Prothena, Roche, Vigil Neuroscience, and Vaxxinity outside the submitted work. Dr Weiner reported grants from NIH/NIA, Department of Defense, California Department of Public Health, Alzheimer's Association, Johnson & Johnson, Kevin & Connie Shanahan, Siemens, Biogen, Hillblom Foundation, GE, VUmc, Australian Catholic University, The Stroke Foundation, and Veterans Administration (also without compensation employer) and personal fees from Acadia Pharmaceuticals, Boxer Capital, Cerecin, Clario/BioClinica, Japan Society for Dementia Research, Eisai, Guidepoint, Health & Wellness Partners, Indiana University, LCN Consulting, Merck Sharp & Dohme (also honoraria and travel support), Duke University, Owkin, Novo Nordisk, ProMIS Neurosciences. Prova Education. Sai MedPartners. T3D Therapeutics, University of Southern California, WebMD, MEDA Corp, China Association for Alzheimer's Disease, Taipei Medical University, Cleveland Clinic, Banner Health, AD/PD Alzheimer's & Parkinson's Diseases Conference, Kenes, Learning Health Society Institute, Japan Society for Dementia Research, University of Toulouse, Korean Dementia Association, University of Madison Wisconsin, CTAD Congress, and National Center for Geriatrics and Gerontology and stock options from Alzeca, Alzheon Inc, and ALZpath outside the submitted work. Dr Petersen reported grants from the NIA, GHR Foundation, and Alexander Family Foundation during the conduct of the study; personal fees from Roche Inc, Genentech Inc, Eli Lilly and Company, Novartis, and Novo Nordisk and nonfinancial support from Eisai Inc outside the submitted work; royalties from Oxford University Press and UpToDate: and educational materials from Medscape. Dr Vemuri reported grants from the NIH outside the submitted work. Dr Schöll reported grants from Knut and Alice Wallenberg Foundation and Swedish Research Council during the conduct of the study; grants from European Union's Horizon Europe, NIH, Gates Ventures, National Research Foundation of Korea, Swedish state under the agreement between the Swedish government and the County Councils, the ALF-agreement, Swedish Brain Foundation. Swedish Alzheimer Foundation, Familien Rönströms Foundation, Sahlgrenska Academy at

Region R&D, Sahlgrenska Science Park, NIHR University College London Hospitals Biomedical Research Centre, Novo Nordisk, and BioArctic and personal fees from Eli Lilly and Company, Novo Nordisk, Roche, BioArctic, Triolabs; and nonfinancial support from Roche and Beckman Coulter outside the submitted work; and being a cofounder and shareholder of Centile Bioscience and serving as associate editor of Alzheimer's Research & Therapy. No other disclosures were reported.

Funding/Support: The Mayo Clinic Study of Aging data were obtained under the research grant UO1 AGO06786. Data collection and sharing for this project was funded in part by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (NIH grant UO1 AGO24904) and Department of Defense ADNI (Department of Defense award No. W81XWH-12-2-0012). ADNI is funded by the NIA, the National Institute of Biomedical Imaging and Bioengineering, and through contributions from the following: AbbVie, Alzheimer's Association, Alzheimer's Drug Discovery Foundation, Araclon Biotech, BioClinica Inc, Biogen, Bristol Myers Squibb Company, CereSpir Inc, Cogstate, Eisai Inc, Elan Pharmaceuticals Inc, Eli Lilly and Company, Eurolmmun, F. Hoffmann-La Roche Ltd and its affiliated company Genentech Inc, Fujirebio, GE Healthcare, IXICO Ltd, Janssen Alzheimer Immunotherapy Research & Development LLC, Johnson & Johnson Pharmaceutical Research & Development LLC, Lumosity, Lundbeck, Merck & Co Inc, Meso Scale Diagnostics LLC, NeuroRx Research, Neurotrack Technologies, Novartis Pharmaceuticals Corporation, Pfizer Inc. Piramal Imaging, Servier. Takeda Pharmaceutical Company, and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (https://fnih.org/). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for NeuroImaging at the University of Southern California. The A4 Study was funded by a public-private-philanthropic partnership, including funding from the NIH/NIA, Eli Lilly and Company, Alzheimer's Association, Accelerating Medicines Partnership, GHR Foundation, an anonymous foundation, and additional private donors, with in-kind support from Avid Radiopharmaceuticals, Cogstate, Albert Einstein College of Medicine, and the Foundation for Neurologic Diseases. The companion observational Longitudinal Evaluation of Amyloid Risk and Neurodegeneration (LEARN) Study was funded by the Alzheimer's Association and GHR Foundation. The A4 and LEARN Studies were led by Dr Sperling at Brigham and Women's Hospital, Harvard Medical School, and Paul Aisen, MD, at the Alzheimer's Therapeutic Research Institute (ATRI) at the University of Southern California. The A4 and LEARN Studies were coordinated by ATRI at the University of Southern California, and the data are made available under the auspices of Alzheimer's Clinical Trial Consortium through the Global Research & Imaging Platform (GRIP). The complete A4 Study Team list is available at https://www.actcinfo.org/a4-study-team-lists/. We acknowledge the dedication of the study participants and their study partners who made the

jama.com JAMA Published online June 16, 2025 E11

the University of Gothenburg, Västra Götaland

© 2025 American Medical Association. All rights reserved, including those for text and data mining, Al training, and similar technologies.

A4 and LEARN Studies possible. Data were provided in part by OASIS-3: Longitudinal Multimodal Neuroimaging (principal investigators [PIs]: Tammie L.S. Benzinger, MD, PhD; Daniel S. Marcus, PhD; John C. Morris, MD; NIH P30 AG066444, P50 AG00561, P30 NS09857781, P01 AG026276, P01 AG003991, R01 AG043434, UL1 TRO00448, R01 EB009352). AV-45 doses were provided by Avid Radiopharmaceuticals, a wholly owned subsidiary of Eli Lilly; and OASIS-3_AV1451 (Pls: Drs Benzinger and Morris; NIH P30 AG066444, AW00006993). AV-1451 doses were provided by Avid Radiopharmaceuticals, a wholly owned subsidiary of Eli Lilly. Data collection and dissemination of the data presented in this article were supported by the LEADS Consortium (R56/ UO1 AGO57195, funded by the NIA). The article has been reviewed by the LEADS Publications Committee for scientific content. The authors acknowledge the invaluable contributions of the participants in LEADS as well as the assistance of the investigators and support staff at each of the participating sites. The AO5, PXO1, and LZAX studies were supported by Avid Radiopharmaceuticals and/or Eli Lilly and Company. The National Alzheimer's Coordinating Center database is funded by NIA/NIH grant U24 AGO72122. NACC data are contributed by the NIAfunded Alzheimer's Disease Research Center: P30 AGO62429 (PL James Brewer, MD, PhD), P30 AG066468 (PI Oscar Lopez, MD), P30 AG062421 (PI Bradley Hyman, MD, PhD), P30 AG066509 (PI Thomas Grabowski, MD), P30 AG066514 (PI Marv Sano, PhD), P30 AGO66530 (PI Helena Chui, MD), P30 AG066507 (PI Marilyn Albert, PhD), P30 AGO66444 (PI John Morris, MD), P30 AGO66518 (PI Jeffrey Kaye, MD), P30 AG066512 (PI Thomas Wisniewski MD) P30 AG066462 (PLScott Small MD), P30 AG072979 (PI David Wolk, MD), P30 AG072972 (PI Charles DeCarli, MD), P30 AG072976 (PI Andrew Saykin, PsyD), P30 AG072975 (PI David Bennett, MD), P30 AG072978 (PI Neil Kowall, MD), P30 AG072977 (PI Robert Vassar, PhD), P30 AGO66519 (PI Frank LaFerla, PhD), P30 AG062677 (PI Ronald Petersen, MD, PhD), P30 AG079280 (PI Eric Reiman, MD), P30 AG062422 (PI Gil Rabinovici, MD), P30 AG066511 (PI Allan Levey, MD, PhD), P30 AG072946 (PI Linda Van Eldik, PhD), P30 AG062715 (PI Sanjay Asthana, MD, FRCP), P30 AG072973 (PI Russell Swerdlow, MD), P30 AG066506 (PI Todd Golde, MD, PhD), P30 AG066508 (PI Stephen Strittmatter, MD, PhD), P30 AG066515 (PI Victor Henderson, MD, MS), P30 AG072947 (PI Suzanne Craft, PhD), P30 AG072931 (PI Henry Paulson, MD, PhD), P30 AGO66546 (PI Sudha Seshadri, MD), P20 AGO68024 (PI Erik Roberson, MD, PhD), P20 AG068053 (PI Justin Miller, PhD), P20 AG068077 (PI Gary Rosenberg, MD), P20 AG068082 (PI Angela Jefferson, PhD), P30 AG072958 (PI Heather Whitson, MD), and P30 AG072959 (PI James Leverenz, MD). Data collected by SCAN and shared by NACC are contributed by the NIA-funded ADRCs as follows: Arizona Alzheimer's Center-P30 AG072980 (PI Eric Reiman, MD): R01 AG069453 (PI Eric Reiman [contact], MD); P30 AG019610 (PI Eric Reiman, MD); and the State of Arizona, which provided additional funding supporting our center; Boston University—P30 AG013846 (PI Neil Kowall MD); Cleveland ADRC-P30 AG062428 (James Leverenz, MD): Cleveland Clinic, Las Vegas-P20AG068053; Columbia-P50 AGO08702 (PI Scott Small MD); Duke/UNC

ADRC-P30 AG072958: Emory University-P30AG066511 (PI Levey Allan, MD, PhD): Indiana University-R01 AG19771 (PI Andrew Saykin, PsyD); P30 AG10133 (PI Andrew Saykin, PsyD); P30 AG072976 (PI Andrew Saykin, PsyD); RO1 AGO61788 (PI Shannon Risacher, PhD): RO1 AG053993 (PI Yu-Chien Wu, MD, PhD); U01 AGO57195 (PI Liana Apostolova, MD); U19 AGO63911 (PI Bradley Boeve, MD); and the Indiana University Department of Radiology and Imaging Sciences: Johns Hopkins-P30 AG066507 (PI Marilyn Albert, Phd); Mayo Clinic-P50 AGO16574 (PI Ronald Petersen MD PhD); Mount Sinai-P30 AGO66514 (PI Mary Sano, PhD); RO1 AGO54110 (PI Trey Hedden, PhD); RO1 AGO53509 (PI Trey Hedden, PhD); New York University-P30AG066512-01S2 (PI Thomas Wisniewski, MD): RO1AGO56031 (PI Ricardo Osorio. MD); RO1AGO56531 (PIs Ricardo Osorio, MD; Girardin Jean-Louis, PhD); Northwestern University-P30 AG013854 (PI Robert Vassar PhD); RO1 AGO45571 (PI Emily Rogalski, PhD); R56 AGO45571, (PI Emily Rogalski, PhD); RO1 AGO67781 (PI Emily Rogalski, PhD); U19 AGO73153 (PI Emily Rogalski, PhD); RO1 DC008552, (M.-Marsel Mesulam, MD); RO1 AG077444, (Pls M.-Marsel Mesulam, MD, Emily Rogalski, PhD); RO1 NSO75075 (PI Emily Rogalski, PhD); RO1 AGO56258 (PI Emily Rogalski, PhD); Oregon Health and Science University-P30 AG008017 (PI Jeffrey Kaye MD); R56 AG074321 (PI Jeffrey Kaye, MD); Rush University-P30 AGO10161 (PI David Bennett MD); Stanford-P30AG066515: P50 AG047366 (PI Victor Henderson MD MS); University of Alabama, Birmingham-P2O; University of California, Davis—P30 AG10129 (PI Charles DeCarli, MD); P30 AG072972 (PI Charles DeCarli, MD); University of California Irvine—P50 AG016573 (PI Frank LaFerla PhD); University of California, San Diego-P30AG062429 (PI James Brewer, MD, PhD); University of California, San Francisco-P30 AGO62422 (Gil D. Rabinovici, MD); University of Kansas-P30 AG035982 (Russell Swerdlow, MD); University of Kentucky-P30 AG028283-15S1 (Pls Linda Van Eldik, PhD, and Brian Gold, PhD); University of Michigan ADRC-P30AG053760 (PI Henry Paulson, MD, PhD), P30AG072931 (PI Henry Paulson, MD, PhD), Cure Alzheimer's Fund 200775 (PI Henry Paulson, MD, PhD), U19 NS120384 (PI Charles DeCarli, MD, University of Michigan Site PI Henry Paulson, MD, PhD), RO1 AGO68338 (PI Bruno Giordani, PhD, Carol Persad, PhD, Yi Murphey, PhD), S100D026738-01 (Pl Douglas Noll, PhD), RO1 AGO58724 (PI Benjamin Hampstead, PhD), R35 AG072262 (PI Benjamin Hampstead, PhD), W81XWH2110743 (PI Benjamin Hampstead, PhD), RO1 AG073235 (PI Nancy Chiaravalloti, University of Michigan Site PI Benjamin Hampstead, PhD), 1101RX001534 (PI Benjamin Hampstead, PhD), IRXO01381 (PI Benjamin Hampstead, PhD); University of New Mexico-P20 AGO68077 (Gary Rosenberg, MD); University of Pennsylvania-State of Pennsylvania project 2019NF4100087335 (PI David Wolk, MD); Rooney Family Research Fund (PI David Wolk, MD); RO1 AGO55005 (PI David Wolk, MD); University of Pittsburgh-P50 AG005133 (PI Oscar Lopez, MD); University of Southern California-P50 AG005142 (PI Helena Chui, MD); University of Washington-P50 AG005136 (PI Thomas Grabowski, MD); University of Wisconsin-P50 AG033514 (PI Sanjay Asthana, MD, FRCP); Vanderbilt University-P20 AG068082;

Wake Forest-P30AG072947 (PI Suzanne Craft. PhD); Washington University, St Louis-P01 AGO3991 (PI John Morris, MD): PO1 AGO26276 (PI John Morris, MD); P20 MH071616 (PI Dan Marcus); P30 AG066444 (PI John Morris, MD); P30 NSO98577 (PI Dan Marcus); RO1 AGO21910 (PI Randy Buckner); RO1 AGO43434 (PI Catherine Roe); RO1 EBOO9352 (PI Dan Marcus); UL1 TROOO448 (PI Brad Evanoff); U24 RRO21382 (PI Bruce Rosen); Avid Radiopharmaceuticals/Eli Lilly; Yale-P50 AGO47270 (PI Stephen Strittmatter, MD. PhD); RO1AGO52560 (PI Christopher van Dyck, MD; Richard Carson, PhD); RO1AGO62276 (PI Christopher van Dyck, MD); 1Florida-P30AG066506-03 (PI Glenn Smith, PhD); and P50 AG047266 (PI Todd Golde, MD, PhD). The data from the Mayo Clinic Alzheimer's Disease Research Center (P30 AG067677) was obtained under research grants from the NIA and GHR Foundation. Data from UCSF ADRC was supported by NIH/NIA P30-AG062422 (UCSF ADRC), NIH/NIA R35 AG072362 (GDR), and Rainwater Charitable Foundation (GDR) grants. Data from the Geneva Memory Center were funded by the following private donors under the supervision of the Private Foundation of Geneva University Hospitals: APRA: Association Suisse pour la Recherche sur la Maladie d'Alzheimer, Genève; Fondation Segré, Genève; Race Against Dementia Foundation, London, UK; Fondation Child Care, Genève: Fondation Edmond J. Safra, Genève; Fondation Minkoff, Genève; Fondazione Agusta, Lugano; McCall Macbain Foundation, Canada: Nicole et René Keller, Genève: and Fondation AETAS, Genève. The BioFINDER study was supported by the NIA (R01AG083740), European Research Council (ADG-101096455), Alzheimer's Association (ZEN24-1069572, SG-23-1061717) GHR Foundation Swedish Research Council (2022-00775), ERA PerMed (ERAPERMED2021-184), Knut and Alice Wallenberg Foundation (2022-0231), Strategic Research Area MultiPark (Multidisciplinary Research in Parkinson's Disease) at Lund University, Swedish Alzheimer Foundation (AF-980907), Swedish Brain Foundation (FO2021-0293), Parkinson Foundation of Sweden (1412/22), Cure Alzheimer's Fund, Rönström Family Foundation, Konung Gustaf V:s och Drottning Victorias Frimurarestiftelse, Skåne University Hospital Foundation (2020-0000028), Regionalt Forskningsstöd (2022-1259), and Swedish federal government under the ALF agreement (2022-Projekt0080). The precursor of flortaucipir F 18 was provided by Avid Radiopharmaceuticals. The precursor of [18F]flutemetamol was sponsored by GE Healthcare. Data from the Samsung Medical Center are supported by a grant from the Korea Health Technology R&D Project through the Korea Health Industry Development Institute, funded by the Ministry of Health & Welfare and Ministry of Science and ICT, Republic of Korea (grant No. HU20C0111, HU22C0170, and HU20C0414), the Korea Health Industry Development Institute (No. HU22C0052), the Ministry of Health Welfare, Republic of Korea (grant No. HR21C0885), the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2019R1A5A2027340 and NRF-2020R1A2C1009778), Institute of Information & communications Technology Planning & Evaluation (IITP) grant funded by the Korea government (MSIT) (No.2021-0-02068, Artificial Intelligence Innovation Hub), and the Korea National Institute of Health research project (2021-ER1006-02). Data

E12 JAMA Published online June 16, 2025 jama.com

from the University of Cologne was funded by the Deutsche Forschungsgemeinschaft (DFG), project ID 431549029-SFB 1451, and partially by DFG, DR 445/9-1. Dr Schöll receives funding from the Knut and Alice Wallenberg Foundation (Wallenberg Centre for Molecular and Translational Medicine; KAW2014.0363 and KAW2023.0371), the Swedish Research Council (2017-02869, 2021-02678, 2021-06545 and 2023-06188), the European Union's Horizon Europe research and innovation program under grant agreement No. 101132933 (AD-RIDDLE) and 101112145 (PROMINENT), the NIH (R01 AGO81394-01). Gates Ventures, the National Research Foundation of Korea (RS-2023-00263612), the Swedish state under the agreement between the Swedish government and the County Councils, the ALF-agreement (ALFGBG-813971 and ALFGBG-965326), the Swedish Brain Foundation (FO2021-0311 and FO2024-0372), the Swedish Alzheimer Foundation (AF-1011738), Familjen Rönström's Foundation, the Sahlgrenska Academy at the University of Gothenburg, the Västra Götaland Region R&D (VGFOUREG-995510) and Innovation platforms, Sahlgrenska Science Park, and the National Institute for Health and Care Research University College London Hospitals Biomedical Research Centre.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Group Information: The PREVENT-AD Research Group, Harvard Aging Brain Study, for the LEADS Consortium, and Alzheimer's Disease Neuroimaging Initiative members are listed in Supplement 2.

Disclaimer: Dr Rabinovici is Associate Editor of JAMA but was not involved in any of the decisions regarding review of the manuscript or its acceptance

Data Sharing Statement: See Supplement 3.

Additional Information: Data used in the preparation of this article were obtained from the Harvard Aging Brain Study (HABS; PO1AGO36694; https://habs.mgh.harvard.edu). The HABS study was launched in 2010, funded by the NIA, and is led by PIs Reisa A. Sperling, MD, and Keith A. Johnson, MD, at Massachusetts General Hospital/Harvard Medical School in Boston. BACS data were supported by AGO34570 and AGO67418 grants. Alzheimer Center Amsterdam is part of the neurodegeneration research program of Amsterdam Neuroscience. Data from Amsterdam UMC are supported by Stichting Alzheimer Nederland and Stichting VUmc fonds. Data from the University of Cambridge cohort are supported by the Cambridge Centre for Parkinson-Plus; the National Institute for Health Research, Cambridge Biomedical Research Centre (BRC-1215-20014; NIHR203312): Medical Research Council (MC_UU_00030/14; MR/T033371/1 JBR; and ref 146281); the Wellcome Trust (220258); the Association of British Neurologists; Patrick Berthoud Charitable Trust (RG99368); the Evelyn Trust (RG84654); and the Race Against Dementia Alzheimer's Research UK (ARUK-RADF2021A-010). Data used in the preparation of this article were obtained in part from the Pre-symptomatic Evaluation of Novel or Experimental Treatments for Alzheimer's Disease (PREVENT-AD) program data release 6.0. PREVENT-AD was launched in 2011 as

a \$13.5 million, 7-year public-private partnership using funds provided by McGill University, the Fonds de Recherche du Ouébec-Santé (FRO-S). an unrestricted research grant from Pfizer Canada, the Levesque Foundation, the Douglas Hospital Research Centre and Foundation, the Government of Canada, and the Canada Fund for Innovation. Private sector contributions are facilitated by the Development Office of the McGill University Faculty of Medicine and by the Douglas Hospital Research Centre Foundation (http://www.douglas.qc.ca/). The founders of the program were John C. S. Breitner, MD, MPH, Judes Poirier, PhD, and Pierre Etienne, MD, Douglas Hospital Research Centre and Faculty of Medicine, McGill University, Montréal, Quebec, Canada. The program's current director is Judes Poirier, PhD, the co-director is Sylvia Villeneuve, PhD, and the study coordinator is Jennifer Tremblay-Mercier, MSc. PREVENT-AD is the result of efforts of many other co-investigators from a range of academic institutions and private corporations, as well as an extraordinarily dedicated and talented clinical and technical assistant staff, students, and postdoctoral fellows. Patients are recruited from the greater Montréal area and more distant locations in Québec. For up-to-date information, see https://prevent-alzheimer.net/? page_id=42&lang=en and https://douglas.research. mcgill.ca/stop-ad-centre. Data used in preparation of this article were obtained from the Pre-symptomatic Evaluation of Novel or Experimental Treatments for Alzheimer's Disease (PREVENT-AD) program (https://douglas.research. mcgill.ca/stop-ad-centre), data release 6.0. A complete listing of PREVENT-AD Research Group can be found in the PREVENT-AD database (https://preventad.loris.ca/acknowledgements/ acknowledgements.php?date=[2024-09-22]). Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni. usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at https:// adni.loni.usc.edu/wp-content/uploads/how_to_ apply/ADNI_Acknowledgement_List.pdf.

REFERENCES

- 1. Blennow K, Mattsson N, Schöll M, Hansson O, Zetterberg H. Amyloid biomarkers in Alzheimer's disease. *Trends Pharmacol Sci.* 2015;36(5):297-309. doi:10.1016/j.tips.2015.03.002
- 2. Hansson O. Biomarkers for neurodegenerative diseases. *Nat Med.* 2021;27(6):954-963. doi:10. 1038/s41591-021-01382-x
- 3. Hyman BT, Phelps CH, Beach TG, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimers Dement*. 2012;8(1): 1-13. doi:10.1016/j.jalz.2011.10.007
- 4. Jagust W. Is amyloid-β harmful to the brain? insights from human imaging studies. *Brain*. 2016; 139(pt 1):23-30. doi:10.1093/brain/awv326
- 5. Rabinovici GD, Gatsonis C, Apgar C, et al. Association of amyloid positron emission tomography with subsequent change in clinical management among Medicare beneficiaries with mild cognitive impairment or dementia. *JAMA*. 2019;321(13):1286-1294. doi:10.1001/jama.2019. 2000

- **6**. Jansen WJ, Ossenkoppele R, Knol DL, et al; Amyloid Biomarker Study Group. Prevalence of cerebral amyloid pathology in persons without dementia: a meta-analysis. *JAMA*. 2015;313(19): 1924-1938. doi:10.1001/jama.2015.4668
- 7. Roberts RO, Aakre JA, Kremers WK, et al. Prevalence and outcomes of amyloid positivity among persons without dementia in a longitudinal, population-based setting. *JAMA Neurol*. 2018;75 (8):970-979. doi:10.1001/jamaneurol.2018.0629
- **8**. Dubois B, Villain N, Frisoni GB, et al. Clinical diagnosis of Alzheimer's disease: recommendations of the International Working Group. *Lancet Neurol*. 2021;20(6):484-496. doi:10.1016/S1474-4422(21) 00066-1
- 9. Jack CR Jr, Andrews JS, Beach TG, et al. Revised criteria for diagnosis and staging of Alzheimer's disease: Alzheimer's Association Workgroup. *Alzheimers Dement*. 2024;20(8):5143-5169. doi:10. 1002/alz.13859
- **10**. Arriagada PV, Growdon JH, Hedley-Whyte ET, Hyman BT. Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease. *Neurology*. 1992;42(3 pt 1): 631-639. doi:10.1212/WNL.42.3.631
- 11. Bierer LM, Hof PR, Purohit DP, et al. Neocortical neurofibrillary tangles correlate with dementia severity in Alzheimer's disease. *Arch Neurol*. 1995; 52(1):81-88. doi:10.1001/archneur.1995. 00540250089017
- 12. US Food and Drug Administration. TAUVID prescribing information. Accessed April 13, 2025. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/212123s000lbl.pdf
- Fleisher AS, Pontecorvo MJ, Devous MD Sr, et al; A16 Study Investigators. Positron emission tomography imaging with [18F]flortaucipir and postmortem assessment of Alzheimer disease neuropathologic changes. *JAMA Neurol*. 2020;77 (7):829-839. doi:10.1001/jamaneurol.2020.0528
- **14**. Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol*. 1991;82(4):239-259. doi:10.1007/BF00308809
- **15.** Haroutunian V, Purohit DP, Perl DP, et al. Neurofibrillary tangles in nondemented elderly subjects and mild Alzheimer disease. *Arch Neurol*. 1999;56(6):713-718. doi:10.1001/archneur.56.6.713
- **16.** Ossenkoppele R, Pichet Binette A, Groot C, et al. Amyloid and tau PET-positive cognitively unimpaired individuals are at high risk for future cognitive decline. *Nat Med.* 2022;28(11):2381-2387. doi:10.1038/s41591-022-02049-x
- 17. Strikwerda-Brown C, Hobbs DA, Gonneaud J, et al; PREVENT-AD, HABS, and AIBL Research Groups. Association of elevated amyloid and tau positron emission tomography signal with near-term development of Alzheimer disease symptoms in older adults without cognitive impairment. *JAMA Neurol.* 2022;79(10):975-985. doi:10.1001/jamaneurol.2022.2379
- **18**. Jack CR Jr, Wiste HJ, Therneau TM, et al. Associations of amyloid, tau, and neurodegeneration biomarker profiles with rates of memory decline among individuals without dementia. *JAMA*. 2019;321(23):2316-2325. doi:10. 1001/jama.2019.7437
- 19. Costoya-Sánchez A, Moscoso A, Silva-Rodríguez J, et al; Alzheimer's Disease Neuroimaging Initiative and the Harvard Aging

- Brain Study. Increased medial temporal tau positron emission tomography uptake in the absence of amyloid- β positivity. *JAMA Neurol.* 2023;80(10): 1051-1061. doi:10.1001/jamaneurol.2023.2560
- **20**. Weigand AJ, Maass A, Eglit GL, Bondi MW. What's the cut-point? a systematic investigation of tau PET thresholding methods. *Alzheimers Res Ther*. 2022;14(1):49. doi:10.1186/s13195-022-00986-w
- **21**. Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med*. 2004;256(3):183-194. doi:10.1111/j.1365-2796.2004.01388.x
- 22. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7(3): 263-269. doi:10.1016/j.jalz.2011.03.005
- 23. McKeith IG, Boeve BF, Dickson DW, et al. Diagnosis and management of dementia with Lewy bodies: fourth consensus report of the DLB Consortium. *Neurology*. 2017;89(1):88-100. doi:10. 1212/WNL.00000000000004058
- **24**. Rascovsky K, Hodges JR, Knopman D, et al. Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain*. 2011;134(pt 9):2456-2477. doi:10.1093/brain/awr179
- **25.** Gorno-Tempini ML, Hillis AE, Weintraub S, et al. Classification of primary progressive aphasia and its variants. *Neurology*. 2011;76(11):1006-1014. doi:10. 1212/WNL.0b013e31821103e6
- **26**. Postuma RB, Berg D, Adler CH, et al. The new definition and diagnostic criteria of Parkinson's disease. *Lancet Neurol*. 2016;15(6):546-548. doi:10. 1016/51474-4422(16)00116-2
- 27. Höglinger GU, Respondek G, Stamelou M, et al; Movement Disorder Society-endorsed PSP Study Group. Clinical diagnosis of progressive supranuclear palsy: the movement disorder society criteria. *Mov Disord*. 2017;32(6):853-864. doi:10.1002/mds.26987
- **28.** Armstrong MJ, Litvan I, Lang AE, et al. Criteria for the diagnosis of corticobasal degeneration. *Neurology*. 2013;80(5):496-503. doi:10.1212/WNL. 0b013e31827f0fd1
- **29**. Besser LM, Kukull WA, Teylan MA, et al. The Revised National Alzheimer's Coordinating Center's Neuropathology Form—available data and new analyses. *J Neuropathol Exp Neurol*. 2018;77(8):717-726. doi:10.1093/jnen/nly049
- **30**. Klunk WE, Koeppe RA, Price JC, et al. The Centiloid Project: standardizing quantitative amyloid plaque estimation by PET. *Alzheimers Dement*. 2015;11(1):1-15.e1, 4. doi:10.1016/j.jalz.2014. 07.003

- **31.** La Joie R, Ayakta N, Seeley WW, et al. Multisite study of the relationships between antemortem [¹¹C]PIB-PET Centiloid values and postmortem measures of Alzheimer's disease neuropathology. *Alzheimers Dement*. 2019;15(2):205-216. doi:10. 1016/j.jalz.2018.09.001
- **32.** Tochel C, Smith M, Baldwin H, et al; ROADMAP consortium. What outcomes are important to patients with mild cognitive impairment or Alzheimer's disease, their caregivers, and health-care professionals? a systematic review. *Alzheimers Dement (Amst)*. 2019;11:231-247. doi:10.1016/j.dadm.2018.12.003
- **33.** Rentz DM, Wessels AM, Annapragada AV, et al. Building clinically relevant outcomes across the Alzheimer's disease spectrum. *Alzheimers Dement* (*N Y*). 2021;7(1):e12181. doi:10.1002/trc2.12181
- **34**. Lanctôt KL, Boada M, Tariot PN, et al. Association between clinical dementia rating and clinical outcomes in Alzheimer's disease. *Alzheimers Dement (Amst)*. 2024;16(1):e12522. doi:10.1002/dad2.12522
- **35.** O'Bryant SE, Waring SC, Cullum CM, et al; Texas Alzheimer's Research Consortium. Staging dementia using Clinical Dementia Rating Scale Sum of Boxes scores: a Texas Alzheimer's research consortium study. *Arch Neurol.* 2008;65(8):1091-1095. doi:10.1001/archneur.65.8.1091
- **36**. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. American Psychiatric Association; 2013.
- **37.** Jack CR Jr, Therneau TM, Lundt ES, et al. Long-term associations between amyloid positron emission tomography, sex, apolipoprotein E and incident dementia and mortality among individuals without dementia: hazard ratios and absolute risk. *Brain Commun.* 2022;4(2):fcac017. doi:10.1093/braincomms/fcac017
- **38**. Seblova D, Berggren R, Lövdén M. Education and age-related decline in cognitive performance: systematic review and meta-analysis of longitudinal cohort studies. *Ageing Res Rev*. 2020;58:101005. doi:10.1016/j.arr.2019.101005
- **39**. Putter H, Fiocco M, Geskus RB. Tutorial in biostatistics: competing risks and multi-state models. *Stat Med*. 2007;26(11):2389-2430. doi:10.1002/sim 2712
- **40**. Brookmeyer R, Abdalla N. Estimation of lifetime risks of Alzheimer's disease dementia using biomarkers for preclinical disease. *Alzheimers Dement*. 2018;14(8):981-988. doi:10.1016/j.jalz.2018.03.005
- **41**. Vassilaki M, Cha RH, Aakre JA, et al. Mortality in mild cognitive impairment varies by subtype, sex, and lifestyle factors: the Mayo Clinic Study of Aging. *J Alzheimers Dis*. 2015;45(4):1237-1245. doi:10. 3233/JAD-143078

- **42**. Schemper M, Smith TL. A note on quantifying follow-up in studies of failure time. *Control Clin Trials*. 1996;17(4):343-346. doi:10.1016/0197-2456(96) 00075-X
- **43.** Rabinovici GD, Knopman DS, Arbizu J, et al. Updated appropriate use criteria for amyloid and tau PET: a report from the Alzheimer's Association and Society for Nuclear Medicine and Molecular Imaging Workgroup. *Alzheimers Dement*. 2025;21 (1):e14338. doi:10.1002/alz.14338
- **44.** Gauthreaux K, Kukull WA, Nelson KB, et al. Different cohort, disparate results: selection bias is a key factor in autopsy cohorts. *Alzheimers Dement*. 2024;20(1):266-277. doi:10.1002/alz.13422
- **45.** Nelson PT, Dickson DW, Trojanowski JQ, et al. Limbic-predominant age-related TDP-43 encephalopathy (LATE): consensus working group report. *Brain*. 2019;142(6):1503-1527. doi:10. 1093/brain/awz099
- **46**. Crary JF, Trojanowski JQ, Schneider JA, et al. Primary age-related tauopathy (PART): a common pathology associated with human aging. *Acta Neuropathol*. 2014;128(6):755-766. doi:10.1007/s00401-014-1349-0
- **47**. Ossenkoppele R, Jansen WJ, Rabinovici GD, et al; Amyloid PET Study Group. Prevalence of amyloid PET positivity in dementia syndromes: a meta-analysis. *JAMA*. 2015;313(19):1939-1949. doi:10.1001/jama.2015.4669
- **48**. Landau SM, Lee J, Murphy A, et al; Alzheimer's Disease Neuroimaging Initiative. Individuals with Alzheimer's disease and low tau burden: characteristics and implications. *Alzheimers Dement*. 2024;20(3):2113-2127. doi:10.1002/alz.13609
- **49**. Jellinger KA, Attems J. Neurofibrillary tangle-predominant dementia: comparison with classical Alzheimer disease. *Acta Neuropathol*. 2007; 113(2):107-117. doi:10.1007/s00401-006-0156-7
- **50**. Van Wambeke É, Gérard T, Lhommel R, Hanseeuw B. Disclosing tau tangles using PET imaging: a pharmacological review of the radiotracers available in 2021. *Acta Neurol Belg*. 2022;122(2):263-272. doi:10.1007/s13760-021-01797-w
- **51.** Buckley RF, Mormino EC, Rabin JS, et al. Sex differences in the association of global amyloid and regional tau deposition measured by positron emission tomography in clinically normal older adults. *JAMA Neurol*. 2019;76(5):542-551. doi:10. 1001/jamaneurol.2018.4693
- **52**. Therriault J, Benedet AL, Pascoal TA, et al. Association of apolipoprotein E ϵ 4 with medial temporal tau independent of amyloid-β. *JAMA Neurol.* 2020;77(4):470-479. doi:10.1001/jamaneurol.2019. 4421

E14 JAMA Published online June 16, 2025