

Core Compendium:

Options on Fitting Cores to Data

Dan Korytov

February 16, 2019

Abstract

This document records the quality of fitting cores to red mapper galaxy clusters. There are too many options and iterations to keep everything in mind.

1 Overview of options

This section is broken down into two sections. The first section deals with the creating the radial galaxy distribution the cores will try to match. The second will go over the options on how to treat the cores.

1.1 Target Galaxy Distribution Options

These options are before we even use any core data. So combining all these options with the options of how the cores are treated give us the full option matrix.

- **Galaxy Luminosity:** The brightness relative to M^* cut off for it to be counted in the stack. I've only looked at the two options below. Most of the new core definition has been with $M_r < M^*$.

- $M_r < M^*$
- $M_r < M^* + 1$

- **Cluster Radial bins:** How the radial profile is stacked. Excluding the center did not change the fit significantly.

- 15 linear out to R_{200}
- 15 linear from $0.05/0.1/0.2R_{200}$ to R_{200}

- **Cluster Mass Stacking:** How the clusters are stacked by SO mass. Currently, it's only done by log linear bins. It might be thinking about making the bins wider in the high mass region to reduce the uncertainty. Empty bins aren't taken into consideration for the fit.

- x log linear bins from $1e14$ to $1e16$ **check**

- **Redshift Bins:** The clusters are stacked in redshift bins. There is almost no redshift evolution, so having only one redshift bin improves the statics.

- One bin from $0 < z < 0.35$
- Three bins from $0 < 0.10 < 0.20 < 0.3$

1.2 Core Option

- **Infall Mass Limit:** The minimum infall mass a core must posses for it be considered a candidate to be galaxy. Varying this infall mass while keeping other variables fixed, changes the overall abundance of cores in a cluster but does not change the shape of the profile significantly.

- On – always required

- **Core Disruption Radius:** How diffuse can a candidate core be before it's considered disrupted and removed as a candidate for a galaxy.

- On

- Off – current best fit

- **Core Merging Linking Length:** If two core candidates or more within are radius they are merged into one object. This uses an FoF merging algorithm for merging 3+ cores. The position is just the average position of the merged cores. **Maybe weighted by infall mass so that centrals are weighted towards the center?**

- On – current best fit

- Off

- **Disrupted Cores Merging:** Allowing the particles of disrupted cores merge into a galaxy. There was a linking length and minimum number of particles required. This option was used to solve the missing central problem which was solved by refreshing central cores.

- Off: not needed any more

- **Infall time pass?:** Maybe cores automatically disrupt/merge after a certain amount of time. Just a thought, not implemented.

- Off – not implement

- **Cluster Radius:** To construct the radial profile, all cores within a radius of the cluster center are used. I've tried some different radii and the parameters of the fit are pretty insensitive to the limit. Cores further out contribute a small fraction to the profile after smoothing for different angles.

- 3 Mpc/h
- 5 Mpc/h
- 8 Mpc/h

1.3 Fit Option

After deciding how the data and cores are treated, there are still several options on how to calculate the likelihood function.

- **Fitting radial profile:** The main mode of fitting. Each stack of clusters try to match the radial surface density profiles of galaxies/cores.
- **Fitting Ngal:** Fitting number of galaxies within R_{200} per each stack of clusters to match. The disagreement is weighted equally as the radial bins. Each stack has 15 radial bins errors and only 1 Ngal error, so it's a bit under weighted. **Maybe this can work well with global abundance?** Just as a side experiment
- **Fitting Accumulated Core profile:** Error bars are highly correlated and don't get the right treatment for the correlation. This is sort of an up-weighted Ngal requirement. Just of exploratory fitting.
- **Minimum Number of Clusters in Stack:** Some stacks of clusters have only a few clusters and might not representative of that stack. The Poisson error will be large, so they don't provide too strong of constraints. They could be excluded from the analysis complete or left in.
 - No Requirement
 - at least X
- **Locking the Global Abundance:** Requires that the abundance of cores per unit volume to match the expected abundance of M^*+x galaxies. This is done by fitting core disruption parameter while adjusting the infall mass to match the expected abundance. **Note:** While fitting for the abundance and profiles, we can't get a good fit to the profile. **Is this only a two parameter fit? Can I look at merging as well?**
- **Fitting Global Abundance:** Instead of locking the parameters to follow the global abundance of galaxies, the global abundance is taking into account for the likelihood evaluation. See link_to_appendix for an example on the effect on the likelihood/ cost function. So far the fit looks pretty good. :)




Figure 1: Model Parameters for fitting different data sets and models. The galaxies modeled in this figure are Mstar+0 galaxies.

- **2-point function:** Not implemented.

2 Fit Procedure

2.1 Error

The RMS of the Poisson noise and error on the mean for each stacked radial bin.

2.2 Posterior

Either through MCMC or grid search. The low dimensionality of the problem (only two or three parameters) lend to faster and more accurate results by a flat grid search.

3 Fit Results

A Mstar+0, M200m

A.1 M200m, Mstar+0, Profile fit, Disruption Only




Figure 2: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1 σ areas shaded in blue. The 2D likelihood distributions have 1 σ and 2 σ contours




Figure 3: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1 σ areas shaded in blue. The 2D likelihood distributions have 1 σ and 2 σ contours




Figure 4: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 5: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

A.2 M200m, Mstar+0, Profile fit, Merging Only




Figure 7: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 8: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 9: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 10: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

A.3 M200m, Mstar+0, Profile fit, Disruption + Merging




Figure 12: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 13: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 14: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 15: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

A.4 M200m, Mstar+0, Profile + Abundance fit, Disruption Only




Figure 17: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 18: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 19: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 20: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

A.5 M200m, Mstar+0, Profile + Abundancefit, Merging Only




Figure 22: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 23: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 24: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 25: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

A.6 M200m, Mstar+0, Profile + Abundance fit, Disruption + Merging




Figure 27: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 28: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 29: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 30: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

B M200c, Mstar+0

B.1 M200c, Mstar+0, Profile fit, Disruption Only




Figure 32: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 33: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 34: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 35: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

B.2 M200c, Mstar+0, Profile fit, Merging Only




Figure 37: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 38: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 39: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 40: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

B.3 M200c, Mstar+0, Profile fit, Disruption + Merging




Figure 42: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 43: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 44: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 45: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

B.4 M200c, Mstar+0, Profile + Abundance, Disruption Only




Figure 47: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 48: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 49: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 50: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

B.5 M200c, Mstar+0, Profile + Abundance, Merging Only




Figure 52: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 53: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 54: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 55: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

B.6 M200c, Mstar+0, Profile + Abundance, Disruption + Merging




Figure 57: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 58: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 59: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 60: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

C Mstar+1, M200m

C.1 M200m, Mstar+1, Profile fit, Disruption Only




Figure 62: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 63: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 64: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 65: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

C.2 M200m, Mstar+1, Profile fit, Merging Only




Figure 67: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 68: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 69: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 70: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a

C.3 M200m, Mstar+1, Profile fit, Disruption + Merging




Figure 72: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 73: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




Figure 74: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours







Figure 75: Likelihood on a parameter grid around the best mode. The marginalized parameter likelihood have 1σ areas shaded in blue. The 2D likelihood distributions have 1σ and 2σ contours




(a) a




(b) a



(c) a



(d) a



(e) a