

Math 6030, Graduate Algebra, Spring 2025, Homework 7

Instructor: Danny Krashen

Discussing the problems with other people is encouraged,
but you must write up your own work independently!

1. Find at least two different minimal primary decompositions of the ideal $I = \langle xy, xz, yz \rangle$ in $\mathbb{C}[x, y, z]$.
2. Let I, J be ideals in a commutative ring R such that $I + J = R$. Prove that:
 - (a) $IJ = I \cap J$.
 - (b) The natural map $\psi : R/IJ \rightarrow R/I \times R/J$, given by $\psi(x + IJ) = (x + I, x + J)$, is an isomorphism.

3. Let $\phi_i : R \rightarrow R_i$ be surjective maps for $i = 1, 2, \dots, n$. Suppose that for every pair of indices i, j , the induced map

$$R \rightarrow R_i \times R_j$$

is surjective. Prove that the map

$$\phi : R \rightarrow \prod_i R_i$$

is also surjective.

4. Prove that $R/IJ \cong R/I + R/J$ (via the natural projection maps) if and only if

$$R/\sqrt{IJ} \cong R/\sqrt{I} \times R/\sqrt{J}.$$

5. Suppose $f_1, \dots, f_r \in \mathbb{C}[x_1, \dots, x_n]$ and let $I = \langle f_1, \dots, f_r \rangle$ be the ideal generated by these. Suppose that $a = (a_1, \dots, a_n) \in \mathbb{C}^n$ and that $f_i(a) = 0$ for all i . Show that if $g \in I^m$, then we have

$$(\partial/\partial x_{i_1} \cdots \partial/\partial x_{i_\ell} g)(a) = 0 \text{ for } \ell < m.$$

6. (optional) For each of the following ideals, determine whether it is primary. Justify your answer.
 - (a) $\langle x^2, xy \rangle$ in $\mathbb{C}[x, y]$.
 - (b) $\langle x^2y, xy^2 \rangle$ in $\mathbb{C}[x, y]$.
 - (c) $\langle x^2 + y^2 \rangle$ in $\mathbb{C}[x, y]$.
 - (d) $\langle x^2 - 2x \rangle$ in $\mathbb{Z}[x]$.
 - (e) $\langle x(x - 1) \rangle$ in $\mathbb{Z}[x]$.