

arXiv Strategy for the Cut-and-Project Paper

Dirk Kunert

May 1, 2026

Purpose

This note collects practical suggestions for submitting the manuscript

Period Formulas for Rational Cut-and-Project Strip Projections: Multiset and Set Cases

to arXiv. It lists suitable subject categories, possible expert readers, and a suggested first-contact email.

The paper proves two complementary closed-form period formulas (one for the multiset of projected gap values, one for the underlying set) and shows that the set period is always at most the multiset period, with equality precisely when $N \leq D$. The algebraic core of both theorems is formalised in Lean 4 ($\sim 2,850$ lines, no `sorry`, `admit`, or `axiom`).

Can an independent researcher submit to arXiv?

Yes. An independent researcher may submit to arXiv. The issue is not the lack of a university affiliation, but the endorsement requirement. A new submitter normally needs endorsement for the relevant arXiv category or subject class.

The affiliation may simply be written as

Independent Researcher

in the paper.

Recommended arXiv categories

The paper is not best framed primarily as deep number theory. It uses number-theoretic tools such as congruences, residue classes, and coprimality, but its main object is a gap sequence arising from a geometric and combinatorial construction.

Recommended categories:

- Primary: `math.CO` or `math.DS`.
- Possible cross-listing: `math.NT` or `math.MG`.

A reasonable first choice is

`math.CO`

because the proofs are essentially combinatorial and residue-class based: both the multiset formula and the set-versus-multiset dichotomy reduce to counting cumulative residue classes modulo $D = \alpha^2 + \beta^2$. If the paper is presented more strongly as a symbolic-dynamical gap-sequence result, then `math.DS` is also plausible.

How to approach potential endorsers or expert readers

It is better not to send a cold message that immediately asks for endorsement. A more professional strategy is:

1. Send a concise email with the manuscript attached or linked.
2. Explain the result in two or three sentences.
3. Ask whether the result appears suitable as an arXiv preprint and whether the recipient knows prior literature containing the same formula.
4. Only if the response is positive, ask whether the person would be willing to endorse the arXiv submission.

This is less intrusive and more likely to receive a serious response.

Most relevant potential readers

The following people are mathematically close to the topic. They should not be viewed as guaranteed endorsers; they are possible expert readers whose published work is near cut-and-project sets, gap problems, symbolic dynamics, or combinatorics on words.

Closest to cut-and-project sets and gap problems

- **Alan Haynes.** Relevant because of work on gap problems and frequencies of patches in cut-and-project sets.
- **Henna Koivusalo.** Relevant because of work on cut-and-project sets, gap problems, and aperiodic order.
- **Lorenzo Sadun.** Relevant because of work on tilings, aperiodic order, and cut-and-project structures.
- **James Walton.** Relevant because of work on aperiodic order, topology of tilings, and cut-and-project structures.
- **Reem Yassawi.** Relevant because of work connecting symbolic dynamics, Meyer sets, and self-similar structures.
- **Valérie Berthé.** Relevant because of work on symbolic dynamics, combinatorics on words, numeration, and Meyer-set related structures.

Gap problems, lattices, and the three-gap theorem

- **Jens Marklof.** Relevant because of work on the three-gap theorem, spaces of lattices, and geometric approaches to gap problems.
- **Andreas Strömbergsson.** Relevant because of work on lattices, dynamics, and the three-gap theorem.
- **Jayadev Athreya.** Relevant through dynamics, lattices, and related gap-distribution questions, although somewhat less directly tied to this specific rational cut-and-project problem.
- **Barak Weiss.** Relevant through homogeneous dynamics, Delone sets, and aperiodic order, although the present result may be more elementary than his usual focus.

Christoffel words, Sturmian sequences, and combinatorics on words

- **Jean Berstel.** Central figure in combinatorics on words, Sturmian words, and Christoffel words.
- **Christophe Reutenauer.** Relevant because of major work on Christoffel words and combinatorics on words.
- **Aaron Lauve.** Relevant because of work on Christoffel words and related combinatorics.
- **Franco V. Saliola.** Relevant because of work on Christoffel words and combinatorics on words.
- **Sébastien Labbé.** Relevant because of work on Christoffel words, multidimensional symbolic dynamics, and cut-and-project related structures.
- **Michel Rigo.** Relevant because of work on combinatorics on words, numeration systems, and symbolic dynamics.

Shortlist

If only a few people are contacted, a sensible order is:

1. Alan Haynes.
2. Henna Koivusalo.
3. James Walton.
4. Valérie Berthé.
5. Sébastien Labbé.
6. Jens Marklof.

The first three are closest to cut-and-project gap questions. Berthé and Labbé are strong choices if the paper is framed through symbolic dynamics and combinatorics on words. Marklof is relevant through the lattice and gap-problem viewpoint.

Suggested first-contact email

Subject: Short note on rational cut-and-project gap periods

Dear Professor [Name],

I am an independent researcher working on a short note about rational one-dimensional cut-and-project strip projections.

The paper proves two complementary closed-form formulas for the minimal period of the projected gap sequence. For coprime positive integers α , β and window parameter $\omega \geq 0$, set

$$\begin{aligned} N &= \text{floor}(\omega \alpha) + \text{floor}(\omega \beta) + 1, \\ D &= \alpha^2 + \beta^2. \end{aligned}$$

Counting projected values with multiplicity gives

$\lambda_{\text{multiset}} = N$ if D does not divide N ,
 $\lambda_{\text{multiset}} = N / D$ if D divides N ,

and discarding multiplicities gives

$\lambda_{\text{set}} = N$ if $N < D$,
 $\lambda_{\text{set}} = 1$ if $N \geq D$.

In particular $\lambda_{\text{set}} \leq \lambda_{\text{multiset}}$, with equality iff $N \leq D$.

The proofs are elementary and reduce the problem to the distribution of consecutive residue classes modulo D . The algebraic core of both theorems has also been formalised in Lean 4 (no sorry, admit, or axiom).

Since your work is close to cut-and-project sets / gap problems / symbolic dynamics, I would be grateful if you were willing to take a brief look at the manuscript. In particular, I would appreciate knowing whether the result appears to be a reasonable arXiv preprint and whether you are aware of prior literature containing this exact formula.

I fully understand if you do not have time.

Best regards,
Dirk Kunert

Possible follow-up email asking for endorsement

This should only be sent if the person has responded positively or has shown interest in the manuscript.

Subject: arXiv endorsement question

Dear Professor [Name],

thank you very much for taking the time to look at my manuscript.

I would now like to submit the note to arXiv, probably under [math.CO/math.DS]. Since I am an independent researcher and a first-time submitter in this area, arXiv requires an endorsement.

If you consider the manuscript suitable as a refereeable mathematical preprint, would you be willing to endorse me for the corresponding arXiv category?

Of course I fully understand if you prefer not to do so.

Best regards,
Dirk Kunert

Checklist before contacting anyone

Before sending the manuscript to potential readers, check the following:

1. The paper compiles without errors.
2. The title names both the multiset and the set case.
3. The abstract states both period formulas and the inequality $\lambda_{\text{set}} \leq \lambda_{\text{multiset}}$.
4. The setup contains an explicit multiplicity convention and is clear that the set case discards multiplicities.
5. The bi-infinite nondecreasing enumeration of projected values is justified for both the multiset and the set side.
6. The proof of $\tilde{p}_s^{(i+N)} = \tilde{p}_s^{(i)} + D$ is explicit.
7. The degenerate case $D \mid N$ clearly explains the role of zero gaps.
8. The set-side dichotomy at $N = D$ (period N below, period 1 at and above) is stated as a separate theorem.
9. Table 1 lists the line-by-line correspondence between paper results and Lean declarations and the cited line numbers are correct.
10. The Lean formalisation is described precisely and not overstated: $\sim 2,850$ lines in `Basic.lean`, no `sorry`, `admit`, or `axiom`; both `main_theorem_concrete` (multiset) and `set_main_theorem_concrete` (set) are unconditional.
11. AI systems are not listed as co-authors.
12. AI assistance is mentioned only in the acknowledgements, with the author taking full responsibility for the mathematical content.
13. The bibliography is cleaned up, preferably replacing Wikipedia with a mathematical reference for the three-gap theorem.
14. The GitHub repository (`cut-and-project`) builds cleanly: `lake build` succeeds against the pinned toolchain (`leanprover/lean4:v4.29.1`, Mathlib commit `5e932f97...`) and `verify_paper.py` passes against the CSV data in `tests/`.

Practical recommendation

The best first step is to contact one or two people from the closest group, for example Alan Haynes and Henna Koivusalo, with a short email and the manuscript attached. If there is no response, proceed to James Walton, Valérie Berthé, or Sébastien Labbé.

The request should initially be for feedback on whether the results appear suitable for arXiv and whether either formula (multiset or set) is known in the literature, not immediately for endorsement. The set-versus-multiset dichotomy is the more distinctive contribution and is worth highlighting explicitly when summarising the paper.