The Hypercubic Manifold in Homotopy Type Theory

Dylan Laird

04/09/2023



Table of contents

- A brief overview of HoTT/cubical type theory
 - Informal Type Theory

Table of contents

- A brief overview of HoTT/cubical type theory
 - Informal Type Theory

We manipulate types A, B and terms a: A of some certain type.

We manipulate types A, B and terms a: A of some certain type.

The inductive type of natural integers

- 0 : N
- suc : $\mathbb{N} \to \mathbb{N}$

We manipulate types A, B and terms a: A of some certain type.

The inductive type of natural integers

- 0:N
- $suc : \mathbb{N} \to \mathbb{N}$

Equality

We distinguish between **definitional equality** ≡ used when defining objects and **propositional equality** which is a type theoretical concept.

We manipulate types A, B and terms a: A of some certain type.

The inductive type of natural integers

- 0:N
- suc : $\mathbb{N} \to \mathbb{N}$

Equality

We distinguish between **definitional equality** = used when defining objects and **propositional equality** which is a type theoretical concept.

Propositional Equality

Given a type A and terms a, b : A there is a type $a =_A b$. We say that elements a and b are (propositionally) equal when $a =_A b$ is inhabited.

Some constructions on types

Function types

Given types A, B one has a type $A \rightarrow B$ of functions from A to B.

Some constructions on types

Function types

Given types A, B one has a type $A \rightarrow B$ of functions from A to B.

Product type

Given types A, B one has a product type $A \times B$.

Some constructions on types

Function types

Given types A, B one has a type $A \rightarrow B$ of functions from A to B.

Product type

Given types A, B one has a product type $A \times B$.

Coproduct type

Give types A, B one has a coproduct type A + B given by the constructors :

- inl : $A \rightarrow A + B$
- inr : $B \rightarrow A + B$

Manipulating types

Introduction rules

They encapsulate how to build an element of a certain type.

- To build an element of $f: A \to B$ one needs an expression $\phi(x)$ such that $a: A \vdash \phi(a): A$ and to set $f: \equiv \lambda x.\phi(x)$
- To build an element of A × B one needs to take elements a: A and
 b: B to form (a, b): A × B

Manipulating types

Introduction rules

They encapsulate how to build an element of a certain type.

- To build an element of $f: A \to B$ one needs an expression $\phi(x)$ such that $a: A \vdash \phi(a): A$ and to set $f: \equiv \lambda x.\phi(x)$
- To build an element of A × B one needs to take elements a: A and
 b: B to form (a, b): A × B

Induction principles

They encapsulate how to build **dependent** functions from a source type A.

• To build a function of type $f: \prod_{x:A\times B} P(x)$ one only needs to give its value on pairs (a, b).

A working example : product types (1)

Projections

We can define projections $\operatorname{pr}_1: A \times B \to A$ and $\operatorname{pr}_2: A \times B \to B$ by the **induction** principle for product types by setting $\operatorname{pr}_1((a,b)) :\equiv a$ and $\operatorname{pr}_2((a,b)) :\equiv b$.

A working example : product types (1)

Projections

We can define projections $\operatorname{pr}_1: A \times B \to A$ and $\operatorname{pr}_2: A \times B \to B$ by the **induction** principle for product types by setting $\operatorname{pr}_1((a,b)) :\equiv a$ and $\operatorname{pr}_2((a,b)) :\equiv b$.

Propositional uniqueness for product types

We have a type theoretic statement that expresses that the type $A \times B$ is the type of "pairs of elements of A and B":

$$\operatorname{uniq}_{A \times B} : \prod_{x: A \times B} \left(x =_{A \times B} \left(\operatorname{pr}_{1}(x), \operatorname{pr}_{2}(x) \right) \right)$$

A working example : product types (2)

Sketch of proof:

• By induction, we only need to build an element :

$$\operatorname{uniq}_{A \times B}((a, b)) : (a, b) = (\operatorname{pr}_1(a, b), \operatorname{pr}_2(a, b))$$

A working example : product types (2)

Sketch of proof:

• By induction, we only need to build an element :

$$\mathrm{uniq}_{A\times B}((a,b)):(a,b)=(\mathrm{pr}_1(a,b),\mathrm{pr}_2(a,b))$$

By the definition of the projections, the goal type reduces to:

$$(a,b) =_{A \times B} (a,b)$$

A working example : product types (2)

Sketch of proof:

• By induction, we only need to build an element :

$$\mathrm{uniq}_{A\times B}((a,b)):(a,b)=(\mathrm{pr}_1(a,b),\mathrm{pr}_2(a,b))$$

By the definition of the projections, the goal type reduces to:

$$(a,b) =_{A \times B} (a,b)$$

• We now have an element refl : $(a, b) =_{A \times B} (a, b)$