


Mass Measurements


Lecture 1 – October 7, 2015

D. Lascar | Postdoctoral Fellow | TRIUMF

Precision v. Accuracy




Not Accurate
Not Precise



Accurate
Not Precise



Not Accurate
Precise



Accurate
Precise

Mass

- The property of a body to attract another body via gravitation
 - Gravitational mass
- Object's resistance to changing acceleration
 - Inertial mass
- The sum of all the rest energy in a body
 - Mass-energy relation

$$F = \frac{Gm_1m_2}{r^2}$$

$$\vec{F} = m\vec{a}$$

$$m = \left(Nm_n + Z(m_p + m_e) - \frac{1}{c^2} \sum_i BE_i \right)$$

Why measure masses?

- Molar mass

- $35.453 \pm 0.002 \text{ g/mol}$
 $35.453(2) \text{ g/mol}$

- $\frac{\delta m}{m} = \frac{0.002}{35.453} \approx 6 \times 10^{-5}$

- $N_A = 6.022140857(74) \times 10^{23} \text{ particles/mol}$


- $\frac{\delta m}{m} = \frac{0.000000074}{6.022140857} \approx 1.2 \times 10^{-8}$

- If you measure out 35.453 g *precisely*

- $6.02214(34) \times 10^{23} \text{ atoms}$

- **Does it matter?**

- **Depends on the application**



Why measure masses?


- Particle/molecular identification
 - What is the difference between:
 - ^{180}Hf
 - ^{179}HfH
 - $\text{C}_6\text{H}_{12}\text{O}_6$ – Glucose
 - $\text{C}_9\text{H}_8\text{O}_4$ – Caffeic Acid (Aspirin)
 - $\text{C}_{10}\text{H}_{12}\text{O}_3$ - Tyrosol, acetate
 - $\text{C}_7\text{H}_7\text{F}_3\text{O}_2$ – No idea but it exists in nature

Why measure masses?

	Mass (AMU)	$m - M(^{180}\text{Hf})$ (AMU)	$\frac{\Delta m}{M}$
^{180}Hf	179.9466	--	--
^{179}HfH	179.9538	0.0071	3.94×10^{-5}
$\text{C}_6\text{H}_{12}\text{O}_6$	180.0648	0.1168	6.49×10^{-4}
$\text{C}_9\text{H}_8\text{O}_4$	180.0432	0.0957	5.32×10^{-4}
$\text{C}_{10}\text{H}_{12}\text{O}_3$	180.0800	0.1321	7.34×10^{-4}
$\text{C}_7\text{H}_7\text{F}_3\text{O}_2$	180.0406	0.0933	5.18×10^{-4}

Atomic and nuclear masses

Masses determine the atomic and nuclear binding energies reflecting all forces in the atom/nucleus.



$$= N \cdot \text{[neutron]} + Z \cdot \text{[proton]} + Z \cdot \text{[electron]} - \text{binding energy}$$

$$M_{\text{Atom}} = N \cdot m_{\text{neutron}} + Z \cdot m_{\text{proton}} + Z \cdot m_{\text{electron}} - (B_{\text{atom}} + B_{\text{nucleus}})/c^2$$

Slide courtesy of K. Blaum
TRIUMF Summer Institute

$$\delta m/m < 10^{-10}$$


$$\delta m/m = 10^{-6} - 10^{-8}$$

Most of physics requires masses

- General physics and chemistry
- Nuclear structure physics
- Astrophysics
- Weak interaction studies
- Fundamental neutrino physics
- Fundamental symmetries
- Tests of QED

Nuclear Astrophysics

- Big Bang
 - ^1H , ^2H , ^3He , ^4He , ^6Li , ^7Li
- Cosmic Ray Spallation (interstellar medium)
 - Many elements, including most Li, Be, B
- Stellar Burning
 - Up to ^{56}Fe
- p Process (core-collapse supernovae)
 - Some heavy elements
- rp/vp Process (x-ray bursts)
- s Process (AGB stars)
 - Heavy elements up to $A=208$ (half of all heavy nuclei)
- r Process (??)
 - Heavy elements $A>56$ (the other half)




Neutron separation energy and mass

$$S_n = M(A, Z) + M(n) - M(A + 1, Z)$$


The Astrophysical *R*-Process

- Temperatures are very high ($T \sim 10^9 \text{K}$)
- Neutron densities are very high ($n_n > 10^{20} \text{cm}^{-3}$)
- (n, γ) rate is high ($< 1 \mu\text{s/capture}$)



The Astrophysical *R*-Process

- Temperatures are very high ($T \sim 10^9 \text{K}$)
- Neutron densities are very high ($n_n > 10^{20} \text{cm}^{-3}$)
- (n, γ) rate is high ($< 1 \mu\text{s/capture}$)
- As region of low neutron separation energy (S_n) reached $(n, \gamma) \rightleftharpoons (\gamma, n)$ equilibrium achieved



[A. Champagne, RIA Summer School '06]

The Astrophysical *R*-Process

- Temperatures are very high ($T \sim 10^9 \text{K}$)
- Neutron densities are very high ($n_n > 10^{20} \text{cm}^{-3}$)
- (n, γ) rate is high ($< 1 \text{ }\mu\text{s/capture}$)
- As region of low neutron separation energy (S_n) reached $(n, \gamma) \leftrightarrow (\gamma, n)$ equilibrium achieved
- Equilibrium point defined by Saha Equation

Partition Function – Ratio can vary between $10^{\pm 4}$ but normally between $10^{\pm 1}$

$$\frac{Y(Z, A+1)}{Y(Z, A)} = n_n \left(\frac{2\pi\hbar^2}{m_u kT} \right)^{3/2} \underbrace{\left(\frac{A+1}{A} \right)^{3/2}}_{\approx 1} \frac{G^*(Z, A+1)}{2G^*(Z, A)} \exp \left[\frac{S_n(Z, A+1)}{kT} \right]$$

The Astrophysical *R*-Process

- Temperatures are very high ($T \sim 10^9 \text{K}$)
- Neutron densities are very high ($n_n > 10^{20} \text{cm}^{-3}$)
- (n, γ) rate is high ($< 1 \text{ } \mu\text{s/capture}$)
- As region of low neutron separation energy (S_n) reached $(n, \gamma) \leftrightarrow (\gamma, n)$ equilibrium achieved
- Equilibrium point defined by Saha Equation


$$S_n^0 \approx kT \ln \left[\frac{2}{n_n} \left(\frac{m_u kT}{2\pi\hbar^2} \right)^{3/2} \right]$$

The Astrophysical *R*-Process

- Temperatures are very high ($T \sim 10^9 \text{K}$)
- Neutron densities are very high ($n_n > 10^{20} \text{cm}^{-3}$)
- (n, γ) rate is high ($< 1 \text{ } \mu\text{s/capture}$)
- As region of low neutron separation energy (S_n) reached $(n, \gamma) \leftrightarrow (\gamma, n)$ equilibrium achieved
- Equilibrium point defined by Saha Equation

$$S_n^0 \approx kT \ln \left[\frac{2}{n_n} \left(\frac{m_u kT}{2\pi\hbar^2} \right)^{3/2} \right] \approx 3 \text{ MeV}$$

So we wait for a β -decay




[A. Champagne, RIA Summer School 06]

The S_n where equilibrium occurs depends only on T and

$$S_n^0 \approx kT \ln \left[\frac{2}{n_n} \left(\frac{m_u kT}{2\pi\hbar^2} \right)^{3/2} \right]$$

n_n

Oct 7,
2015




Nucleosynthesis in the r-process


Joint Institute for Nuclear Astrophysics 2012

Time: 4.1×10^{-1} s

Temperature: 0.25 GK



Evidence From Elemental Abundances




A	N
195	126
130	82

Y.-Z. Qian, *Prog. Part. Nucl. Phys.*, 50 (2003) 153

Neutron separation energy and mass

$$S_n = M(A, Z) + M(n) - M(A + 1, Z)$$

Test of nuclear mass models...(Hint: They stink!)



How to measure masses? $\vec{B} \otimes$


Dipole mass spectrometry

- Charged particle in a uniform magnetic field
- $\vec{F} = q\vec{v} \times \vec{B}$ (Magnetic force)
- $\vec{F} \perp \vec{v}$ (F is a centripetal force)

$$qvB = \frac{mv^2}{\rho} \quad \left| \begin{array}{l} \text{Radius of} \\ \text{curvature} \end{array} \right.$$

- Magnetic Rigidity ($B\rho$):

$$B\rho = \frac{p}{q}$$
- Best $\frac{\delta m}{m} \sim 10^{-5}$




How to measure masses?

Time of flight mass spectrometry

- Measure the time that an ion at known energy will travel a known distance.
- Can be linear
 - Space limitation
- Storage rings better
 - Distance unlimited
- Best precision:

$$\frac{\delta m}{m} \sim 10^{-6}$$



We can do better than $\delta m/m \sim 10^{-6}$

- Actually we need
 - For nuclear astrophysics $\frac{\delta m}{m} \sim 10^{-7}$ *minimum*
 - For fundamental symmetries: $\frac{\delta m}{m} \sim 10^{-10}$ *minimum*
- What is the physical quantity that we can measure to the highest precision?

Time

- Time is the physical quantity we can measure most precisely
- For this clock:
- We can do better than $\frac{\delta t}{t} \sim 10^{-15}$
- When making a precision measurement, the goal is to measure time (frequency)

WIRED

ADAM MANN SCIENCE 04.04.14 6:30 AM

HOW THE U.S. BUILT THE WORLD'S MOST RIDICULOUSLY ACCURATE ATOMIC CLOCK

$$\frac{\delta t}{t} \sim 10^{-14}$$

For next class:

- How can you relate a time (frequency) measurement to mass?

Thank you!

Merci



Canada's national laboratory for particle and nuclear physics

Laboratoire national canadien pour la recherche en physique nucléaire et en physique des particules

TRIUMF: Alberta | British Columbia |
Calgary | Carleton | Guelph | Manitoba |
McGill | McMaster | Montréal | Northern
British Columbia | Queen's | Regina |
Saint Mary's | Simon Fraser | Toronto |
Victoria | Western | Winnipeg | York

