

통계수학1 과제#3 풀이

*연습문제 1.3 - 문제 6(d) (역함수만 구할 것, $x = f^{-1}(y)$ 꼴로 역함수를 표시할 것),
문제 11

$$6(d) \quad f(x) = y = \sqrt{4-x^2}, \quad 0 \leq x \leq 2$$

$$x = f^{-1}(y) = \sqrt{4-y^2}, \quad 0 \leq y \leq 2$$

$$11 \quad f(x) = \frac{1}{8}x - 3 \text{ and } g(x) = x^3 \Rightarrow f^{-1}(x) = 8(x+3) \text{ and } g^{-1}(x) = \sqrt[3]{x}$$

$$(a) \quad (f^{-1} \circ g^{-1})(1) = f^{-1}(g^{-1}(1)) = f^{-1}(1) = 32$$

$$(b) \quad (f^{-1} \circ f^{-1})(6) = f^{-1}(f^{-1}(6)) = f^{-1}(72) = 600$$

*연습문제 1.4 - 문제 15 (f의 역함수만 구할 것, $x = f^{-1}(y)$ 꼴로 역함수를 표시할 것)

답:

$$y = \ln(x + \sqrt{x^2 + 1})$$

$$e^y = x + \sqrt{x^2 + 1}$$

$$(e^y - x)^2 = x^2 + 1$$

$$e^{2y} - 2xe^y + x^2 = x^2 + 1$$

$$2xe^y = e^{2y} - 1$$

$$x = \frac{e^{2y} - 1}{2e^y}$$

$$\therefore f^{-1}(y) = \frac{e^{2y} - 1}{2e^y}$$

*연습문제 1.6 - 문제 13

답: $\lim_{x \rightarrow 0} \frac{\sin x (1 - \cos x)}{2x^2} = \lim_{x \rightarrow 0} \left[\frac{1}{2} \cdot \frac{\sin x}{x} \cdot \frac{1 - \cos x}{x} \right] = \frac{1}{2}(1)(0) = 0$

*연습문제 1.7 - 문제 24 ($x = 0$ 에서 연속이 되기 위한 $f(0)$ 의 값만 제시할 것)

답:

$$f(x) = \frac{\sqrt{x+c^2} - c}{x}, \quad c > 0$$

$$\text{Domain: } x + c^2 \geq 0 \Rightarrow x \geq -c^2 \text{ and } x \neq 0, \quad [-c^2, 0) \cup (0, \infty)$$

$$\lim_{x \rightarrow 0} \frac{\sqrt{x+c^2} - c}{x} = \lim_{x \rightarrow 0} \frac{\sqrt{x+c^2} - c}{x} \cdot \frac{\sqrt{x+c^2} + c}{\sqrt{x+c^2} + c}$$

$$= \lim_{x \rightarrow 0} \frac{(x+c^2) - c^2}{x[\sqrt{x+c^2} + c]} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{x+c^2} + c} = \frac{1}{2c}$$

Define $f(0) = 1/(2c)$ to make f continuous at $x = 0$.

*연습문제 2.1 - 문제 14(b) (미분가능성만 판단하여 제시), 문제 15

14(b) $x=1$ 에서 미분가능

15 $x=2$ 에서 미분가능

$$f(x) = \begin{cases} (x-1)^3, & x \leq 1 \\ (x-1)^2, & x > 1 \end{cases}$$

The derivative from the left is

$$\begin{aligned} \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^-} \frac{(x-1)^3 - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^-} (x-1)^2 = 0. \end{aligned}$$

The derivative from the right is

$$\begin{aligned} \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} &= \lim_{x \rightarrow 1^+} \frac{(x-1)^2 - 0}{x - 1} \\ &= \lim_{x \rightarrow 1^+} (x-1) = 0. \end{aligned}$$

These one-sided limits are equal. Therefore, f is differentiable at $x = 1$. ($f'(1) = 0$)

Note that f is continuous at $x = 2$.

$$f(x) = \begin{cases} x^2 + 1, & x \leq 2 \\ 4x - 3, & x > 2 \end{cases}$$

The derivative from the left is

$$\begin{aligned} \lim_{x \rightarrow 2^-} \frac{f(x) - f(2)}{x - 2} &= \lim_{x \rightarrow 2^-} \frac{(x^2 + 1) - 5}{x - 2} \\ &= \lim_{x \rightarrow 2^-} (x + 2) = 4. \end{aligned}$$

The derivative from the right is

$$\lim_{x \rightarrow 2^+} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2^+} \frac{(4x - 3) - 5}{x - 2} = \lim_{x \rightarrow 2^+} 4 = 4.$$

The one-sided limits are equal. Therefore, f is differentiable at $x = 2$, ($f'(2) = 4$).

*연습문제 2.2 - 문제 3(b), 문제 18

3(b) $f(x) = \sqrt{x} - 6\sqrt[3]{x} = x^{1/2} - 6x^{1/3}$

$$f'(x) = \frac{1}{2}x^{-1/2} - 2x^{-2/3} = \frac{1}{2\sqrt{x}} - \frac{2}{x^{2/3}}$$

18 $y = 2x^2 - 3x + 1$

$$y = ax^2 + bx + c$$

Since the parabola passes through (0, 1) and (1, 0), we have:

$$(0, 1): 1 = a(0)^2 + b(0) + c \Rightarrow c = 1$$

$$(1, 0): 0 = a(1)^2 + b(1) + 1 \Rightarrow b = -a - 1$$

Thus, $y = ax^2 + (-a - 1)x + 1$. From the tangent line $y = x - 1$, we know that the derivative is 1 at the point (1, 0).

$$y' = 2ax + (-a - 1)$$

$$1 = 2a(1) + (-a - 1)$$

$$1 = a - 1$$

$$a = 2$$

$$b = -a - 1 = -3 \quad \text{Therefore, } y = 2x^2 - 3x + 1.$$

*연습문제 2.3 - 문제 3(c), 문제 9

답:

3(c) $f'(x) = \frac{2x-5}{2x\sqrt{x}}$

$$f(x) = \frac{2x+5}{\sqrt{x}} = 2x^{1/2} + 5x^{-1/2}$$

$$f'(x) = x^{-1/2} - \frac{5}{2}x^{-3/2} = x^{-3/2} \left[x - \frac{5}{2} \right]$$

$$= \frac{2x-5}{2x\sqrt{x}} = \frac{2x-5}{2x^{3/2}}$$

9 $y = -\frac{1}{2}x - \frac{1}{2}, y = -\frac{1}{2}x + \frac{7}{2}$

$$f(x) = \frac{x+1}{x-1}$$

$$f'(x) = \frac{(x-1) - (x+1)}{(x-1)^2} = \frac{-2}{(x-1)^2}$$

$$2y + x = 6 \Rightarrow y = -\frac{1}{2}x + 3; \text{ Slope: } -\frac{1}{2}$$

$$\frac{-2}{(x-1)^2} = -\frac{1}{2}$$


$$(x-1)^2 = 4$$

$$x-1 = \pm 2$$

$$x = -1, 3; f(-1) = 0, f(3) = 2$$

$$y - 0 = -\frac{1}{2}(x+1) \Rightarrow y = -\frac{1}{2}x - \frac{1}{2}$$

$$y - 2 = -\frac{1}{2}(x-3) \Rightarrow y = -\frac{1}{2}x + \frac{7}{2}$$



***연습문제 2.4 - 문제 14(b)**

답: $f''(0) = 0$

$$f(x) = \cos(x^2), \quad (0, 1)$$

$$f'(x) = -\sin(x^2)(2x) = -2x \sin(x^2)$$

$$f''(x) = -2x \cos(x^2)(2x) - 2 \sin(x^2)$$

$$= -4x^2 \cos(x^2) - 2 \sin(x^2)$$

$$f''(0) = 0$$

***연습문제 2.5 - 문제 5(b) (주어진 점에서 접선을 구할 것)**

답: $y' = 0$

$$(x^2 + y^2)^2 = 4x^2y$$

$$2(x^2 + y^2)(2x + 2yy') = 4x^2y' + y(8x)$$

$$4x^3 + 4x^2yy' + 4xy^2 + 4y^3y' = 4x^2y' + 8xy$$

$$4x^2yy' + 4y^3y' - 4x^2y' = 8xy - 4xy^2$$

$$4y'(x^2y + y^3 - x^2) = 4(2xy - x^3 - xy^2)$$

$$y' = \frac{2xy - x^3 - xy^2}{x^2y + y^3 - x^2}$$

At (1, 1): $y' = 0$

*연습문제 2.6 - 문제 1, 문제 6

답:

1(a) $\frac{1}{5}$

$$f(x) = x^3 + 2x - 1, f(1) = 2 = a$$

$$f'(x) = 3x^2 + 2$$

$$(f^{-1})'(2) = \frac{1}{f'(f^{-1}(2))} = \frac{1}{f'(1)} = \frac{1}{3(1)^2 + 2} = \frac{1}{5}$$

(b) $\frac{2\sqrt{3}}{3}$

$$f(x) = \sin x, f\left(\frac{\pi}{6}\right) = \frac{1}{2} = a$$

$$f'(x) = \cos x$$

$$(f^{-1})'\left(\frac{1}{2}\right) = \frac{1}{f'(f^{-1}(1/2))} = \frac{1}{f'(\pi/6)} = \frac{1}{\cos(\pi/6)} = \frac{1}{\sqrt{3}/2} = \frac{2\sqrt{3}}{3}$$

6 $y = \frac{4\sqrt{3}}{3}x + \frac{\pi}{3} - \frac{2\sqrt{3}}{3}$

$$y = 2 \arcsin x, \left(\frac{1}{2}, \frac{\pi}{3}\right)$$

$$y' = \frac{2}{\sqrt{1-x^2}}$$

$$\text{At } \left(\frac{1}{2}, \frac{\pi}{3}\right), y' = \frac{2}{\sqrt{1-(1/4)}} = \frac{4}{\sqrt{3}}$$

$$\text{Tangent line: } y - \frac{\pi}{3} = \frac{4}{\sqrt{3}}\left(x - \frac{1}{2}\right)$$

$$y = \frac{4}{\sqrt{3}}x + \frac{\pi}{3} - \frac{2}{\sqrt{3}}$$

$$y = \frac{4\sqrt{3}}{3}x + \frac{\pi}{3} - \frac{2\sqrt{3}}{3}$$

*연습문제 3.1 - 문제 4(c)

답: Max (-1,5) Min (0,0)

$$y = 3x^{2/3} - 2x, [-1, 1]$$

$$y' = 2x^{-1/3} - 2 = \frac{2(1 - \sqrt[3]{x})}{\sqrt[3]{x}}$$

Left endpoint: (-1, 5) Maximum

Critical number: (0, 0) Minimum

Right endpoint: (1, 1)