

# Priorities to Pairwise

Bill Adams

Aug 2017

## Abstract

Given a priority vector  $\vec{p}$  there is a method to create a pairwise comparison matrix  $M_{\vec{p}}$  that has  $\vec{p}$  as its priority vector. In this paper we provide context for why this calculation could be useful, the definition of the calculation itself, and a few example calculations.

## 1 Introduction

Given  $n$  items whose  $n \times n$  pairwise comparison matrix is  $M$ , standard AHP theory uses the largest eigenvector calculation to find the priority vector  $\vec{p}_M$  for those  $n$  items. In addition, standard AHP theory provides a way to work in reverse; i.e. given a priority vector  $\vec{p}$  we can create a pairwise comparison matrix  $M_{\vec{p}}$  whose largest eigenvector is  $\vec{p}$ .

This method of working in reverse can be useful in the following circumstance. If one knows the priority vector for a group of users, but the software that gives that priority does not provide the group pairwise comparison matrix, one can reconstruct a pairwise comparison matrix for the group that results in the given priority vector. This circumstance is what prompted the writing of this paper.

## 2 Calculation and proof

Given a priority vector  $\vec{p} = (p_1, \dots, p_n)$ , we construct the matrix  $M$  whose priority vector is  $\vec{p}$  by taking the matrix made of the ratios of the  $p_i$ 's. We state this result as the following lemma.

**Lemma 1.** *Let  $\vec{p} = (p_1, \dots, p_n)$  be the priority vector of  $n$  items. If we define the  $n \times n$  matrix  $M_{\vec{p}} = (m_{ij})$  by the formula*

$$m_{ij} = \frac{p_i}{p_j}$$

*then the priority vector for  $M_{\vec{p}}$  is  $\vec{p}$ .*

*Proof.* Let largest eigenvector of  $M_{\vec{p}}$  can be calculated from the following formula

$$\lim_{n \rightarrow \infty} \frac{b^n}{\sum_{i=1}^n b_i^n}$$

where  $b^i$  is an  $n$  dimensional vector defined using the following recursive formula:

$$\begin{aligned} b^0 &= (1, \dots, 1) \\ b^{i+1} &= M_{\vec{p}} \cdot b^i \end{aligned}$$

Thus we need only understand  $b^i$ . Let us begin by calculating  $b^1$ .

$$\begin{aligned} b^1 &= M_{\vec{p}} \cdot b^0 \\ &= \begin{bmatrix} 1 & p_1/p_2 & \dots & p_1/p_n \\ p_2/p_1 & 1 & \dots & p_2/p_n \\ \vdots & \vdots & \ddots & \vdots \\ p_n/p_1 & p_n/p_2 & \dots & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} \sum_{i=1}^n p_1/p_i \\ \sum_{i=1}^n p_2/p_i \\ \dots \\ \sum_{i=1}^n p_n/p_i \end{bmatrix} \\ &= \begin{bmatrix} p_1 \sum_{i=1}^n 1/p_i \\ p_2 \sum_{i=1}^n 1/p_i \\ \vdots \\ p_n \sum_{i=1}^n 1/p_i \end{bmatrix} \\ &= \sum_{i=1}^n 1/p_i \begin{bmatrix} p_1 \\ p_2 \\ \dots \\ p_n \end{bmatrix} \end{aligned}$$

Since we are dividing by the sum of the elements of  $b^i$  in the limit, we can multiply or divide  $b^i$  by any constant and not change the limit. Thus effectively  $b^1 = (p_1, \dots, p_n)$ .

Next we claim that  $b^2 = (p_1, \dots, p_n)$ . If that is the case, then we know that all of the  $b^i$ 's for  $i \geq 1$  are that as well, and we will have shown that the largest eigenvector of  $M_{\vec{p}}$  is  $\vec{p}$ . Let us calculate the value of  $b^2$  to complete the proof.

$$\begin{aligned}
b^2 &= M_{\vec{p}} \cdot b^1 \\
&= \begin{bmatrix} 1 & p_1/p_2 & \dots & p_1/p_n \\ p_2/p_1 & 1 & \dots & p_2/p_n \\ \vdots & \vdots & \ddots & \vdots \\ p_n/p_1 & p_n/p_2 & \dots & 1 \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \\ \vdots \\ p_n \end{bmatrix} \\
&= \begin{bmatrix} np_1 \\ np_2 \\ \vdots \\ np_n \end{bmatrix}
\end{aligned}$$

Again, since we are dividing by the sum of the elements of  $b^2$  in the limit, we can multiply or divide by any constant and not effect the limit calculation. Thus, effectively,  $b^2$  is effectively  $(p_1, \dots, p_n)$ .  $\square$

### 3 Examples

We calculate the matrix  $M_{\vec{p}}$  for several priority vectors  $\vec{p}$  below.

1. If  $\vec{p} = (1/2, 1/3, 1/6)$ , the ratio matrix  $M_{\vec{p}}$  is:

$$\begin{aligned}
M_{\vec{p}} &= \begin{bmatrix} 1 & \frac{1/2}{1/3} & \frac{1/2}{1/6} \\ \frac{1/3}{1/2} & 1 & \frac{1/3}{1/6} \\ \frac{1/6}{1/2} & \frac{1/6}{1/3} & 1 \end{bmatrix} \\
&= \begin{bmatrix} 1 & 3/2 & 3 \\ 2/3 & 1 & 2 \\ 1/3 & 1/2 & 1 \end{bmatrix}
\end{aligned}$$

2. If  $\vec{p} = (3, 2, 1, 4)$  we can calculate the ratio matrix as below. Notice that the priority vector need not be normalized (i.e. its components need not sum to 1).

$$\begin{aligned}
M_{\vec{p}} &= \begin{bmatrix} 1 & \frac{3}{2} & \frac{3}{1} & \frac{3}{4} \\ \frac{2}{3} & 1 & \frac{2}{1} & \frac{2}{4} \\ \frac{1}{3} & \frac{1}{2} & 1 & \frac{1}{4} \\ \frac{4}{3} & \frac{4}{2} & \frac{4}{1} & 1 \end{bmatrix} \\
&= \begin{bmatrix} 1 & \frac{3}{2} & 3 & \frac{3}{4} \\ \frac{2}{3} & 1 & 2 & \frac{1}{2} \\ \frac{1}{3} & \frac{1}{2} & 1 & \frac{1}{4} \\ \frac{4}{3} & 2 & 4 & 1 \end{bmatrix}
\end{aligned}$$