

A CLAS Proposal for PAC44

Transition Form Factors of the η' and ϕ Mesons with CLAS12

M. C. Kunkel^{*†1}, M. J. Amarian², D. Lersch¹, J. Ritman^{†1}, S.
Schadmand^{†1}, X. Song¹

¹Forschungszentrum Jülich, Jülich (Germany)

²Old Dominion University (U.S.A.)

Abstract

Dalitz decays are radiative decays in which the photon is virtual and subsequently produces an electron positron pair, $P \rightarrow l^+l^-X$. Such decays serve as an important tool used to reveal the internal structure of hadrons and the interaction mechanisms between photons and hadrons. Furthermore, assuming point-like particles, the electromagnetic interaction is calculable within QED by the Kroll-Wada formula. Transition form factors quantify deviations from the QED decay rate. They characterize modifications of the point-like photon-meson vertex due to the structure of the meson. For the η' meson this deviation represents the internal structure of the meson, while for the ϕ meson the deviation represents the transition from $\phi \rightarrow \eta$. The transition form factor can be characterized as $|F(q^2)|$, where q^2 is the square of the invariant mass of the lepton pair, and can be determined by comparing QED predictions to the experimentally measured rate.

Measurements with the highest scientific impact on the determination of the transition form factor have been performed in the space-like region ($q^2 < 0$) in collider experiments. However, due to experimental limitations (e.g. π^\pm contamination in lepton sample, low branching fractions, external conversion contamination), transition form factors in the time-like region ($q^2 > 0$) have not yet been precisely determined. Recent measurements of the time-like transition form factor for $\eta' \rightarrow e^+e^-\gamma$ have been performed by the BESIII collaboration with insufficient statistical precision, therefore the proper theoretical description cannot be determined.

From previous CLAS analyses using the g12 data set, it was preliminarily shown that measurements of the time-like transition form factor were achievable, but without the statistical precision needed to be competitive. Therefore, we propose to use CLAS12 to focus

^{*}Contact person, email: m.kunkel@fz-juelich.de

[†]Spokesperson

on the dilepton decay channels from the reactions $ep \rightarrow e'p\eta'$ and $ep \rightarrow e'p\phi$, where $\eta' \rightarrow e^+e^-\gamma$ and $\phi \rightarrow \eta e^+e^-$. The CLAS12 detector will be used to identify and measure the e^+e^- decay products by means of the High Threshold Cherenkov Counter (HTCC), Pre-Calorimeter (PCAL) and Electromagnetic Calorimeter (EC). The combination of HTCC+PCAL+EC can provide a rejection factor for single e^\pm/π^\pm of up to 10^6 for momenta less than 4.9 GeV/c with $\approx 100\%$ efficiency. For dileptons (e^+e^- pairs), this rejection factor will be $\approx 10^{12}$, which enables dilepton studies for branching ratios $\approx 10^{-9}$. Precise determination of momenta and angles of the e^+e^- decay products are the key features available to CLAS12. The momentum and angle of final state photons will be determined in CLAS12 by using the PCAL and EC. Consequently, the photon in the process $\eta' \rightarrow e^+e^-\gamma$ and the photons in the process $\phi \rightarrow e^+e^-\eta \rightarrow e^+e^-\gamma\gamma$ will be detected. Preliminary studies using the CLAS12 simulation suite have shown that a beam time of 100 days, at full luminosity, will accumulate a data sample at least one order of magnitude larger in statistics than the most current $\eta' \rightarrow e^+e^-\gamma$ and $\phi \rightarrow \eta e^+e^-$ measurement.

1 Introduction

In the year 1951, Richard Dalitz published a letter [1] in which he calculated the rate for the π^0 decaying into an electron-positron pair (dilepton) and a photon, $\pi^0 \rightarrow e^+e^-\gamma$. The calculation assumed that the decay proceeded through a two-photon decay in which one of the photons was virtual and converted internally into an electron-positron pair. This kind of reaction is now known as a Dalitz decay. The experimental evidence of this decay process was first observed in emulsion plates exposed to the Chicago cyclotron in 1952 [2] and a number of experiments performed over the next ten years verified Dalitz's hypothesis that the $\pi^0 \rightarrow e^+e^-\gamma$ decay resulted from internal conversion of a virtual photon [3, 4, 5]. A few years later N. Kroll and W. Wada calculated the framework for Dalitz decays within the QED framework [6], and extended the framework to double Dalitz Decays, in which the π^0 decays into two electron-positron pairs via emission of two virtual photons. Throughout the following years, much work was done to extend the framework of Dalitz decays to heavier mesons, such as η , ω , η' , and ϕ . With numerous experimental data taken, it was shown that the shape of the dilepton mass spectrum deviated from the QED predictions. Such deviations are attributed to the meson not being point-like, as calculated in QED, but instead to the internal structure of the meson. The virtual photon, that decayed into a dilepton pair, has the ability to probe the structure of meson because, like its on-shell counterpart, emission of a virtual photon is radiation, which decouples from any strong interaction within the meson when the meson transitions into its decay. Therefore, the information of the transition is encoded into the virtual photon, known as the Transition Form

Factor (TFF), and can be characterized as $|F(q^2)|$, where q^2 is the square of the invariant mass of the lepton pair. The transition form factor can be determined by comparing QED predictions to the experimentally measured rate.

In this proposal we present an experiment to study two channels of which decay via Dalitz decays, $\eta \rightarrow e^+e^-\gamma$ and $\phi \rightarrow e^+e^-\eta$. The η and ϕ are produced via electro-production, $ep \rightarrow ep\eta'$ and $ep \rightarrow ep\phi$ in Hall B, using the CLAS12 detector. The superior $e^+e^-/\pi^+\pi^-$ discrimination of the CLAS12 detector will give access to measurements for which $e^+e^-/\pi^+\pi^-$ branching ratios of 10^{12} is achievable. This proposal is organized as follows. In Section 2 we summarize the current knowledge of Dalitz decays and transition form factors, challenges in dilepton signal quality and how the CLAS12 detector can surpass the current challenges in measuring a TFF of low statistical error. In Section 3, an explanation of the kinematics of the decay processes will be given as well as kinematics of main competing backgrounds. In Section 4 a brief discussion on past CLAS analysis will be given, along with a description of analysis techniques that have been used and will be used in a CLAS12 measurement. Also in Section 4, an explanation of the Monte-Carlo simulations that were performed to extract the acceptances will be given as well as a calculation of expected yield and a validity check on the expected yield from previous CLAS analyses. In Section 5 we present the beam time request and a summary of the experiment.

2 Motivation

Here I will briefly write about the motivation to use dalitz decays to study the structure of meson. I will include the g-2 measurements Previous results etc

3 Kinematics

Write the kinematics of dalitz decays, and also of gamma gamma decays. Illustrate how the gamma can externally convert into a e^+e^- pair and discuss how the use of the SVT, with 50-65 μ m can this overcome background since a resolution of 1mm vertex resolution is needed.

3.1 η' Decays

The main decays studied for this analysis are when a pseudoscalar meson, $P_p(\eta')$, decays via 2 photons $\gamma\gamma$ or a photon γ and a dilepton (l^+l^-) and when a vector meson $V_p(\phi)$, decays via an η photons $\gamma\gamma$ or a photon γ and a dilepton (l^+l^-) Figure ?? illustrates the Feynman diagrams for the “Two photon decay” and the “Dalitz decay”.

4 Measurement

Here I will write about how we plan on using CLAS12

4.1 Previous CLAS6 analysis

4.2 Simulating

Write about using PLUTO++

4.3 Lepton Acceptance

Write about the acceptance

4.4 Calculating Expected Yield

4.4.1 Calculating Virtual Photon Flux

Expected yield

4.5 Realistic Yield

5 Summary

Give beam time request here.

References

- [1] R H Dalitz. On an alternative decay process for the neutral π^0 -meson. *Proceedings of the Physical Society. Section A*, 64(7):667, 1951.
- [2] J. J. Lord, Joseph Fainberg, D. M. Haskin, and Marcel Schein. Narrow angle pairs of particles from nuclear interactions. *Phys. Rev.*, 87:538–539, Aug 1952.
- [3] N. P. Samios. Dynamics of internally converted electron-positron pairs. *Phys. Rev.*, 121:275–281, Jan 1961.
- [4] P. Lindenfeld, A. Sachs, and J. Steinberger. The Internal Pair Production of γ -Rays of Mesonic Origin; Alternate Modes of π^0 Decay. *Physical Review*, 89:531–537, February 1953.
- [5] C. P. Sargent, R. Cornelius, M. Rinehart, L. M. Lederman, and K. Rogers. Diffusion cloud-chamber study of very slow mesons. i. internal pair formation. *Phys. Rev.*, 98:1349–1354, Jun 1955.
- [6] Norman M. Kroll and Walter Wada. Internal pair production associated with the emission of high-energy gamma rays. *Phys. Rev.*, 98:1355–1359, Jun 1955.