

DS 4400

Machine Learning and Data Mining I
Spring 2024

David M. Liu

Khoury College of Computer Science
Northeastern University

Tuesday January 16 2024

Today's Outline

- Probability review
 - Random variables (discrete)
 - Expectation and variance
 - Conditional probabilities and independence
 - Bayes Theorem
 - Marginalization
- **Linear algebra review**
 - Matrices
 - Vectors
 - Linear independence
 - Rank of a matrix and matrix inverse

Announcements

- Office hours schedule starting this week.
 - Please create a Khoury user account if you don't have one.
 - For this week only, we will use Zoom (link on Piazza)

Office Hours:

- **Jai:** Tuesdays 2:00 – 3:30pm
- **Dhanush:** Wednesdays 12:00 – 1:30pm
- **David:** Thursdays 2:00 – 3:30pm
- **Caleb:** Fridays 4:00 – 5:30pm

Office hours held virtually on the [Khoury Office Hours Application](#). If you do not have a Khoury account please apply for one [here](#).

- Homework 1 posted to Canvas. Due in 2 weeks on Gradescope. Two problems require coding.
- Please submit project teams by Friday January 26. There will be a forum on Piazza to help with searching for teammates.

COMPUTER SCIENCE


Unveiling intra-person fingerprint similarity via deep contrastive learning

Gabe Guo^{1*}, Aniv Ray¹, Miles Izydorczak², Judah Goldfeder¹, Hod Lipson³, Wenyao Xu⁴

Fingerprint biometrics are integral to digital authentication and forensic science. However, they are based on the unproven assumption that no two fingerprints, even from different fingers of the same person, are alike. This renders them useless in scenarios where the presented fingerprints are from different fingers than those on record. Contrary to this prevailing assumption, we show above 99.99% confidence that fingerprints from different fingers of the same person share very strong similarities. Using deep twin neural networks to extract fingerprint representation vectors, we find that these similarities hold across all pairs of fingers within the same person, even when controlling for spurious factors like sensor modality. We also find evidence that ridge orientation, especially near the fingerprint center, explains a substantial part of this similarity, whereas minutiae used in traditional methods are almost nonpredictive. Our experiments suggest that, in some situations, this relationship can increase forensic investigation efficiency by almost two orders of magnitude.

Copyright © 2024 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).

A



<https://www-science-org.ezproxy.neu.edu/content/article/do-prints-two-different-fingers-belong-same-person-ai-can-tell>

Probability review

Marginalization

- We can also show that:

$$P(B) = P(B \wedge [A = v_1 \vee A = v_2 \vee \dots \vee A = v_k])$$

$$P(B) = \sum_{i=1}^k P(B \wedge A = v_i) = \sum_{i=1}^k P(B | A = v_i) P(A = v_i)$$

- This is called **marginalization** over A

EXAMPLE

Marginalization

Consider this *joint* distribution $P(A \text{ and } B)$

		B		
		1	2	3
A	1	0.1	0.05	0.2
	2	0.15	0	0.1
	3	0	0.2	0.2

$$\begin{aligned}P(A = 2) &= P(A = 2 \cap B = 1) + P(A = 2 \cap B = 2) + P(A = 2 \cap B = 3) \\&= \sum_{i=1}^3 P(A = 2 \cap B = i) \\&= 0.15 + 0 + 0.1 \\&= 0.25\end{aligned}$$

Recap of Probability

- Discrete Random Variables
 - e.g. Bernoulli random variables
- Expectation and Variance

$$E[X] = \sum_v v \Pr[X = v] \quad \text{Var}[X] = E(X^2) - E^2(X)$$

- Conditional Probability and Bayes Rule

$$P(A \mid B) = \frac{P(B \mid A) \times P(A)}{P(B)}$$

- Marginalization

Probability Resources

- Review notes from Stanford's machine learning class
 - <http://cs229.stanford.edu/section/cs229-prob.pdf>
- David Blei's probability review
 - https://khoury.neu.edu/home/eelhami/courses/CS6140_Fall16/lecture0_review_probability_1.pdf
- Books:
 - Sheldon Ross, A First course in probability

Linear algebra review

Resources


- Zico Kolter, Linear algebra review
 - <http://cs229.stanford.edu/section/cs229-linalg.pdf>
- Books:
 - O. Bretscher, Linear Algebra with Applications

Vectors and matrices


- **Vector** in \mathbb{R}^n is an ordered set of n real numbers.

- e.g. $v = (1, 6, 3, 4)$ is in \mathbb{R}^4


- A column vector:


$$\begin{pmatrix} 1 \\ 6 \\ 3 \\ 4 \end{pmatrix}$$

- A row vector:


$$(1 \ 6 \ 3 \ 4)$$

- m -by- n **matrix** is an object in $\mathbb{R}^{m \times n}$ with m rows and n columns, each entry filled with a (typically) real number:


$$\begin{pmatrix} 1 & 2 & 8 \\ 4 & 78 & 6 \\ 9 & 3 & 2 \end{pmatrix}$$

Vector operations

- Addition component by component

$$[a_1, a_2, \dots, a_n] + [b_1, b_2, \dots, b_n] = [a_1 + b_1, \dots, a_n + b_n]$$

$$[1, -2, 5] + [0, 3, 7] =$$

- Subtraction is also done component by component

$$[a_1, a_2, \dots, a_n] - [b_1, b_2, \dots, b_n] = [a_1 - b_1, \dots, a_n - b_n]$$

– Can add and subtract row or column vectors of same dimension

- Dot product

– Only works for row and column vector of same size

$$[a_1, a_2, \dots, a_n] \cdot \begin{bmatrix} b_1 \\ \dots \\ b_n \end{bmatrix} = [a_1 b_1 + \dots + a_n b_n]$$

$$[1, -2, 5] \cdot \begin{bmatrix} 0 \\ 3 \\ 7 \end{bmatrix} =$$

Matrix Operations

Matrix multiplication

We will use upper case letters for matrices. The elements are referred by $A_{i,j}$.

- **Matrix product:**

$$A \in \mathbb{R}^{m \times n} \quad B \in \mathbb{R}^{n \times p}$$

$$C = AB \in \mathbb{R}^{m \times p}$$

$$C_{ij} = \sum_{k=1}^n A_{ik} B_{kj}$$

e.g.

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$$

$$AB = \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{pmatrix}$$

Matrix transpose

Transpose: You can think of it as

- “flipping” the rows and columns

OR

- “reflecting” vector/matrix on line

e.g. $\begin{pmatrix} a \\ b \end{pmatrix}^T = (a \quad b)$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^T = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$$

A is a **symmetric matrix** if $A = A^T$

Linear Independence

Linear independence

- A set of vectors is **linearly independent** if none of them can be written as a *linear combination* of the others.
 - Vectors x_1, \dots, x_k are linearly independent if $c_1x_1 + \dots + c_kx_k = 0$ implies $c_1 = \dots = c_k = 0$
- Otherwise they are **linearly dependent**

$$x_1 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \quad x_2 = \begin{pmatrix} 0 \\ 3 \\ 3 \end{pmatrix}$$

Linear independence

- A set of vectors is **linearly independent** if none of them can be written as a linear combination of the others.
- Vectors x_1, \dots, x_k are linearly independent if $c_1x_1 + \dots + c_kx_k = 0$ implies $c_1 = \dots = c_k = 0$
- Otherwise they are **linearly dependent**

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad x_2 = \begin{bmatrix} 4 \\ 1 \\ 5 \end{bmatrix} \quad x_3 = \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix}$$

**Whiteboard for Visual
Geometric Understanding**

Rank of a Matrix

- $\text{rank}(A)$ (the rank of a m -by- n matrix A) is
 - number of linearly independent columns
 - number of linearly independent rows

- If A is n by m , then
 - $\text{rank}(A) \leq \min(m, n)$

- Examples $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ $\begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix}$ $\begin{pmatrix} 2 & 1 & 3 \\ 0 & 5 & 2 \end{pmatrix}$

Matrix Inverse and System of Equations

Inverse of a matrix

- Inverse of a square matrix A , denoted by A^{-1} is the *unique* matrix s.t.
 - $AA^{-1} = A^{-1}A = I$ (identity matrix)
- Inverse of a square matrix exists only if the matrix is **full rank**
- If A^{-1} and B^{-1} exist, then
 - $(AB)^{-1} = B^{-1}A^{-1}$
 - $(A^T)^{-1} = (A^{-1})^T$

Diagonal matrices

A diagonal matrix is a square matrix with zeros everywhere except the diagonal.

$$D = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$D^T = D$$

D^{-1} : replace all entries of the diagonal with their reciprocal

$$D^{-1} = \begin{bmatrix} \frac{1}{3} & 0 & 0 \\ 0 & \frac{1}{9} & 0 \\ 0 & 0 & \frac{1}{2} \end{bmatrix}$$

System of linear equations

$$\begin{array}{rclcl} 4x_1 & - & 5x_2 & = & -13 \\ -2x_1 & + & 3x_2 & = & 9. \end{array}$$

Matrix formulation

$$Ax = b$$

$$A = \begin{bmatrix} 4 & -5 \\ -2 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} -13 \\ 9 \end{bmatrix}.$$

If A has an inverse, solution is $x = A^{-1}b$

Recap of Linear Algebra

- Vector and Matrix Operations
 - Add, multiply, transpose
- Linear Independence
- Rank
- Matrix Inverse and System of Equations
- Continuous Random Variables
 - PDFs and CDFs

Not covered in this lecture: determinants, eigenvalues/eigenvectors, matrix calculus

For more, see [Stanford CS 229 notes](#) pages 1-12

Announcements

- Office hours schedule starting this week.
 - Please create a Khoury user account if you don't have one.
 - For this week only, we will use Zoom (link on Piazza)

Office Hours:

- **Jai:** Tuesdays 2:00 – 3:30pm
- **Dhanush:** Wednesdays 12:00 – 1:30pm
- **David:** Thursdays 2:00 – 3:30pm
- **Caleb:** Fridays 4:00 – 5:30pm

Office hours held virtually on the [Khoury Office Hours Application](#). If you do not have a Khoury account please apply for one [here](#).

- Homework 1 posted to Canvas. Due in 2 weeks on Gradescope. Two problems require coding.
- Please submit project teams by Friday January 26. There will be a forum on Piazza to help with searching for teammates.