

Page_5e	Section	Problems
213	4.1	3,6,10,14
221	4.2	5,6,13,16,36, (40 from 6th edition)
229-230	4.3	5,12,22,24,41,45
240-241	4.4	4,7,10,45, 46,52
245	4.5	3,9,12
248-249	4.6	4,10,14
659	12.1	3,10,18, 21
664	12.2	10,12,16
669-670	12.3	4,6,12,30,33
692	13.1	3,8,24
697	13.2	3,10,12
699	13.3	2,5
703-704	13.4	3,5,10
709-710	13.5	4,10,13
716	13.6	1,6
758	15.3	2,10

Section 4.1 · ed. 5 p.213

In Problems 1–18, use Definition 4.1.1 to find $\mathcal{L}\{f(t)\}$.

$$3. f(t) = \begin{cases} t, & 0 \leq t < 1 \\ 1, & t \geq 1 \end{cases}$$

$$6. f(t) = \begin{cases} \sin t, & 0 \leq t < \pi/2 \\ 0, & t \geq \pi/2 \end{cases}$$




FIGURE 4.1.9 Graph for Problem 10

$$14. f(t) = t^2 e^{-2t}$$

Section 4.2 · ed. 5 p.221

In Problems 1–30, use Theorem 4.2.1 to find the given inverse

$$5. \mathcal{L}^{-1}\left\{\frac{(s+1)^3}{s^4}\right\}$$

$$6. \mathcal{L}^{-1}\left\{\frac{(s+2)^2}{s^3}\right\}$$

$$13. \mathcal{L}^{-1}\left\{\frac{4s}{4s^2+1}\right\}$$

$$16. \mathcal{L}^{-1}\left\{\frac{s+1}{s^2+2}\right\}$$

In Problems 31–40, use the Laplace transform to solve the given initial-value problem.

$$36. y'' - 4y' = 6e^{3t} - 3e^{-t}, \quad y(0) = 1, \quad y'(0) = -1$$

In Problems 33–44, use the Laplace transform to solve the given initial-value problem.

$$40. y'' + 9y = e^t, \quad y(0) = 0, \quad y'(0) = 0 \quad \text{from 6th ed.}$$

Section 4.3 · ed. 5 p.229, p.230

In Problems 1–20, find either $F(s)$ or $f(t)$, as indicated.

5. $\mathcal{L}\{t(e^t + e^{2t})^2\}$

12. $\mathcal{L}^{-1}\left\{\frac{1}{(s-1)^4}\right\}$

In Problems 21–30, use the Laplace transform to solve the given

22. $y' - y = 1 + te^t, \quad y(0) = 0$

24. $y'' - 4y' + 4y = t^3 e^{2t}, \quad y(0) = 0, \quad y'(0) = 0$

41. $\mathcal{L}\{\cos 2t \cdot \mathcal{U}(t - \pi)\}$

45. $\mathcal{L}^{-1}\left\{\frac{e^{-\pi s}}{s^2 + 1}\right\}$

Section 4.4 · ed. 5 p.240, p.241

In Problems 1–8, use Theorem 4.4.1 to evaluate the given Laplace transform.

4. $\mathcal{L}\{t \sinh 3t\}$

7. $\mathcal{L}\{te^{2t} \sin 6t\}$

8. $\mathcal{L}\{te^{-3t} \cos 3t\}$

In Problems 9–14, use the Laplace transform to solve the given

10. $y' - y = te^t \sin t, \quad y(0) = 0$

In Problems 37–46, use the Laplace transform to solve the given integral equation or integrodifferential equation.

45. $y'(t) = 1 - \sin t - \int_0^t y(\tau) d\tau, \quad y(0) = 0$

46. $\frac{dy}{dt} + 6y(t) + 9 \int_0^t y(\tau) d\tau = 1, \quad y(0) = 0$

In Problems 51–56, use Theorem 4.4.3 to find the Laplace transform of the given periodic function.




FIGURE 4.4.7 Graph for Problem 52

Section 4.5 · ed. 5 p.245

In Problems 1–12, use the Laplace transform to solve the given differential equation subject to the indicated initial conditions.

3. $y'' + y = \delta(t - 2\pi), \quad y(0) = 0, y'(0) = 1$

9. $y'' + 4y' + 5y = \delta(t - 2\pi), \quad y(0) = 0, y'(0) = 0$

12. $y'' - 7y' + 6y = e^t + \delta(t - 2) + \delta(t - 4), \quad y(0) = 0, y'(0) = 0$

Section 4.6 · ed. 5 p.248, p.249

In Problems 1–12, use the Laplace transform to solve the given system of differential equations.

4.
$$\begin{aligned} \frac{dx}{dt} + 3x + \frac{dy}{dt} &= 1 \\ \frac{dx}{dt} - x + \frac{dy}{dt} - y &= e^t \\ x(0) &= 0, \quad y(0) = 0 \end{aligned}$$

10.
$$\begin{aligned} \frac{dx}{dt} - 4x + \frac{d^3y}{dt^3} &= 6 \sin t \\ \frac{dx}{dt} + 2x - 2\frac{d^3y}{dt^3} &= 0 \\ x(0) &= 0, \quad y(0) = 0, \\ y'(0) &= 0, \quad y''(0) = 0 \end{aligned}$$

14. Derive the system of differential equations describing the straight-line vertical motion of the coupled springs shown in equilibrium in **FIGURE 4.6.4**. Use the Laplace transform to solve the system when $k_1 = 1$, $k_2 = 1$, $k_3 = 1$, $m_1 = 1$, $m_2 = 1$ and $x_1(0) = 0$, $x_1'(0) = -1$, $x_2(0) = 0$, $x_2'(0) = 1$.




FIGURE 4.6.4 Coupled springs in Problem 14

Section 12.1 · ed. 5 p.659 In Problems 1–6, show that the given functions are orthogonal on the indicated interval.

3. $f_1(x) = e^x$, $f_2(x) = xe^{-x} - e^{-x}$; $[0, 2]$
6. $f_1(x) = e^x$, $f_2(x) = \sin x$; $[\pi/4, 5\pi/4]$

In Problems 7–12, show that the given set of functions is orthogonal on the indicated interval. Find the norm of each function in the set.

10. $\left\{ \sin \frac{n\pi}{p}x \right\}$, $n = 1, 2, 3, \dots$; $[0, p]$
18. From Problem 1 we know that $f_1(x) = x$ and $f_2(x) = x^2$ are orthogonal on $[-2, 2]$. Find constants c_1 and c_2 such that $f_3(x) = x + c_1x^2 + c_2x^3$ is orthogonal to both f_1 and f_2 on the same interval.
21. A real-valued function f is said to be **periodic** with period

See later in document.

Section 12.2 · ed. 5 p.664

In Problems 1–16, find the Fourier series of f on the given

$$10. f(x) = \begin{cases} 0, & -\pi/2 < x < 0 \\ \cos x, & 0 \leq x < \pi/2 \end{cases}$$

$$12. f(x) = \begin{cases} 0, & -2 < x < 0 \\ x, & 0 \leq x < 1 \\ 1, & 1 \leq x < 2 \end{cases}$$

$$16. f(x) = \begin{cases} 0, & -\pi < x < 0 \\ e^x - 1, & 0 \leq x < \pi \end{cases}$$

Section 12.3 · ed. 5 p.669, p.670

In Problems 1–10, determine whether the function is even, odd

$$4. f(x) = x^3 - 4x$$

$$6. f(x) = e^x - e^{-x}$$

In Problems 11–24, expand the given function in an appropriate cosine or sine series.

$$12. f(x) = \begin{cases} 1, & -2 < x < -1 \\ 0, & -1 < x < 1 \\ 1, & 1 < x < 2 \end{cases}$$

In Problems 25–34, find the half-range cosine and sine expansions of the given function.

$$30. f(x) = \begin{cases} 0, & 0 < x < \pi \\ x - \pi, & \pi \leq x < 2\pi \end{cases}$$

$$33. f(x) = x^2 + x, \quad 0 < x < 1$$

Section 13.1 · ed. 5 p.692

In Problems 1–16, use separation of variables to find, if possible, product solutions for the given partial differential equation.

$$3. u_x + u_y = u$$

$$8. y \frac{\partial^2 u}{\partial x \partial y} + u = 0$$

In Problems 17–26, classify the given partial differential equation as hyperbolic, parabolic, or elliptic.

$$24. \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = u$$

$$25. a^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2}$$

$$26. k \frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}, \quad k > 0$$

Section 13.2 · ed. 5 p.697

In Problems 1–6, a rod of length L coincides with the interval $[0, L]$ on the x -axis. Set up the boundary-value problem for the temperature $u(x, t)$.

3. The left end is held at temperature 100° , and there is heat transfer from the right end into the surrounding medium at temperature zero. The initial temperature is $f(x)$ throughout.

In Problems 7–10, a string of length L coincides with the interval $[0, L]$ on the x -axis. Set up the boundary-value problem for the displacement $u(x, t)$.

10. The ends are secured to the x -axis, and the string is initially at rest on that axis. An external vertical force proportional to the horizontal distance from the left end acts on the string for $t > 0$.

From 12.1

21. A real-valued function f is said to be **periodic** with period T if $f(x + T) = f(x)$. For example, 4π is a period of $\sin x$ since $\sin(x + 4\pi) = \sin x$. The smallest value of T for which $f(x + T) = f(x)$ holds is called the **fundamental period** of f . For example, the fundamental period of $f(x) = \sin x$ is $T = 2\pi$. What is the fundamental period of each of the following functions?

(a) $f(x) = \cos 2\pi x$

(b) $f(x) = \sin \frac{4}{L} x$

(c) $f(x) = \sin x + \sin 2x$

(d) $f(x) = \sin 2x + \cos 4x$

(e) $f(x) = \sin 3x + \cos 2x$

(f) $f(x) = A_0 + \sum_{n=1}^{\infty} \left(A_n \cos \frac{n\pi}{p} x + B_n \sin \frac{n\pi}{p} x \right), A_n \text{ and } B_n$
depend only on n

12. A semi-infinite plate coincides with the region defined by $0 \leq x \leq \pi, y \geq 0$. The left end is held at temperature e^{-y} , and the right end is held at temperature 100° for $0 < y \leq 1$ and temperature zero for $y > 1$. The bottom of the plate is held at temperature $f(x)$.

Section 13.3 · ed. 5 p.699


In Problems 1 and 2, solve the heat equation (1) subject to the given conditions. Assume a rod of length L .

2. $u(0, t) = 0, \quad u(L, t) = 0$
 $u(x, 0) = x(L - x)$

5. Suppose heat is lost from the lateral surface of a thin rod of length L into a surrounding medium at temperature zero. If the linear law of heat transfer applies, then the heat equation takes on the form

$$k \frac{\partial^2 u}{\partial x^2} - hu = \frac{\partial u}{\partial t}, \quad 0 < x < L, \quad t > 0,$$

h a constant. Find the temperature $u(x, t)$ if the initial temperature is $f(x)$ throughout and the ends $x = 0$ and $x = L$ are insulated. See FIGURE 13.3.3.



Section 13.4 · ed. 5 p.703, p.704

In Problems 1–6, solve the wave equation (1) subject to the given conditions.


3. $u(0, t) = 0, \quad u(\pi, t) = 0, \quad t > 0$

$$u(x, 0) = 0, \quad \left. \frac{\partial u}{\partial t} \right|_{t=0} = \sin x, \quad 0 < x < \pi$$

5. $u(0, t) = 0, \quad u(1, t) = 0, \quad t > 0$

$$u(x, 0) = x(1 - x), \quad \left. \frac{\partial u}{\partial t} \right|_{t=0} = x(1 - x), \quad 0 < x < 1$$

10. $f(x)$



In Problems 7–10, a string is tied to the x -axis at $x = 0$ and at $x = L$ and its initial displacement $u(x, 0) = f(x)$, $0 < x < L$, is shown in the figure. Find $u(x, t)$ if the string is released from rest.

Section 13.5 · ed. 5 p.709. p.710

In Problems 1–10, solve Laplace's equation (1) for a rectangular plate subject to the given boundary conditions.

4. $\frac{\partial u}{\partial x} \Big|_{x=0} = 0, \quad \frac{\partial u}{\partial x} \Big|_{x=a} = 0$

$u(x, 0) = x, \quad u(x, b) = 0$

10. $u(0, y) = 10y, \quad \frac{\partial u}{\partial x} \Big|_{x=1} = -1$

$u(x, 0) = 0, \quad u(x, 1) = 0$

In Problems 13 and 14, solve Laplace's equation (1) for a rectangular plate subject to the given boundary conditions.

13. $u(0, y) = 0, \quad u(a, y) = 0$

$u(x, 0) = f(x), \quad u(x, b) = g(x)$

Section 13.6 · ed. 5 p.716

In Problems 1 and 2, solve the heat equation $ku_{xx} = u_t$, $0 < x < 1$, $t > 0$ subject to the given conditions.

1. $u(0, t) = 100, \quad u(1, t) = 100$

$u(x, 0) = 0$

6. Solve the boundary-value problem

$$k \frac{\partial^2 u}{\partial x^2} - hu = \frac{\partial u}{\partial t}, \quad 0 < x < \pi, \quad t > 0$$

$$u(0, t) = 0, \quad u(\pi, t) = u_0, \quad t > 0$$

$$u(x, 0) = 0, \quad 0 < x < \pi.$$

In Problems 1–6, find the Fourier integral representation of the given function.

$$\mathbf{2.} \quad f(x) = \begin{cases} 0, & x < \pi \\ 4, & \pi < x < 2\pi \\ 0, & x > 2\pi \end{cases} \quad \mathbf{3.} \quad f(x) = \begin{cases} 0, & x < 0 \\ x, & 0 < x < 3 \\ 0, & x > 3 \end{cases}$$

In Problems 7–12, represent the given function by an appropriate cosine or sine integral.

$$\mathbf{10.} \quad f(x) = \begin{cases} x, & |x| < \pi \\ 0, & |x| > \pi \end{cases}$$