



Tomography of the Solar Corona with Multiple Instruments: First Steps

D.G. Lloveras¹, A.M. Vásquez^{1,2}, Enrico Landi³, Richard A. Frazin³

¹ *Instituto de Astronomía y Física del Espacio, CONICET-UBA, Argentina*

² *Departamento de Ciencia y Tecnología, UNTREF, Argentina*

³ *Department of Climate and Space Sciences and Engineering, University of Michigan, USA*

Contact / dlloveras@iafe.uba.ar

Resumen / La tomografía solar rotacional es una técnica observacional de la corona solar que permite la reconstrucción de la distribución tri-dimensional (3D) global de algunos de sus parámetros fundamentales, como por ejemplo la densidad electrónica. La tomografía basada en EUV es aplicada regularmente a datos obtenidos con telescopios espaciales, cubriendo típicamente el rango de alturas heliocéntricas $1.02 - 1.25 R_{\odot}$. Este rango solapa parcialmente el del campo de visión del coronógrafo de luz blanca K-coronagraph (KCOR), en el High Altitude Observatory (HAO), que cubre el rango $1.05 - 3.0 R_{\odot}$. En este trabajo presentamos resultados preliminares de la primer comparación de una reconstrucción de la densidad electrónica coronal obtenida utilizando imágenes EUV con la obtenida utilizando imágenes en luz blanca. Los resultados obtenidos son discutidos en términos de las diversas características de los datos utilizados, así como de los diferentes factores físicos que afectan cada análisis.

Abstract / Solar rotational tomography is an observational technique of the solar corona that allows reconstruction of the global three-dimensional distribution of some of its fundamental physical parameters, such as electron density. EUV tomography is routinely applied to data provided by spaceborne telescopes, typically covering the range of heliocentric heights $\approx 1.02 - 1.25 R_{\odot}$. This range partially overlaps that of the field of view of the white light K-coronagraph (KCOR) instrument, at the High Altitude Observatory (HAO), which covers the range $\approx 1.05 - 3.0 R_{\odot}$. In this work we show preliminary results of the first comparison of a tomographic reconstruction of the coronal electron density based on EUV images against that reconstructed based on white light images. Results are discussed in terms of the diverse characteristics of the used data sets, as well as the different physical factors that affect each analysis.

Keywords / Sun: corona — Sun: fundamental parameters — Sun: UV radiation — Sun: abundances

1. Introduction

Solar rotational tomography (SRT) was initially developed by Altschuler & Perry (1972) for reconstruction of the three-dimensional (3D) distribution of the coronal electron density based on white light (WL) data. Frazin & Janzen (2002) developed a modern, robust, regularized, positive method for tomographic inversion of the coronal electron density from WL data, used in this work. A thorough review on WL tomography can be found in those references.

More recently, Frazin et al. (2009) developed the differential emission measure tomography (DEMT) technique. DEMT combines EUV tomography in several pass-bands with local DEM analysis, to reconstruct the 3D distribution of both the coronal electron density and temperature. A recent review by Vásquez (2016) summarizes the existing solar physics literature based on DEMT analysis. More recent DEMT-based works include Lloveras et al. (2017), who carried out a comparative analysis of the coronal structure at the solar minima between solar cycles (SC-)22/23 and 23/24.

DEMT research has been carried out based on data provided by several generations of spaceborne EUV telescopes. The latest one is the Atmospheric Imaging As-

sembly (AIA) instrument, on board the Solar Dynamics Observatory (SDO), whose data is used in this work.

While the electron density determined by WL tomography is of an absolute nature (Frazin et al., 2010), that obtained from EUV tomography is dependent on the assumed iron coronal abundance, as well as affected by the so-called coronal filling factor (Frazin et al., 2009). So far, no comparison had been carried out between results obtained with both techniques. Here, the tomographic reconstruction of the coronal electron density for a given period is carried out using both methods and their results are compared for the first time.

2. Data and Methodology

Carrington rotation (CR-)2198 (2017, December 03 UT 14:37 - December 30 UT 22:25) was selected as target for analysis, a relatively quiet rotation in the declining activity phase of SC-24. Low latitudes were dominated by the equatorial streamer belt, high latitudes by polar coronal holes (CHs), and a complex of active regions (ARs) was located in the longitude range $\approx 80^{\circ} - 200^{\circ}$.

For the selected rotation, KCOR and AIA images were obtained for the period 2017, December 03 through December 17, allowing to observe the off-limb corona at




Figure 1: Example of images used for tomographic reconstruction of the coronal electron density of CR-2198 (see text), both corresponding to 2017 December 03 UT 18:00-19:00. Left panel: SDO/AIA coronal EUV image in the 211 Å band. Right panel: HAO/KCOR coronal pB image, with white rings indicating heliocentric heights 1.09, 1.50, and 2.0 R_{\odot} .

all Carrington longitudes. In the case of AIA, images for all the coronal bands 171, 193, and 211 Å are obtained every 1 hr, and processed with our own tomography data pre-processing tools, which make use of the SolarSoft AIA software in its latest version. In particular, images are averaged over 6 hr long bins, so that a total of about 55 images are used in the end. In the case of KCOR we requested to the HAO team one 10-minute average coronal polarized brightness (pB) image for each day, obtained using its best observational window.

Figure 1 shows examples of the coronal images used for this work. The left panel shows an AIA coronal EUV image taken in its 211 Å band. The right panel shows a KCOR coronal pB WL image. Both images were taken nearly-simultaneously on 2017 December 03 UT 18:00-19:00. This is roughly the beginning of CR-2198, so the longitude of the disk center in these images is $\approx 0^\circ$, so the ARs are not seen and the images are dominated by the quiet sun streamer belt and the CHs.

Both the WL and EUV tomographic reconstructions were performed on the same resolution, with computational voxels having uniform radial size of 0.01 R_{\odot} and angular size of 2° in both latitude and longitude. Solutions were computed with a 3D regularization scheme.

3. Results

For the specific data sets of this rotation, the coronal electron density structure was reconstructed in the height range 1.02 – 1.20 R_{\odot} from EUV tomography, and 1.09 – 1.80 R_{\odot} from WL tomography. Figure 2 shows Carrington maps of both reconstructions at heliocentric height $r = 1.105 R_{\odot}$.

The white boxes in the maps indicate two regions selected for quantitative comparison. The region in the Southern hemisphere is a high-density quiet Sun region within the equatorial streamer belt during this rotation. The region in the Northern hemisphere is a lower density region at subpolar latitudes in the northern CH.

The quantitative comparison of the electron density determined from both tomographic analysis at each

computational voxel of the two selected regions is shown in Figure 3. For each region, the top panel shows the histogram of the ratio of electron density as determined from both tomographic analysis for all voxels within the common reconstructed coronal volume, i.e. in the range of heights 1.09 – 1.20 R_{\odot} . For each region, the bottom panel shows the average radial dependence of the electron density determined from both analysis.

4. Discussion and Future Work

The coronal electron density determined from WL tomography is systematically larger than that determined from EUV tomography. The characteristic (median) WL-to-EUV electron density ratio is ~ 2.0 in the quiet sun streamer belt region and ~ 1.6 in the subpolar region within the CH.

This difference may be partly explained by the fact that the KCOR data is affected by the sky brightness which was not subtracted in the specific data set used for this first work, as the procedures that are required for that were being revised by the HAO team at the moment of this analysis. Once applied a sky subtraction, the electron density as determined from WL tomography will systematically decrease. However, once corrected a difference is still expected to remain in the electron density as determined from the two analysis. In that case, systematic differences can be attributable too physical factors.

Firstly, the EUV emissivity determined at each voxel from tomographic analysis is proportional to the local mean squared electron density, i.e. $E_{\text{EUV}} \propto \langle N_e^2 \rangle = f \langle N_e \rangle^2$, where the *filling factor* is defined as $f \equiv \langle N_e^2 \rangle / \langle N_e \rangle^2$. As the WL emission is due to Thompson scattering, its voxel emissivity is proportional to the local average electron density, i.e. $E_{\text{WL}} \propto \langle N_e \rangle$. Then: $\langle N_e \rangle_{\text{WL}} / \langle N_e \rangle_{\text{EUV}} \propto \sqrt{f}$.

If differences in the results are solely attributed to the filling factor f , then $f \sim 4.0$ the quiet sun region of the streamer belt, and $f \sim 2.5$ in the subpolar region of the CH. Note that, as for any probability distribution,



Figure 2: Example of results of the tomographic reconstruction of the electron density of the solar corona for CR-2198. Carrington maps of the reconstructed electron density are shown at heliocentric height $r = 1.105 R_{\odot}$. Left panel: reconstruction based on EUV data. Right panel: reconstruction based on WL data. The white boxes indicate two ranges of longitudes and latitudes selected for quantitative comparison (see text).




Figure 3: Quantitative comparison between the results of the two tomographic reconstructions of the coronal electron density of CR-2198 in the two selected regions indicated in Figure 2. Left panels show the results in the quiet sun region of the southern hemisphere within the streamer belt, and right panels show the results in the subpolar open region within the northern CH. For each region, the top panel shows the histogram of the ratio of the electron density value obtained in each computational voxel from the WL and EUV tomographic analysis. For each region, the bottom panel shows the average radial profile of the electron density based on the EUV (red) and WL (blue) tomographic reconstructions.

$\sigma_{N_e}^2 \equiv \text{Var}N_e = \langle N_e^2 \rangle - \langle N_e \rangle^2 = \langle N_e \rangle^2 (f - 1)$, then $\sigma_{N_e} / \langle N_e \rangle = \sqrt{f - 1}$. With this interpretation, where f is larger (quiet sun region) the electron density probability distribution has larger variance.

This work is a first step towards implementation of a new technique dubbed multi-instrument tomography (MIT), currently under development. MIT will aim at simultaneously reconstructing the 3D distribution of different coronal parameters through joint analysis of tomographic results based on data provided by multiple instruments, including white-light coronagraphs, EUV telescopes and visible emission line coronagraphs. In particular, MIT will attempt to combine KCOR and AIA data, with that to be provided by the, soon to be operative, Upgraded Coronal Multichannel Polarimeter (UCoMP) instrument (Landi et al., 2016).

References

- Altschuler M. D., Perry R. M., 1972, SoPh, 23, 410
- Frazin R. A., Janzen P., 2002, ApJ, 570, 408
- Frazin R. A., et al., 2010, SoPh, 265, 19
- Frazin R. A., Vázquez A. M., Kamalabadi F., 2009, ApJ, 701, 547
- Landi E., Habbal S. R., Tomczyk S., 2016, Journal of Geophysical Research (Space Physics), 121, 8237
- Lloveras D. G., et al., 2017, SoPh, 292, 153
- Vázquez A. M., 2016, Advances in Space Research, 57, 1286