## Intermediate Test 2 – Solutions

## Stellenbosch Camp 2018

1. For any real number x, let  $\lfloor x \rfloor$  denote the greatest integer less than or equal to x, and let  $\{x\} = x - \lfloor x \rfloor$  be the fractional part of x. Find all real numbers a, b, and c such that

$$\lfloor a \rfloor + \{b\} = -2.3,$$

$$[b] + \{c\} = 8.9,$$
 and

$$|c| + \{a\} = 23.4.$$

The first equation gives us that  $\lfloor a \rfloor = -3$  and  $\{b\} = 0.7$ . The second equation gives us tht  $\lfloor b \rfloor = 8$  and  $\{c\} = 0.9$ . The last equation implies that  $\lfloor c \rfloor = 23$  and  $\{a\} = 0.4$ . It follows that a = -2.6, b = 8.7 and c = 23.9.

2. Lets PQRS be a cyclic quadrilateral such that PQ = QR, PR = SR and  $PQ \parallel SR$ . Let ASB be a tangent at S where A lies on PQ. If  $\angle BSR = 72^{\circ}$ , find the value of  $\angle RPQ$ .

By the tan-chord theorem, we have that  $\angle SPR = \angle BSR = 72^{\circ}$ . Since PR = SR, we find that  $\angle RSP = \angle SPR = 72^{\circ}$ . Since PQRS is a cyclic quadrilateral, we have that  $\angle PQR = 180^{\circ} - \angle RSP = 180^{\circ} - 72^{\circ} = 108^{\circ}$ .

Now note that  $\angle PQR + \angle RPQ + \angle QRP = 180^{\circ}$ . However, PQR is an isosceles triangle, so  $\angle RPQ = \angle QRP$ , and so we have that

$$2\angle RPQ + 108^{\circ} = \angle RPQ + \angle QRP + \angle PQR = 180^{\circ}$$
.


and so  $\angle RPQ = 36^{\circ}$ .

3. Let n be a positive integer. Find the last digit of

$$n^{2018} + (n+1)^{2018} + \dots + (n+99)^{2018}$$
.

4. Let ABC be a triangle, and let the midpoint of AC be M. The circle tangent to BC at B and passing through M meets the line AB again at P. Prove that  $AB \times BP = 2BM^2$ .

1



Construct point N, the midpoint of AB. By midpoint  $\operatorname{Th}^{\underline{m}}$ , NM||BC. Using this and Tan-Chord  $\operatorname{Th}^{\underline{m}}$ , we get:  $\angle BMN = \angle MBC = \angle MPB$ . Since we also have that  $\angle PBM = \angle MBN$ , we have that  $\triangle BMN \sim \triangle BPM$ . So

$$\frac{BM}{BP} = \frac{BN}{BM}$$
 
$$\Rightarrow BM^2 = BN \cdot BP = \left(\frac{1}{2}AB\right) \cdot BP$$

5. Show that for all positive real numbers x and y,

$$\frac{x^2}{x+2y}+\frac{y^2}{2x+y}\geq \frac{x+y}{3}.$$

By the Cauchy-Schwarz inequality (in "Engel Form") we have that

$$\frac{x^2}{x+2y} + \frac{y^2}{2x+y} \ge \frac{(x+y)^2}{(x+2y) + (2y+x)} = \frac{x+y}{3}.$$