

# ON A QUESTION OF KÜLSHAMMER FOR REDUCTIVE ALGEBRAIC GROUPS

DANIEL LOND AND BENJAMIN MARTIN

ABSTRACT.

## 1. INTRODUCTION

Let  $G$  and  $H$  be linear algebraic groups over an algebraically closed field of characteristic  $p > 0$ . Pick a maximal unipotent subgroup  $U$  of  $H$  [this is unique up to conjugacy]. By a *representation* of  $H$  in  $G$  we mean a homomorphism of algebraic groups from  $H$  to  $G$ . The group  $G$  acts on the set of representations  $\text{Hom}(H, G)$  by  $(g \cdot \rho)(h) = g\rho(h)g^{-1}$  for  $h \in H$  and  $g \in G$ ; we call the orbits *conjugacy classes*. We consider the following question.

**Question 1.1.** *Let  $\sigma: U \rightarrow G$  be a representation. Are there only finitely many conjugacy classes of representation  $\rho: H \rightarrow G$  such that  $\rho|_U$  is  $G$ -conjugate to  $\sigma$ ?*

Külshammer raised this question for finite  $H$  [1]; in this case the maximal unipotent subgroups are the Sylow  $p$ -subgroups. An example of Cram shows that the answer to Question 1.1 is no for  $H = S_3$  and  $G$  a certain 3-dimensional non-connected group with  $G^0$  unipotent [ref: Slodowy]. If  $G = \text{GL}_n$  and  $p$  does not divide  $|H|$  then the answer to Question 1.1 is yes, by Maschke's Theorem. It follows from a well-known geometric argument of Richardson that if  $p$  does not divide  $|H|$  then the answer to Question 1.1 is yes for any connected reductive  $G$ : one embeds  $G$  in some  $\text{GL}_n$  and studies the behavior of the induced map  $\text{Hom}(H, G) \rightarrow \text{Hom}(H, \text{GL}_n)$ . In fact, standard representation-theoretic results imply that the answer is yes for any finite group when  $G = \text{GL}_n$ , and Slodowy used Richardson's argument to show the answer is yes for arbitrary finite  $H$  and connected reductive  $G$  under mild hypotheses on  $\text{char}(p)$ . [General case of finite  $H$ : there is now a counterexample in type  $G_2$  [ref].]

In this paper we instead consider the case when  $H$  is connected and semisimple. We settle Question 1.1 as follows.

**Theorem 1.2.** *The answer to Question 1.1 is yes if  $H$  is connected and semisimple.*

Note that we allow  $G$  to be non-connected, but it is clear we can reduce immediately to the case when  $G$  is connected and  $Z(G)^0$  is trivial. Also, it is clear that the answer to Question 1.1 is no for arbitrary connected reductive  $H$  (e.g., just take  $H$  to be a torus,  $U = 1$  and  $G^0$  non-unipotent).

An important tool in the work of [above] is nonabelian 1-cohomology. Let  $\rho \in \text{Hom}(H, G)$  and let  $P$  be a parabolic subgroup of  $G$  such that  $\rho(H) \subseteq P$ . Then  $H$  acts on  $V := R_u(P)$  via  $h \cdot u = \rho(h)u\rho(h)^{-1}$ , and representations near  $\rho$  in an appropriate sense can be understood

---

1991 *Mathematics Subject Classification.* ??? (???)

*Key words and phrases.* ???

in terms of the nonabelian 1-cohomology  $H^1(H, \rho, V)$ . In Section [below] we discuss this cohomological approach and study its behavior with respect to restriction of representations from  $H$  to  $U$ . [Further discussion.]

[Based on D's PhD thesis. Related work by David Stewart: see below. He uses high-powered cohomology techniques; our methods are more elementary.]

[Acknowledgements: Marsden grants; Günter.]

## 2. PRELIMINARIES

[Some useful results from ‘‘On unipotent algebraic groups and 1-cohomology’’, David Stewart:

Cor. 3.4.3: Let  $B = TU$  be connected solvable group with unipotent radical  $U$  and maximal torus  $T$ . Let  $Q$  be a unipotent group on which  $B$  acts. Then the restriction map  $H^1(B, Q) \rightarrow H^1(U, Q)$  is injective.

Thm 3.5.2: Let  $G$  be connected reductive and let  $Q$  be a unipotent group on which  $G$  acts. Then for any parabolic subgroup  $P$  of  $G$ , the restriction map  $H^1(G, Q) \rightarrow H^1(P, Q)$  is an isomorphism of pointed sets.

Corollary 3.6.2. Let  $H$  be a closed, connected, reductive subgroup of  $G$  contained in a parabolic  $P = LR_u(P)$  of  $G$  and let  $P_1$  be any parabolic subgroup of  $H$ . Then  $H$  is  $G$ -conjugate to a subgroup of  $L$  if and only if  $P_1$  is as well.]

We assume  $G$  is a possibly non-connected algebraic group over  $k$  and  $H$  is a linear algebraic group over  $k$ . We fix a maximal unipotent subgroup  $U$  of  $H$ .

Suppose  $H$  is connected, let  $B$  be a Borel subgroup of  $H$ , let  $X$  be an affine variety and let  $f: H \rightarrow X$  be a morphism such that  $f(hb) = f(h)$  for all  $h \in H$  and all  $b \in B$ . Then  $f$  gives rise to a morphism  $\bar{f}$  from the projective variety  $H/B$  to  $X$ . Since  $H/B$  is connected and  $X$  is affine,  $\bar{f}$  must be constant, so  $f$  is constant. In particular, if  $V$  is an affine  $H$ -variety,  $v \in V$  and the stabiliser  $H_v$  contains  $B$  then  $H_v = H$ .

**Lemma 2.1.** *Suppose  $G$  is connected and reductive. Let  $H_1, H_2$  be connected reductive subgroups of  $G$ . Suppose  $B$  is a common Borel subgroup of both  $H_1$  and  $H_2$ . Then  $H_1 = H_2$ .*

*Proof.* The quotient variety  $G/H_1$  is affine since  $H_1$  is reductive [ref], and  $H_2$  acts on  $G/H_1$  by left multiplication. The stabiliser in  $H_2$  of the coset  $H_1$  contains  $B$ , so it must equal the whole of  $H_2$ . Hence  $H_2 \subseteq H_1$ . The reverse inequality follows similarly, so  $H_1 = H_2$ .  $\square$

**Lemma 2.2.** *Let  $B$  be a Borel subgroup of  $H$ . Let  $\rho_1, \rho_2 \in \text{Hom}(H, G)$  such that  $\rho_1|_B = \rho_2|_B$ . Then  $\rho_1 = \rho_2$ .*

*Proof.* Define  $f: H \rightarrow G$  by  $f(h) = \rho_1(h)\rho_2(h)^{-1}$ . For any  $h \in H$ ,  $b \in B$ ,  $f(hb) = \rho_1(hb)\rho_2(hb)^{-1} = \rho_1(h)\rho_1(b)\rho_2(b)^{-1}\rho_2(h)^{-1} = \rho_1(h)\rho_2(h)^{-1} = f(h)$ . So  $f$  gives rise to a morphism  $\bar{f}$  from the connected projective variety  $H/B$  to  $G$  given by  $\bar{f}(hB) = f(h)$ . Then  $\bar{f}$  is constant, so  $f$  is constant with value  $f(1) = 1$ , and the result follows.  $\square$

## 3. PROOF OF THEOREM 1.2 FOR $G$ REDUCTIVE

In this section we assume  $G$  and  $H$  are connected and semisimple.

**Lemma 3.1.** *Let  $B$  be the Borel subgroup of  $H$  that contains  $U$ , and let  $T$  be a maximal torus of  $B$ . Let  $\rho_1, \rho_2: H \rightarrow G$  be representations such that  $\rho_1(B) = \rho_2(B)$  and  $\rho_1(T) = \rho_2(T)$ . Set  $U' = \rho_1(U) = \rho_2(U)$ . Suppose that for all  $\alpha \in \Phi_T(B)$ ,  $\rho_1(U_\alpha) = \rho_2(U_\alpha)$  and there exists  $\phi_\alpha \in \text{Aut}(\rho_1(U_\alpha))$  such that for all  $u \in U_\alpha$ ,  $\rho_2(u) = \phi_\alpha(\rho_1(u))$ . Then there exists  $t' \in T$  such that  $t' \cdot \rho_2 = \rho_1$ .*

*Proof.* Since  $B$  is a Borel subgroup of  $H$ ,  $\rho_i(B)$  is a Borel subgroup of  $\rho_i(H)$  for  $i = 1, 2$ . Lemma 2.1 implies that  $\rho_1(H) = \rho_2(H)$ . Hence we can assume that  $\rho_1$  and  $\rho_2$  are surjective. Set  $T' = \rho_1(T) = \rho_2(T)$ .

Pick a base  $\{\alpha'_1, \dots, \alpha'_m\}$  for  $\Phi_{T'}(B')$ . Since  $\rho_1$  and  $\rho_2$  map a given root group of  $U$  with respect to  $T$  to a root group of  $U'$  with respect to  $T'$ , there exist  $\alpha_1, \dots, \alpha_m \in \Phi_T(B)$  such that  $\rho_1(U_{\alpha_i}) = \rho_2(U_{\alpha_i}) = U_{\alpha'_i}$  and  $\alpha'_i(\rho_1(t)) = \alpha'_i(\rho_2(t))$  for all  $i$ . [This is not quite true as stated: some of the  $U_{\alpha_i}$  may be mapped by  $\rho_1$  and  $\rho_2$  to 1; we're only interested in the case when this doesn't happen. A bit more argument is needed here.] As  $H$  is semisimple, this implies that  $\rho_1|_T = \rho_2|_T$ .

By hypothesis, each  $\phi_{\alpha_i}$  is an automorphism of  $U_{\alpha'_i}$ , so there exist  $b_1, \dots, b_m \in k^*$  such that  $\phi_{\alpha_i}(\epsilon_{\alpha'_i}(x)) = \epsilon_{\alpha'_i}(b_i x)$  for each  $i$  and for all  $x \in k$ ; so there exist  $a_1, \dots, a_m \in k^*$  such that  $\rho_2(\epsilon_{\alpha_i}(x)) = \rho_1(\epsilon_{\alpha_i}(a_i x))$  for each  $i$  and for all  $x \in k$  [check!].

The weights  $\alpha'_1, \dots, \alpha'_m$  are linearly independent as  $H$  is semisimple, so there exists  $t' \in T'$  such that  $\alpha_i(t') = \alpha_i^{-1}$  for  $1 \leq i \leq m$ . We then have  $(t' \cdot \rho_2)(\epsilon_{\alpha_i}(x)) = \rho_1(\epsilon_{\alpha_i}(x))$  for each  $i$  and for all  $x \in k$ . It follows that  $\rho_1|_{\tilde{U}} = (t' \cdot \rho_2)|_{\tilde{U}}$ , where  $\tilde{U}$  is the subgroup of  $U$  generated by the  $U_{\alpha_i}$ . But  $\rho_1(\tilde{U}) = \rho_2(\tilde{U}) = (t' \cdot \rho_2)(\tilde{U})$  since the  $U_{\alpha'_i}$  generate  $U'$  [ref: Humphreys?], and  $\ker(\rho_1|_U) = \ker(\rho_2|_U) = \ker((t' \cdot \rho_2)|_U)$ , so  $\rho_1|_U = (t' \cdot \rho_2)|_U$ .

To complete the proof, it is enough by Lemma 2.2 to show that  $\rho_1|_B = (t' \cdot \rho_2)|_B$ . But  $(t' \cdot \rho_2)|_T = \rho_2|_T = \rho_1|_T$  from the discussion above, so we are done.  $\square$

[Note: In the situation of the above lemma (and with  $\rho_1, \rho_2$  assumed surjective), suppose we don't assume that  $\rho_1(U_\alpha) = \rho_2(U_\alpha)$  for every  $\alpha$ . Then  $\rho_1$  and  $\rho_2$  need no longer be  $G$ -conjugate. I suspect, however, that in this case they are  $\text{Aut}(G)$ -conjugate. If this is the case then we should get a reasonable bound in Theorem 1.2 involving  $|\text{Out}(G)|$  and the size of the Weyl group of  $T'$  in  $N_G(U')$  (cf. below).]

**Theorem 3.2.** *The answer to Külshammer's question is yes for  $H$  and  $G$  connected and semisimple.*

*Proof.* Let  $\sigma: U \rightarrow G$  and let  $U' = \sigma(U)$ . Let  $B$  be the Borel subgroup of  $H$  that contains  $U$  and fix a maximal torus  $T$  of  $B$ . Fix a maximal torus  $T'$  of  $N_G(U')$ . Let  $C = \{\rho \in \text{Hom}(H, G) \mid \rho|_U = \sigma\}$ . If  $\rho \in C$  then  $\rho(T)$  normalizes  $\rho(U) = U'$ , so  $\rho(T)$  is a torus of  $N_G(U')$ . By conjugacy of maximal tori of  $N_G(U')$ , there exists  $g \in N_G(U')$  such that  $(g \cdot \rho)(T) \subseteq T'$ . If  $h \in N_G(U')$  and  $(h \cdot \rho)(T) \subseteq T'$  then there exists  $n \in N_{N_G(U')}(T')$  such that  $(h \cdot \rho)(T) = ((ng) \cdot \rho)(T)$ . The group  $N_{N_G(U')}(T')/C_{N_G(U')}(T')$  is finite, so there is a finite set of subtori  $S_1, \dots, S_r$  of  $T'$  such that  $\rho(T)$  is  $N_G(U')$ -conjugate to one of  $S_1, \dots, S_r$ .

Define a relation  $\equiv$  on  $C$  by  $\rho_1 \equiv \rho_2$  if there exist  $i \in \{1, \dots, r\}$  and  $g_1, g_2 \in N_G(U')$  such that  $(g_1 \cdot \rho_1)(T) = (g_2 \cdot \rho_2)(T) = S_i$  and  $\rho_1(U_\alpha) = \rho_2(U_\alpha)$  for all  $\alpha \in \Phi_T(U)$ . It is clear that  $\equiv$  is an equivalence relation. Define  $C_i = \{\rho \in G \mid (g \cdot \rho)(T) = S_i \text{ for some } g \in N_G(U')\}$ . If  $(g \cdot \rho)(T) = S_i$  then  $(g \cdot \rho)$  must map any root group  $U_\alpha$  of  $U$  with respect to  $T$  to a root group of  $U'$  with respect to  $S_i$ , so there are only finitely many possibilities for  $(g \cdot \rho)(U_\alpha)$ . It follows that  $\equiv$  has only finitely many equivalence classes.

To complete the proof it is enough, therefore, to show that if  $\rho_1 \equiv \rho_2$  then  $\rho_1$  and  $\rho_2$  are  $G$ -conjugate. So suppose  $g_1$  and  $g_2$  are [as in the defn of  $\equiv$ ]. Set  $U_{\alpha'} = \rho_1(U_{\alpha}) = \rho_2(\alpha)$  if this is nontrivial [recall that  $\rho_1(U_{\alpha})$  might be just 1]. Then the map  $\phi: G \rightarrow G$  given by  $\phi(g) = g_2 g_1^{-1} g g_1 g_2^{-1}$  gives rise to an automorphism of  $U'_{\alpha}$  for all  $\alpha' \in \Phi_{S_i}(U')$ . It follows by Lemma 3.1 that  $g_1 \cdot \rho_1$  and  $g_2 \cdot \rho_2$  are  $T$ -conjugate, so we are done.  $\square$

[Consequence: Look at **images** of representations and prove an analogous finiteness result, for connected reductive  $H$  as well?]

#### 4. 1-COHOMOLOGY

[Describe Daniel's approach using 1-cohomology.]

[We've proved Theorem 1.2 for reductive  $G$  (it's enough to take  $G$  to be connected and semisimple, since  $H$  is connected and semisimple). I think the theorem follows for arbitrary non-reductive  $G$  by taking the result for the special case of reductive groups and applying it to the reductive group  $G/V$ , then using David Stewart's result that the restriction map  $H^1(B, V) \rightarrow H^1(U, V)$  is injective. Here  $V = R_u(G)$  and  $B$  is a Borel subgroup of  $G$  with unipotent radical  $U$ .]

[An application:

Theorem: Let  $G$  be a reductive group such that the simple components of  $G$  are all of type  $A_n$  or  $B_2$ . Then the answer to K's question is yes for any  $H$  with  $H^0$  semisimple.

Proof (sketch): We can approximate  $H$  with a sufficiently large finite subgroup, so assume without loss that  $H$  is finite. It's enough to study the simple components of  $G$  separately, so without loss we assume  $G$  is simple. If  $G$  is of type  $B_2$  and  $p \neq 2$  then we're done by Slodowy's paper, so without loss we assume that either  $G$  is of type  $A_n$ , or  $G$  is of type  $B_2$  and  $p = 2$ . Note that maximal parabolics of  $G$  have abelian unipotent radicals under this hypothesis.

We now use an inductive proof. If  $\rho$  is  $G$ -irreducible then we're done. Otherwise put  $\rho$  into a maximal parabolic  $P$ . The key point is that we can transfer the problem from  $P$  into a Levi of  $P$  by the usual arguments, because the restriction map  $H^1(H, V) \rightarrow H^1(U, V)$  is injective for any finite  $H$  if  $V = R_u(P)$  is abelian. (Here  $U$  is a Sylow  $p$ -subgroup of  $H$ .) By induction on  $\dim(G)$ , the result is true for  $L$ —for the simple components of  $L$  are all of type  $A_n$  or  $B_2$ —and we get what we want.

This result shows that the  $G_2$  counterexample to K's question for finite  $H$  is in a sense the smallest possible: there is no such counterexample for any other  $G$  of rank 1 or 2.]

#### REFERENCES

- [1] B. Külshammer. Donovan's conjecture, crossed products and algebraic group actions. *Israel J. Math.*, 92(1–3):295–306, 1995.

???

*E-mail address:* ???

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ABERDEEN, KING'S COLLEGE, FRASER NOBLE BUILDING, ABERDEEN AB24 3UE, UNITED KINGDOM

*E-mail address:* B.Martin@abdn.ac.uk