

Analysis 3 : Measure Theory

Lecture by Ikhan Choi

Notes by Ikhan Choi

Contents

Chapter 1. Carathéodory's theory	5
1. σ -algebras	6
2. Carathéodory's extension theorem	7
Chapter 2.	9
Chapter 3. Topological measures	11
1. Descriptive set theory	12
2. Borel measures	13
2.1. Regular measures	13
2.2. Radon measures	13
3. Riesz-Markov-Kakutani representation theorem	15
3.1. Positive linear functional	15
Proof of the first theorem	15
Chapter 4. Hmmm	17
0.1. Convergence in measure	17

CHAPTER 1

Carathéodory's theory

1. σ -algebras

DEFINITION 1.1. Let X be a set. A *ring of sets* is a family of subsets of X that is closed under finite union and finite relative complement; in other words, $\mathcal{R} \subset \mathcal{P}(X)$ is called a ring of sets if the following two conditions are satisfied:

- (1) if $A, B \in \mathcal{R}$, then $A \cup B \in \mathcal{R}$,
- (2) if $A, B \in \mathcal{R}$, then $A \setminus B \in \mathcal{R}$.

PROPOSITION 1.1. *Let X be a set and $\mathcal{R} \subset \mathcal{P}(X)$. Then, the followings are equivalent:*

- (1) \mathcal{R} is a ring of sets,
- (2) \mathcal{R} is closed under symmetric difference and finite intersection,
- (3) \mathcal{R} is a ring,
- (4) \mathcal{R} is a Boolean ring.

For the ring structure, we take the symmetric difference as addition and the intersection as multiplication.

PROPOSITION 1.2. *A ring of sets is a distributive lattice.*

If a ring of sets contains a multiplicative identity, the entire set, then we call the ring of sets as follows:

DEFINITION 1.2. An *algebra of sets* is a ring of sets with the entire set.

PROPOSITION 1.3. *Let X be a set and $\mathcal{R} \subset \mathcal{P}(X)$. Then, the followings are equivalent:*

- (1) \mathcal{R} is an algebra of sets,
- (2) \mathcal{R} is closed under finite union, finite intersection, and complement,
- (3) \mathcal{R} is a Boolean algebra.

An algebra of sets is sometimes called a field of sets.

2. Carathéodory's extension theorem

THEOREM 2.1 (Carathéodory's extension theorem). *Let \mathcal{R} be a ring of sets over X . Let $\sigma(\mathcal{R})$ be the σ -algebra generated by \mathcal{R} . A set function $\mu : \mathcal{R} \rightarrow [0, \infty]$ is extended to a measure on $\sigma(\mathcal{R})$ if and only if it is a premeasure.*

CHAPTER 2

CHAPTER 3

Topological measures

1. Descriptive set theory

2. Borel measures

2.1. Regular measures.

THEOREM 2.1. *Let μ be a Borel measure. Then, the followings are equivalent:*

- (1) μ is inner regular on σ -bounded sets
- (2) μ is outer regular on σ -bounded sets.

2.2. Radon measures. Locally compact Hausdorff spaces have at least two important applications in abstract analysis related to measure theory: one is locally compact groups and the associated Harr measures in abstract harmonic analysis, the other is the Gelfand-Naimark theorem which states every commutative C^* -algebra can be represented as a function space on a locally compact Hausdorff space. In the set of this section, we assume every base space X is locally compact Hausdorff.

Note that locally finite measures are compact finite but the converser holds only if in locally compact Hausdorff spaces. We want to consider locally finite Borel measures as the minimally compatible measures with a given topology on X . For locally finite Borel measures, a set is finite-measured if and only if it is contained in a compact set.

DEFINITION 2.1. (Folland's) A *Radon measure* is a Borel measure on X which satisfies the following three conditions:

- (1) locally finite,
- (2) outer regular on all Borel sets,
- (3) inner regular on all open sets.

Radon measures are rather simply characterized when the base space X is σ -compact. The following proposition proves the equivalence between regularity and Radonness of locally finite Borel measure on a σ -compact space.

PROPOSITION 2.2. *A Radon measure is inner regular on all σ -finite Borel sets. (Folland's)*

PROOF. First we approximate Borel sets of finite measure, with compact sets. Let E be a Borel set with $\mu(E) < \infty$ and U be an open set containing E . By outer regularity, there is an open set $V \supset U - E$ such that

$$\mu(V) < \mu(U - E) + \frac{\varepsilon}{2}.$$

By inner regularity, there is a compact set $K \subset U$ such that

$$\mu(K) > \mu(U) - \frac{\varepsilon}{2}.$$

Then, we have a compact set $K - V \subset K - (U - E) \subset E$ such that

$$\begin{aligned} \mu(K - V) &\geq \mu(K) - \mu(V) \\ &> \left(\mu(U) - \frac{\varepsilon}{2}\right) - \left(\mu(U - E) + \frac{\varepsilon}{2}\right) \\ &\geq \mu(E) - \varepsilon. \end{aligned}$$

It implies that a Radon measure is inner regular on Borel sets of finite measures.

Suppose E is a σ -finite Borel set so that $E = \bigcup_{n=1}^{\infty} E_n$ with $\mu(E_n) < \infty$. We may assume E_n are pairwise disjoint. Let K_n be a compact subset of E_n such that

$$\mu(K_n) > \mu(E_n) - \frac{\varepsilon}{2^n},$$

and define $K = \bigcup_{n=1}^{\infty} K_n \subset E$. Then,

$$\mu(K) = \sum_{n=1}^{\infty} \mu(K_n) > \sum_{n=1}^{\infty} \left(\mu(E_n) - \frac{\varepsilon}{2^n} \right) = \mu(E) - \varepsilon.$$

Therefore, a Radon measure is inner regular on all σ -finite Borel sets. \square


We get a corollary:

COROLLARY 2.3. *If X is σ -compact, then a locally finite Borel measure is Radon if and only if it is regular.*

THEOREM 2.4. *If every open set in X is σ -compact (i.e. Borel sets and Baire sets coincide), then every locally finite Borel measure is regular.*

PROPOSITION 2.5. *In a second countable space, every open set is σ -compact (i.e. Borel sets and Baire sets coincide).*

Two corollaries are presented as follows:



Many applications assume X is an open subset of a Euclidean space, so X is usually second countable. In this case, the followings will be synonym: A measure is

(1)

$$L_{\text{loc}}^1 = \text{absolutely continuous measures} \subset \text{Radon measures} \subset \mathcal{D}'.$$

THEOREM 2.6. *Every finite Radon measure is regular.*

3. Riesz-Markov-Kakutani representation theorem

In this section, we always assume X is a locally compact Hausdorff space. Hence we can use the Urysohn lemma in the following way: If a compact subset K and a closed subset F are disjoint, then by applying the Urysohn lemma on a compact neighborhood of K , we can find a continuous function $f : X \rightarrow [0, 1]$ such that $f|_K = 1$ and $f|_F = 0$. In particular, there always exists a “continuous characteristic function” $\phi \in C_c(X)$ with $\phi|_K = 1$.

There are two Riesz-Markov-Kakutani theorems: the first theorem describes the positive elements in $C_c(X)^*$ as Radon measures when the natural colimit topology is assumed, and the second theorem describes $C_c(X)^*$ as the space of finite Radon measures when uniform topology is assumed.

3.1. Positive linear functional. Positivity of linear functional itself implies a rather strong continuity property.

THEOREM 3.1. *Let X be a locally compact Hausdorff space. If a linear functional on $C_c(X)$ is positive, then it is continuous with respect to the inductive topology.*

PROOF. Let I be a positive linear functional on $C_c(X)$. Take an open subset U of X that has a compact closure. Choose a nonnegative $\phi \in C_c(X)$ such that $\phi|_{\bar{U}} = 1$ using the Urysohn lemma. Since

$$|I(f)| \leq I(|f|) = I(\phi|f|) \leq I(\phi\|f\|) = I(\phi)\|f\|_{\sup}$$

for $f \in C_c(U)$,

$$\|I\|_{C_c^*(U)} \leq I(\phi) < \infty.$$

□

Borel regular measure is tight on every measurable set.

Proof of the first theorem. A regular measure of an open set U can be realized by the norm of integration functional on $C_c(U)$.

LEMMA 3.2. *Let μ be a Borel regular measure on a locally compact Hausdorff space X . Then,*

$$\mu(U) = \|I_\mu\|_{C_c^*(U)}$$

for every open U in X , where I_μ is a linear functional on $C_c(U)$ defined by

$$I_\mu(f) := \int f d\mu.$$

PROOF. (\geq) For $f \in C_c(U)$, we have

$$|\int f d\mu| = |\int_U f d\mu| \leq \mu(U)\|f\|_{\sup}.$$

(\leq) Since μ is tight on U , there is $K \in U$ such that $\mu(U \setminus K) < \varepsilon$. By the construction of Urysohn, we can find a nonnegative function $f \in C_c(U)$ with $f|_K \equiv 1$ and $\|f\|_{C_c(U)} = 1$ so that

$$\mu(K) \leq \int f d\mu.$$

Thus,

$$\mu(U) - \varepsilon < \int f d\mu \leq \|I_\mu\|_{C_c(U)}$$

for arbitrary $\varepsilon > 0$. □

THEOREM 3.3 (The Riesz-Markov-Kakutani representation theorem). *Let X be a locally compact Hausdorff space. Let $I : C_c(X) \rightarrow \mathbb{R}$ be a positive linear functional. Then, there is a unique Radon measure μ on X such that*

$$I(f) = \int f d\mu.$$

Note that any continuity of I is not assumed except positivity.

PROOF. For existence, define μ by

$$\mu(U) := \|I\|_{C_c^*(U)}$$

for open sets and

$$\mu(E) := \inf_{U \supset E} \mu(U)$$

for Borel sets. We need to show

- (1) countable additivity
- (2) inner regularity
- (3) $I(f) = \int f d\mu$.

□

Jordan decomposition: $(C_0(X), \|\cdot\|)^* = (C_c(X), \|\cdot\|)^* \subset (C_c(X), \lim_{\rightarrow})^*$ converse?

CHAPTER 4

Hmmmm

0.1. Convergence in measure. Since $\{f_n(x)\}_n$ diverges if and only if

$$\exists k > 0, \quad \forall n_0 > 0. \quad \exists n > n_0 : \quad |f_n(x) - f(x)| > n^{-1},$$

we have

$$\begin{aligned} \{x : \{f_n(x)\}_n \text{ diverges}\} &= \bigcup_{k>0} \bigcap_{n_0>0} \bigcup_{n>n_0} \{x : |f_n(x) - f(x)| > n^{-1}\} \\ &= \bigcup_{k>0} \limsup_n \{x : |f_n(x) - f(x)| > n^{-1}\}. \end{aligned}$$

Since for every k

$$\limsup_n \{x : |f_n(x) - f(x)| > k^{-1}\} \subset \limsup_n \{x : |f_n(x) - f(x)| > n^{-1}\},$$

we have

$$\{x : \{f_n(x)\}_n \text{ diverges}\} \subset \limsup_n \{x : |f_n(x) - f(x)| > n^{-1}\}.$$

THEOREM 0.1. *Let f_n be a sequence of measurable functions on a measure space (X, μ) . If f_n converges to f in measure, then f_n has a subsequence that converges to f μ -a.e.*

PROOF. Since $d_{f_n-f}(1/k) \rightarrow 0$ as $n \rightarrow \infty$, we can extract a subsequence f_{n_k} such that

$$\mu(\{x : |f_{n_k}(x) - f(x)| > k^{-1}\}) < 2^{-k}.$$

Since

$$\sum_{k=1}^{\infty} \mu(\{x : |f_{n_k}(x) - f(x)| > k^{-1}\}) < \infty,$$

by the Borel-Canteli lemma, we get

$$\mu(\limsup_k \{x : |f_{n_k}(x) - f(x)| > k^{-1}\}) = 0.$$

Therefore, f_{n_k} converges μ -a.e. □