

Finite Group Theory

IKHAN CHOI

CONTENTS

1. Sylow game	1
2. Simple groups	3
2.1. Symmetric groups	3
2.2. Linear groups	3
3. Extensions	3

1. SYLOW GAME

Definition 1.1 (Sylow p -subgroup). Let G be a finite group of order $n = p^a m$ for a prime $p \nmid m$. A *Sylow p -subgroup* is a subgroup of order p^a . We are going to denote the set of Sylow p -subgroups by $\text{Syl}_p(G)$ and the number of Sylow p -subgroups by $n_p(G)$.

Theorem 1.1 (The Sylow theorem). *Let G be a finite group of order $n = p^a m$ for a prime $p \nmid m$. Then,*

$$p \mid n_p - 1, \quad n_p \mid m$$

for some $k \in \mathbb{N}$.

Proof. Step 1: Sylow p -subgroups exist. We apply mathematical induction. The base step is trivial. Suppose every finite group of order less than n possesses a Sylow p -subgroup.

By applying the orbit-stabilizer theorem for the action $G \curvearrowright G$ by conjugation, build the class equation

$$|G| = |Z(G)| + \sum_i |G : C_G(g_i)|.$$

There are two cases: $p \mid |Z(G)|$ or $p \nmid |Z(G)|$.

Case 1: $p \mid |Z(G)|$. The group G has a normal subgroup of order p by applying Cauchy's theorem for abelian groups on the center. Then, the inverse image of a Sylow p -subgroup of the quotient group is also a Sylow p -subgroup of G .

Case 2: $p \nmid |Z(G)|$. Since $p \mid n$, we have $p \nmid |G : C_G(g)|$ for some $g \in G$. Then, a Sylow p -subgroup of the centralizer is also a Sylow p -subgroup of G .

Therefore, we are done for Step 1.

First Written : November 7, 2019.

Last Updated : November 7, 2019.

Step 2: Sylow p -subgroup that is normal is unique. Note that p does not divide the order of the quotient group. Every p -subgroup should be contained in the Sylow p -subgroup, the kernel of the quotient map. The Sylow p -subgroup is clearly unique.

Step 3: Sylow p -subgroups get action by conjugation. Let P be a Sylow p -subgroup of G . We construct class equations via the orbit-stabilizer theorem for various actions to extract information on n_p . Note that stabilizers in any setwise conjugation action is exactly normalizers.

- (1) The action $P \curvearrowright \text{Syl}_p(G)$ gives

$$n_p = 1 + \sum_i |P : N_P(P_i)|$$

since $P = N_P(P_i)$ implies $P \trianglelefteq N_G(P_i)$ and $P = P_i$.

- (2) Suppose the action $G \curvearrowright \text{Syl}_p(G)$ is not transitive. Take another Sylow p -subgroup P' is not conjugate with P in G . The two actions $P \curvearrowright \text{Orb}_G(P)$ and $P' \curvearrowright \text{Orb}_G(P)$ gives

$$|\text{Orb}_G(P)| = 1 + \sum_i |P : N_P(P_i)| = \sum_i |P' : N_{P'}(P_i)|.$$

It deduces $|\text{Orb}_G(P)| \equiv 0, 1 \pmod{p}$ simultaneously, which leads a contradiction.

- (3) The action $G \curvearrowright \text{Syl}_p(G)$ gives

$$n_p = |G : N_G(P_i)|$$

for all $P_i \in \text{Syl}_p(G)$ because the action is transitive.

Then, (1) proves $p \mid n_p - 1$, and (3) proves $n_p \mid m$. □

Corollary 1.2. *Let G be a finite group. Then,*

- (1) *every pair of two Sylow p -subgroup is conjugate.*
- (2) *every p -subgroup is contained in a Sylow p -subgroup.*
- (3) *a Sylow p -subgroup is normal if and only if $n_p = 1$.*

Theorem 1.3. *Alternative proof for existence of p -groups.*

Proof. Let $|G| = p^{a+b}m$. Let \mathcal{P}_{p^a} be the set of all p^a -sets in G . Give $G \curvearrowright \mathcal{P}_{p^a}$ by left multiplication. Since $v_p(|\mathcal{P}_{p^a}|) = v_p\left(\binom{p^{a+b}m}{p^a}\right) = b$, there is an orbit \mathcal{O} such that $v_p(|\mathcal{O}|) \leq b$. We have transitive action $G \curvearrowright \mathcal{O}$ and the stabilizer H satisfies $p^a \mid |G|/|\mathcal{O}| = |H|$. Since $H \curvearrowright \mathcal{O}$ trivially, $H \curvearrowright A$ for $A \in \mathcal{O} \subset \mathcal{P}_{p^a}$. It is only possible when $H \subset A$, hence $|H| = p^a$. □

What we want to find is subgroup lattices. A subgroup lattice particularly contains data about orders and conjugacy classes of subgroups.

In order to show the existence of subgroups of particular orders:

- (1) p -group theory, (including Cauchy and Sylow)
- (2) extension theory, (what can subgroups of subgroups do?)
- (3) normalizers,
- (4) kernel of permutation representation

In order to find the size of conjugacy classes:

- (1) measure the order of normalizers, (find some groups normalize a subgroup)
- (2) count elements,

2. SIMPLE GROUPS

2.1. Symmetric groups.

2.2. Linear groups.

3. EXTENSIONS

Proposition 3.1. *Let N and H be groups. Then, the following objects have one-to-one correspondences among each other.*

- (1) *isomorphic types of groups G such that a sequence*

$$0 \rightarrow N \rightarrow G \rightarrow H \rightarrow 0$$

is exact and right split,

- (2) *isomorphic types of groups G such that $N \trianglelefteq G \geq H$ with $G = NH$ and $N \cap H = 1$,*
- (3) *group actions $H \curvearrowright N$ preserving the group structure of N .*

Definition 3.1. The group G in the previous proposition is called the *semidirect product* of N and H .