

# **Analysis 1 : Foundations of Calculus**

Lecture by Ikhan Choi

Notes by Ikhan Choi



# Contents

Preface	i
Chapter 1. Real numbers	1
1. Fields	2
2. Orders	3
3. Norms	4
4. Completeness	5
5. The real numbers	6
5.1. Constructions of the real numbers	6
5.2. Cardinality	6
5.3. Elementary functions	6
Chapter 2. Sequences	7
Chapter 3. Functions	9
Chapter 4. Integration	11
1. Lebesgue spaces	12
1.1. $1 \leq p \leq \infty$	12
1.2. $L^2$ space	12
Chapter 5. Multivariable calculus	13
0.1. Inverse function theorem	13



## Preface

Before 19th century, theory of limits, infinite series, differentiation, and integration was so focused on calculating particular values of expressions, such as infinite series, that they do not have sufficient rigor. As an example, A. L. Cauchy, a pioneer of mathematical analysis, is known to have made many mistakes on his theses. Thereby, establishing a rigid framework for analysis has emerged as one of the central problems of mathematics in 19th century.



## CHAPTER 1

# **Real numbers**

## 1. Fields



## **2. Orders**

### **3. Norms**

## 4. Completeness

## **5. The real numbers**

**5.1. Constructions of the real numbers.**

**5.2. Cardinality.**

**5.3. Elementary functions.**

## CHAPTER 2

# Sequences



## CHAPTER 3

# **Functions**





## CHAPTER 4

# **Integration**

### 1. Lebesgue spaces

#### 1.1. $1 \leq p \leq \infty$ .

DEFINITION 1.1. Let  $f$  be a measurable real-valued function on a measure space  $(X, \mu)$ . The  $L^p$  norm of  $f$  is defined as a real number

$$\|f\|_{L^p(X)} := \left( \int |f|^p d\mu \right)^{\frac{1}{p}}$$

for  $0 < p < \infty$ , and

$$\|f\|_{L^\infty(X)} := \operatorname{ess\,sup}_{x \in X} |f(x)|$$

for  $p = \infty$ .

DEFINITION 1.2. The *essential supremum* is defined by the smallest number  $M$  such that

$$|f(x)| \leq M \quad a.e.$$

We often abbreviate  $\|f\|_{L^p(X)}$  as  $\|f\|_{L^p}$  or  $\|f\|_p$ .

#### 1.2. $L^2$ space.

## CHAPTER 5

### Multivariable calculus

$T < \min\{1/K, R/M\}$  and  $|f(t, x)| \leq M$  on  $[0, T] \times \overline{B(x_0, R)}$

$$x' = f(t, x), \quad x(0) = x_0.$$

$$\phi(x)(t) = x_0 + \int_0^t f(s, x(s)) ds.$$

$$\begin{aligned} |\phi(x)(t) - x_0| &\leq \int_0^t |f(s, x(s))| ds \\ &\leq TM \end{aligned}$$

$$\begin{aligned} |\phi(g)(t) - \phi(h)(t)| &= \left| \int_0^t (f(s, g(s)) - f(s, h(s))) ds \right| \\ &\leq \int_0^t |f(s, g(s)) - f(s, h(s))| ds \\ &\leq \int_0^t K |g(s) - h(s)| ds \\ &\leq TK \sup_s |g(s) - h(s)| \end{aligned}$$

#### 0.1. Inverse function theorem.

PROOF 1. Take  $y$ . We must first define  $x$  such that  $f(x) = y$ . Let

$$\varphi_y(x) = x + df_p^{-1}(y - f(x)).$$

Note that  $f(x) = y$  if and only if  $x$  is a fixed point of  $\varphi_y$ .

Since  $df$  is continuous, there is an open ball  $U$  such that

$$\|df(x) - df(p)\| < \frac{1}{2\|df(p)^{-1}\|}.$$

For  $x \in U$ ,

$$d\varphi_y(x) = \text{id} - df_p^{-1}df_x$$

□