

Diachrony of Spectra

Ikhan Choi

Postech - Unist - Kaist Joint Seminar

July 30, 2019

Introduction

Introduction

Definition

Let R be a commutative ring. The *spectrum* of R is the set of prime ideals of R .

Introduction

Definition

Let R be a commutative ring. The *spectrum* of R is the set of prime ideals of R .

Example

$$\operatorname{Spec}(\mathbb{Z}) = \{2\mathbb{Z}, 3\mathbb{Z}, 5\mathbb{Z}, 7\mathbb{Z}, 11\mathbb{Z}, \dots\}.$$

Introduction

Definition

Let R be a commutative ring. The *spectrum* of R is the set of prime ideals of R .

Example

$$\operatorname{Spec}(\mathbb{Z}) = \{2\mathbb{Z}, 3\mathbb{Z}, 5\mathbb{Z}, 7\mathbb{Z}, 11\mathbb{Z}, \dots\}.$$

Question

Why is it defined like this?

Contents


Hydrogen atom

Spectral theory on Hilbert spaces


Gelfand theory

Algebraic geometry

Hydrogen spectral series



Hydrogen spectral series




410.2nm

468.1nm

656.3 nm

434.0nm

Hydrogen spectral series



410.2nm

468.1nm

656.3 nm

434.0nm

Question

How can we explain and compute this phenomenon?

Rydberg's formula : Bohr model

Bohr's postulates:

Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.

Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.
- ▶ The stationary orbits are computed by the old quantization assumption for angular momenta:

$$mvr = n\hbar.$$

Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.
- ▶ The stationary orbits are computed by the old quantization assumption for angular momenta:

$$mvr = n\hbar.$$

- ▶ An electron absorbs or emits light frequency f when they jump from an orbit to another, satisfying

$$\Delta E = hf.$$

Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.
- ▶ The stationary orbits are computed by the old quantization assumption for angular momenta:

$$mvr = n\hbar.$$

- ▶ An electron absorbs or emits light frequency f when they jump from an orbit to another, satisfying

$$\Delta E = hf.$$

The constant h is called the Planck constant and $\hbar := \frac{h}{2\pi}$.

Rydberg's formula : Bohr model

From the three relations

$$mvr = n\hbar, \quad \frac{mv^2}{r} = -k \frac{(+e)(-e)}{r^2}, \quad E = K + V = \frac{1}{2}mv^2 - k \frac{e^2}{r},$$

Rydberg's formula : Bohr model

From the three relations

$$mvr = n\hbar, \quad \frac{mv^2}{r} = -k \frac{(+e)(-e)}{r^2}, \quad E = K + V = \frac{1}{2}mv^2 - k \frac{e^2}{r},$$

we deduce

$$E = -\frac{k^2 e^4 m}{2\hbar^2} \frac{1}{n^2} \approx -13.6 \frac{1}{n^2} \text{ (eV)}.$$

Rydberg's formula : Bohr model

From the three relations

$$mvr = n\hbar, \quad \frac{mv^2}{r} = -k \frac{(+e)(-e)}{r^2}, \quad E = K + V = \frac{1}{2}mv^2 - k \frac{e^2}{r},$$

we deduce

$$E = -\frac{k^2 e^4 m}{2\hbar^2} \frac{1}{n^2} \approx -13.6 \frac{1}{n^2} \text{ (eV)}.$$

Proposition (Rydberg formula)

The wavelengths λ of absorbed or emitted photons from a hydrogen atom is estimated by the following formula:

$$\frac{1}{\lambda} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right), \quad \text{for } n_1, n_2 \in \mathbb{N},$$

where $R := \frac{k^2 e^4 m}{4\pi\hbar^3 c}$ is the Rydberg constant.

Rydberg's formula : Schrödinger equation

More mathematically!

Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where V is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where V is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

By solving it, we obtain the probability distribution $|\Psi(t, x)|^2$ of the electron at time t ,

Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where V is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

By solving it, we obtain the probability distribution $|\Psi(t, x)|^2$ of the electron at time t , hence the assumption $\forall t, \int |\Psi(t, x)|^2 dx < \infty$.

Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where V is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

By solving it, we obtain the probability distribution $|\Psi(t, x)|^2$ of the electron at time t , hence the assumption $\forall t, \int |\Psi(t, x)|^2 dx < \infty$.

Let's solve.

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, \mathbf{x})}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, \mathbf{x}) + V(\mathbf{x}) \Psi(t, \mathbf{x}).$$

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, \mathbf{x})}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, \mathbf{x}) + V(\mathbf{x}) \Psi(t, \mathbf{x}).$$

“Mathematization”:

$$i\partial_t \Psi(t, \mathbf{x}) = (-\Delta + V(\mathbf{x})) \Psi(t, \mathbf{x}).$$

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form $\Psi(t, x) = \phi(t)\psi(x)$, then

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form $\Psi(t, x) = \phi(t)\psi(x)$, then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)}$$

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form $\Psi(t, x) = \phi(t)\psi(x)$, then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form $\Psi(t, x) = \phi(t)\psi(x)$, then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

The constant E is interpreted as the energy of electrons.

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, \mathbf{x})}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, \mathbf{x}) + V(\mathbf{x}) \Psi(t, \mathbf{x}).$$

“Mathematization”:

$$i\partial_t \Psi(t, \mathbf{x}) = (-\Delta + V(\mathbf{x})) \Psi(t, \mathbf{x}).$$

Ansatz: if the solutions has the form $\Psi(t, \mathbf{x}) = \phi(t)\psi(\mathbf{x})$, then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(\mathbf{x}))\psi(\mathbf{x})}{\psi(\mathbf{x})} = E.$$

The constant E is interpreted as the energy of electrons.

We have two *eigenvalue problems* with *shared eigenvalue* E :

Rydberg's formula : Schrödinger equation

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form $\Psi(t, x) = \phi(t)\psi(x)$, then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

The constant E is interpreted as the energy of electrons.

We have two *eigenvalue problems* with *shared eigenvalue* E :

$$i\frac{d}{dt}\phi(t) = E\phi(t), \quad (-\Delta + V(x))\psi(x) = E\psi(x).$$

Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions $\phi_E(t)$, $\psi_E(x)$ of the eigenvalue problems for each complex number E .

Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions $\phi_E(t)$, $\psi_E(x)$ of the eigenvalue problems for each complex number E .

Some facts:

Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions $\phi_E(t)$, $\psi_E(x)$ of the eigenvalue problems for each complex number E .

Some facts:

- ▶ For a given E , ϕ_E and ψ_E are of course not unique. In fact they form a vector space which is called the eigenspace.

Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions $\phi_E(t)$, $\psi_E(x)$ of the eigenvalue problems for each complex number E .

Some facts:

- ▶ For a given E , ϕ_E and ψ_E are of course not unique. In fact they form a vector space which is called the eigenspace.
- ▶ Note that for some E we probably cannot find the solution $\psi_E(x)$ that satisfies $\int |\psi_E(x)|^2 dx = 1$: the eigenspace is trivial.

Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions $\phi_E(t)$, $\psi_E(x)$ of the eigenvalue problems for each complex number E .

Some facts:

- ▶ For a given E , ϕ_E and ψ_E are of course not unique. In fact they form a vector space which is called the eigenspace.
- ▶ Note that for some E we probably cannot find the solution $\psi_E(x)$ that satisfies $\int |\psi_E(x)|^2 dx = 1$: the eigenspace is trivial.
- ▶ All functions that look like

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation and *all solutions are found in this way*(think physically!).

Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions $\phi_E(t)$, $\psi_E(x)$ of the eigenvalue problems for each complex number E .

Some facts:

- ▶ For a given E , ϕ_E and ψ_E are of course not unique. In fact they form a vector space which is called the eigenspace.
- ▶ Note that for some E we probably cannot find the solution $\psi_E(x)$ that satisfies $\int |\psi_E(x)|^2 dx = 1$: the eigenspace is trivial.
- ▶ All functions that look like

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation and *all solutions are found in this way* (think physically!).

- ▶ Since $\phi_E(t) \propto e^{-iEt}$ is easily solved, the main difficulty is ψ_E .

Rydberg's formula : Schrödinger equation

So, here are what we need to investigate seriously:

Rydberg's formula : Schrödinger equation

So, here are what we need to investigate seriously: *what are the eigenvalues and eigenvectors of the linear operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

defined by

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

Also, how can we compute them?

Rydberg's formula : Schrödinger equation

So, here are what we need to investigate seriously: *what are the eigenvalues and eigenvectors of the linear operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

defined by

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

Also, how can we compute them?

*** **The Beginning of Spectral Theory** ***

Rydberg's formula : Schrödinger equation

So, here are what we need to investigate seriously: *what are the eigenvalues and eigenvectors of the linear operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

defined by

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

Also, how can we compute them?

*** The Beginning of Spectral Theory ***

Remark

Simply, $L^2(\mathbb{R}^3)$ is the space of $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $\int |f|^2 < \infty$.

Rydberg's formula : Schrödinger equation

So, here are what we need to investigate seriously: *what are the eigenvalues and eigenvectors of the linear operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

defined by

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

Also, how can we compute them?

*** The Beginning of Spectral Theory ***

Remark

Simply, $L^2(\mathbb{R}^3)$ is the space of $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $\int |f|^2 < \infty$.

Remark

In fact, there are many technical issues to formalize this problem, for example, L^2 function is in general not differentiable.

Rydberg's formula : Schrödinger equation

So, here are what we need to investigate seriously: *what are the eigenvalues and eigenvectors of the linear operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

defined by

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

Also, how can we compute them?

*** The Beginning of Spectral Theory ***

Remark

Simply, $L^2(\mathbb{R}^3)$ is the space of $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $\int |f|^2 < \infty$.

Remark

In fact, there are many technical issues to formalize this problem, for example, L^2 function is in general not differentiable.

Don't be so pedantic in doing physics.

Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts have found the following result:

Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts have found the following result:

Proposition

The eigenvalues of the operator $\mathcal{H} = -\Delta - 2|x|^{-1}$ are

Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts have found the following result:

Proposition

The eigenvalues of the operator $\mathcal{H} = -\Delta - 2|x|^{-1}$ are

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts have found the following result:

Proposition

The eigenvalues of the operator $\mathcal{H} = -\Delta - 2|x|^{-1}$ are

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

with multiplicity 1, 4, 9, 16 and so on.

- Eigenvalues embody the possible energies of an electron, so we can give the Rydberg formula a reasonable explanation.

Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts have found the following result:

Proposition

The eigenvalues of the operator $\mathcal{H} = -\Delta - 2|x|^{-1}$ are

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

with multiplicity 1, 4, 9, 16 and so on.

- ▶ Eigenvalues embody the possible energies of an electron, so we can give the Rydberg formula a reasonable explanation.
- ▶ This result explains not only the discretized energy spectrum but also the number of each orbitals!

Conclusion of Section 1

Partial Differential Equations with Time Evolution



Separation of variables

Simultaneous Eigenvalue Problems



Study of Eigenvalues = Study of Hydrogen Spectrum

Contents

Hydrogen atom

Spectral theory on Hilbert spaces

Gelfand theory

Algebraic geometry

Spectral theory?

So far, we have seen that spectral theory refers to the theory about eigenvalues and eigenvectors,

Spectral theory?

So far, we have seen that spectral theory refers to the theory about eigenvalues and eigenvectors, especially for INFINITE dimensional linear operators.

Spectral theory?

So far, we have seen that spectral theory refers to the theory about eigenvalues and eigenvectors, especially for INFINITE dimensional linear operators.

In this section,

- ▶ we review the spectral theory on finite dimensional vector spaces
- ▶ and introduce Hilbert spaces — a typical example of infinite dimensional vector spaces — to state some results which extend the spectral theory to infinite dimensional spaces.

Spectral theory?

So far, we have seen that spectral theory refers to the theory about eigenvalues and eigenvectors, especially for INFINITE dimensional linear operators.

In this section,

- ▶ we review the spectral theory on finite dimensional vector spaces
- ▶ and introduce Hilbert spaces — a typical example of infinite dimensional vector spaces — to state some results which extend the spectral theory to infinite dimensional spaces.

From now, we basically assume \mathbb{C} as the scalar field for vector spaces.

Spectral theorem for normal matrices

The term “spectral theorem” is given to several theorems that show a condition for a linear operator to be diagonalizable.

Spectral theorem for normal matrices

The term “spectral theorem” is given to several theorems that show a condition for a linear operator to be diagonalizable. (diagonalizability is the firstly important “spectral property”!)

Spectral theorem for normal matrices

The term “spectral theorem” is given to several theorems that show a condition for a linear operator to be diagonalizable.

(diagonalizability is the firstly important “spectral property”!)

In particular, spectral theorems state the relation

condition related to adjoint \iff a kind of diagonalizability.

Spectral theorem for normal matrices

The term “spectral theorem” is given to several theorems that show a condition for a linear operator to be diagonalizable.

(diagonalizability is the firstly important “spectral property”!)

In particular, spectral theorems state the relation

condition related to adjoint \iff a kind of diagonalizability.

The most famous one is:

Definition

Let V be a finite dimensional inner product space of complex scalars and $A : V \rightarrow V$ be linear. (i.e., let A be a complex square matrix.) Then, A is said to be *normal* if $AA^* = A^*A$.

Spectral theorem for normal matrices

The term “spectral theorem” is given to several theorems that show a condition for a linear operator to be diagonalizable.

(diagonalizability is the firstly important “spectral property”!)

In particular, spectral theorems state the relation

condition related to adjoint \iff a kind of diagonalizability.

The most famous one is:

Definition

Let V be a finite dimensional inner product space of complex scalars and $A : V \rightarrow V$ be linear. (i.e., let A be a complex square matrix.) Then, A is said to be *normal* if $AA^* = A^*A$.

Note that the complex conjugate depends on the inner product structure of V .

Spectral theorem for normal matrices

The term “spectral theorem” is given to several theorems that show a condition for a linear operator to be diagonalizable.

(diagonalizability is the firstly important “spectral property”!)

In particular, spectral theorems state the relation

condition related to adjoint \iff a kind of diagonalizability.

The most famous one is:

Definition

Let V be a finite dimensional inner product space of complex scalars and $A : V \rightarrow V$ be linear. (i.e., let A be a complex square matrix.) Then, A is said to be *normal* if $AA^* = A^*A$.

Note that the complex conjugate depends on the inner product structure of V .

Theorem (Spectral theorem for normal matrices)

A complex square matrix A is normal if unitarily normalizable.

Spectral theorem of normal matrices

Hilbert space

Bounded operators

Spectral theorem of compact normal operators

Spectral theorem of elliptic operators (Bonus)

The operator $-\Delta - 2|x|^{-1}$ is an example of what we call *elliptic operators*.

Spectral theorem of elliptic operators (Bonus)

The operator $-\Delta - 2|x|^{-1}$ is an example of what we call *elliptic operators*. We will not deal with this, but:

Spectral theorem of elliptic operators (Bonus)

The operator $-\Delta - 2|x|^{-1}$ is an example of what we call *elliptic operators*. We will not deal with this, but:

The operator $-\Delta - 2|x|^{-1}$ is unbounded,

Spectral theorem of elliptic operators (Bonus)

The operator $-\Delta - 2|x|^{-1}$ is an example of what we call *elliptic operators*. We will not deal with this, but:

The operator $-\Delta - 2|x|^{-1}$ is unbounded, but it is known that we can view it as the inverse of a compact self-adjoint operator.

Spectral theorem of elliptic operators (Bonus)

The operator $-\Delta - 2|x|^{-1}$ is an example of what we call *elliptic operators*. We will not deal with this, but:

The operator $-\Delta - 2|x|^{-1}$ is unbounded, but it is known that we can view it as the inverse of a compact self-adjoint operator.



The eigenvalues are distributed like

$$0 < \lambda_1 < \lambda_2 < \cdots \rightarrow \infty.$$

Contents

Hydrogen atom

Spectral theory on Hilbert spaces

Gelfand theory

Algebraic geometry

C^* -algebras

First we give definitions:

Pseudo-definition

An algebra is a vector space with multiplication.

Definition

A C^* -algebra is a complex associative algebra with norm $\| \cdot \|$ such that:

1. $\|x^*x\| = \|x\|^2$

Example 1 : Bounded operators

Example 2 : Continuous functions

Spectra, multiplicative homomorphisms, maximal ideals

Gelfand-Naimark theorem

Algebraic variety

Coordinate ring

Maximal ideal is a point

Problem of unified codomains

Functoriality