

# Diachrony of Spectra

Ikhan Choi

Postech - Unist - Kaist Joint Seminar

July 25, 2019

# Introduction

# Introduction

## Definition

Let  $R$  be a commutative ring. The *spectrum* of  $R$  is the set of prime ideals of  $R$ .

# Introduction

## Definition

Let  $R$  be a commutative ring. The *spectrum* of  $R$  is the set of prime ideals of  $R$ .

## Example

$$\operatorname{Spec}(\mathbb{Z}) = \{2\mathbb{Z}, 3\mathbb{Z}, 5\mathbb{Z}, 7\mathbb{Z}, 11\mathbb{Z}, \dots\}.$$

# Introduction

## Definition

Let  $R$  be a commutative ring. The *spectrum* of  $R$  is the set of prime ideals of  $R$ .

## Example

$$\operatorname{Spec}(\mathbb{Z}) = \{2\mathbb{Z}, 3\mathbb{Z}, 5\mathbb{Z}, 7\mathbb{Z}, 11\mathbb{Z}, \dots\}.$$

## Question

Why is it defined like this?

# Contents

Hydrogen atom

Spectral theory on Hilbert spaces


Gelfand theory

Algebraic geometry

# Hydrogen spectral series



# Hydrogen spectral series



410.2nm


468.1nm

656.3 nm

434.0nm



# Hydrogen spectral series



410.2nm

468.1nm

656.3 nm

434.0nm

## Question

How can we explain and compute this phenomenon?

# Rydberg's formula : Bohr model

Bohr's postulates:

# Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.

# Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.
- ▶ The stationary orbits are computed by the old quantization assumption for angular momenta:

$$mvr = n\hbar.$$

# Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.
- ▶ The stationary orbits are computed by the old quantization assumption for angular momenta:

$$mvr = n\hbar.$$

- ▶ An electron absorbs or emits light frequency  $f$  when they jump from an orbit to another, satisfying

$$\Delta E = hf.$$

# Rydberg's formula : Bohr model

Bohr's postulates:

- ▶ The electrons are on certain stable orbits.
- ▶ The stationary orbits are computed by the old quantization assumption for angular momenta:

$$mvr = n\hbar.$$

- ▶ An electron absorbs or emits light frequency  $f$  when they jump from an orbit to another, satisfying

$$\Delta E = hf.$$

The constant  $h$  is called the Planck constant and  $\hbar := \frac{h}{2\pi}$ .

# Rydberg's formula : Bohr model

From the three relations

$$mvr = n\hbar, \quad \frac{mv^2}{r} = -k \frac{(+e)(-e)}{r^2}, \quad E = K + V = \frac{1}{2}mv^2 - k \frac{e^2}{r},$$

# Rydberg's formula : Bohr model

From the three relations

$$mvr = n\hbar, \quad \frac{mv^2}{r} = -k \frac{(+e)(-e)}{r^2}, \quad E = K + V = \frac{1}{2}mv^2 - k \frac{e^2}{r},$$

we deduce

$$E = -\frac{k^2 e^4 m}{2\hbar^2} \frac{1}{n^2} \approx -13.6 \frac{1}{n^2} \text{ (eV)}.$$



# Rydberg's formula : Bohr model

From the three relations

$$mvr = n\hbar, \quad \frac{mv^2}{r} = -k \frac{(+e)(-e)}{r^2}, \quad E = K + V = \frac{1}{2}mv^2 - k \frac{e^2}{r},$$

we deduce

$$E = -\frac{k^2 e^4 m}{2\hbar^2} \frac{1}{n^2} \approx -13.6 \frac{1}{n^2} \text{ (eV)}.$$

## Proposition (Rydberg formula)

*The wavelengths  $\lambda$  of absorbed or emitted photons from a hydrogen atom is estimated by the following formula:*

$$\frac{1}{\lambda} = R \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right), \quad \text{for } n_1, n_2 \in \mathbb{N},$$

where  $R := \frac{k^2 e^4 m}{4\pi\hbar^3 c}$  is the Rydberg constant.

# Rydberg's formula : Schrödinger equation

More mathematically!

# Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for  $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

# Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for  $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

# Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for  $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where  $V$  is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

# Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for  $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where  $V$  is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

By solving it, we obtain the probability distribution  $|\Psi(t, x)|^2$  of the electron at time  $t$ .

# Rydberg's formula : Schrödinger equation

More mathematically!

In quantum mechanics, an electron around a hydrogen atom is described by the Schrödinger equation: for  $(t, x) \in \mathbb{R}^{1+3}$

$$i\hbar \frac{\partial}{\partial t} \Psi(t, x) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x),$$

energy

kinetic energy

potential energy

where  $V$  is given by the Coulomb potential

$$V(x) = -\frac{2}{|x|}.$$

By solving it, we obtain the probability distribution  $|\Psi(t, x)|^2$  of the electron at time  $t$ . Let's solve.

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$



# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form  $\Psi(t, x) = \phi(t)\psi(x)$ , then

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form  $\Psi(t, x) = \phi(t)\psi(x)$ , then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)}$$

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form  $\Psi(t, x) = \phi(t)\psi(x)$ , then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form  $\Psi(t, x) = \phi(t)\psi(x)$ , then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

The constant  $E$  is interpreted as the energy of electrons.

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form  $\Psi(t, x) = \phi(t)\psi(x)$ , then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

The constant  $E$  is interpreted as the energy of electrons. We have two *eigenvalue problems* with *shared eigenvalue*  $E$ :

# Rydberg's formula : Schrödinger equation

Schrödinger equation:

$$i\hbar \frac{\partial \Psi(t, x)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(t, x) + V(x) \Psi(t, x).$$

“Mathematization”:

$$i\partial_t \Psi(t, x) = (-\Delta + V(x)) \Psi(t, x).$$

Ansatz: if the solutions has the form  $\Psi(t, x) = \phi(t)\psi(x)$ , then

$$\frac{i\partial_t \phi(t)}{\phi(t)} = \frac{(-\Delta + V(x))\psi(x)}{\psi(x)} = E.$$

The constant  $E$  is interpreted as the energy of electrons. We have two *eigenvalue problems* with *shared eigenvalue*  $E$ :

$$\begin{aligned} i\frac{d}{dt}\phi(t) &= E\phi(t), \\ (-\Delta + V(x))\psi(x) &= E\psi(x). \end{aligned}$$

## Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions  $\phi_E(t)$ ,  $\psi_E(x)$  of the eigenvalue problems for each complex number  $E$ .



## Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions  $\phi_E(t)$ ,  $\psi_E(x)$  of the eigenvalue problems for each complex number  $E$ .

- ▶ All functions of the form

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation (in fact we can find all solutions in this way).

## Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions  $\phi_E(t)$ ,  $\psi_E(x)$  of the eigenvalue problems for each complex number  $E$ .

- ▶ All functions of the form

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation (in fact we can find all solutions in this way).

- ▶ For a given  $E$ ,  $\phi_E$  and  $\psi_E$  are of course not unique. In fact they form a vector space which is called the eigenspace.

## Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions  $\phi_E(t)$ ,  $\psi_E(x)$  of the eigenvalue problems for each complex number  $E$ .

- ▶ All functions of the form

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation (in fact we can find all solutions in this way).

- ▶ For a given  $E$ ,  $\phi_E$  and  $\psi_E$  are of course not unique. In fact they form a vector space which is called the eigenspace.
- ▶ Note that for some  $E$  we probably cannot find the solution  $\psi_E(x)$  that satisfies  $\int |\psi_E(x)|^2 dx = 1$ : the eigenspace is trivial.

## Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions  $\phi_E(t)$ ,  $\psi_E(x)$  of the eigenvalue problems for each complex number  $E$ .

- ▶ All functions of the form

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation (in fact we can find all solutions in this way).

- ▶ For a given  $E$ ,  $\phi_E$  and  $\psi_E$  are of course not unique. In fact they form a vector space which is called the eigenspace.
- ▶ Note that for some  $E$  we probably cannot find the solution  $\psi_E(x)$  that satisfies  $\int |\psi_E(x)|^2 dx = 1$ : the eigenspace is trivial.
- ▶ Since  $\phi_E(t) \propto e^{-iEt}$  is easily solved, the main difficulty is  $\psi_E$ .

## Rydberg's formula : Schrödinger equation

Suppose we already have found the solutions  $\phi_E(t)$ ,  $\psi_E(x)$  of the eigenvalue problems for each complex number  $E$ .

- ▶ All functions of the form

$$\Phi(t, x) = \sum_E c_E \psi_E(t) \phi_E(x)$$

are solutions of the original Schrödinger equation (in fact we can find all solutions in this way).

- ▶ For a given  $E$ ,  $\phi_E$  and  $\psi_E$  are of course not unique. In fact they form a vector space which is called the eigenspace.
- ▶ Note that for some  $E$  we probably cannot find the solution  $\psi_E(x)$  that satisfies  $\int |\psi_E(x)|^2 dx = 1$ : the eigenspace is trivial.
- ▶ Since  $\phi_E(t) \propto e^{-iEt}$  is easily solved, the main difficulty is  $\psi_E$ .

### Remark

The first one is not mathematically correct statement because we should resolve some technical issues on convergence.

# Rydberg's formula : Schrödinger equation

So, what we need to investigate seriously is:

# Rydberg's formula : Schrödinger equation

So, what we need to investigate seriously is: *what are the eigenvalues and eigenvectors of the operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

*defined by*

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

*Also, how can we compute them?*

# Rydberg's formula : Schrödinger equation

So, what we need to investigate seriously is: *what are the eigenvalues and eigenvectors of the operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

*defined by*

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

*Also, how can we compute them?*

## Remark

Simply,  $L^2(\mathbb{R}^3)$  is the space of  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  such that  $\int |f|^2 < \infty$ .



# Rydberg's formula : Schrödinger equation

So, what we need to investigate seriously is: *what are the eigenvalues and eigenvectors of the operator*

$$\mathcal{H} : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$$

*defined by*

$$\mathcal{H}\psi(x) := (-\Delta - 2|x|^{-1})\psi(x)?$$

*Also, how can we compute them?*

## Remark

Simply,  $L^2(\mathbb{R}^3)$  is the space of  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  such that  $\int |f|^2 < \infty$ .

## The Beginning of Spectral Theory

# Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts found the following result:

# Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts found the following result:

## Proposition

*The eigenvalues of  $\mathcal{H} = -\Delta - 2|x|^{-1}$  are*

# Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts found the following result:

## Proposition

*The eigenvalues of  $\mathcal{H} = -\Delta - 2|x|^{-1}$  are*

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

# Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts found the following result:

## Proposition

*The eigenvalues of  $\mathcal{H} = -\Delta - 2|x|^{-1}$  are*

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

*with multiplicity 1, 4, 9, 16 and so on.*

# Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts found the following result:

## Proposition

*The eigenvalues of  $\mathcal{H} = -\Delta - 2|x|^{-1}$  are*

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

*with multiplicity 1, 4, 9, 16 and so on.*

Since the eigenvalues embody the possible energy values of an electron, we just gave the Rydberg formula a reasonable explanation.

# Rydberg's formula : Schrödinger equation

With long long calculations and hard hard mathematics, experts found the following result:

## Proposition

*The eigenvalues of  $\mathcal{H} = -\Delta - 2|x|^{-1}$  are*

$$-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, \dots$$

*with multiplicity 1, 4, 9, 16 and so on.*

Since the eigenvalues embody the possible energy values of an electron, we just gave the Rydberg formula a reasonable explanation. This result explains not only the discretized energy spectrum but also the number of each orbitals!

# Contents

Hydrogen atom

Spectral theory on Hilbert spaces

Gelfand theory

Algebraic geometry



# Separation of variables

# Spectral theorem of normal matrices

# Spectral theorem of compact operators

# Spectral theorem of elliptic operators

# Contents

Hydrogen atom

Spectral theory on Hilbert spaces

Gelfand theory

Algebraic geometry

# Banach algebras and $C^*$ -algebras

## Example 1 : Bounded operators

## Example 2 : Continuous functions



# Spectra, multiplicative homomorphisms, maximal ideals

# Gelfand-Naimark theorem

# Algebraic variety

# Coordinate ring

Maximal ideal is a point

# Problem of unified codomains

# Functoriality