

VLASOV-POISSON SYSTEM

IKHAN CHOI

CONTENTS

1. Vlasov-Poisson system	1
1.1. Representation of electric field	1
1.2. Volume preservation	2
1.3. Conservative laws	2
1.4. Moment propagation	3
2. Local existence	4
2.1. Approximate solution	4
2.2. A priori estimates of ρ_n and E_n	4
2.3. A priori estimates of $\nabla_x \rho_n$ and $\nabla_x E_n$	5
2.4. Convergence of f_n	5
2.5. Uniqueness	6
3. Global existence	7
3.1. Prolongation criterion	7
3.2. Some observations	7
3.3. Divide and conquer	8
3.4. Polynomial decay	9

1. VLASOV-POISSON SYSTEM

Consider a Cauchy problem of the *Vlasov-Poisson system*:

$$\begin{cases} f_t + v \cdot \nabla_x f + \gamma E \cdot \nabla_v f = 0, & (t, x, v) \in \mathbb{R}_t^+ \times \mathbb{R}_x^3 \times \mathbb{R}_v^3, \\ E(t, x) = -\nabla_x \Phi, \\ \Phi(t, x) = (-\Delta_x)^{-1} \rho, \\ \rho(t, x) = \int f dv, \\ f(0, x, v) = f_0(x, v), \end{cases}$$

where $\gamma = \pm 1$. For example, we have *repulsive problem* $\gamma = +1$ for electrons in plasma and *attractive problems* $\gamma = -1$ for galaxies. (ρ denotes mass density.)

1.1. Representation of electric field. For the boundaryless problem in which the potential function vanishes at infinity, we have

$$\Phi = \frac{1}{4\pi|x|} * \rho,$$

$$E = -\nabla_x \Phi = -\nabla_x \left(\frac{1}{4\pi|x|} * \rho \right) = \frac{x}{4\pi|x|^3} * \rho.$$

$$E(t, x) = \frac{1}{4\pi} \int \frac{(x - y)f(t, y, v)}{|x - y|^3} dy.$$

1.2. Volume preservation.

1.3. Conservative laws. The following results cannot be used in any local existence proof because they assume the existence of solutions, but they help understand the fundamental nature of solutions of the Vlasov-Poisson system and are used in the proof of global existence.

Lemma 1.1. *Let $f \in C_c^1([0, T] \times \mathbb{R}^6)$.*

(1) *(Continuity equation)*

$$\rho_t + \nabla_x \cdot j = 0, \quad \text{where } j = \int v f dv.$$

(2) *(Energy conservation)*

$$\iint |v|^2 f dv dx + \gamma \int |E|^2 dx = \text{const}.$$

Proof.

(1) Integrate with respect to v to get

$$\begin{aligned} 0 &= \int f_t dv + \int v \cdot \nabla_x f dv \\ &= \rho_t + \nabla_x \cdot \int v f dv \\ &= \rho_t + \nabla_x \cdot j. \end{aligned}$$

(2) Multiply $|v|^2$ and integrate with respect to v and x to get

$$\begin{aligned} \frac{d}{dt} \iint |v|^2 f dv dx &= \iint |v|^2 f_t dv dx = - \iint |v|^2 \gamma E \cdot \nabla_v f dv dx \\ &= \iint 2v \cdot \gamma E f dv dx = -2\gamma \int \nabla_x \Phi \cdot j dx \\ &= 2\gamma \int \Phi \nabla_x \cdot j dx = 2\gamma \int \Phi \Delta_x \Phi_t dx \\ &= -\frac{d}{dt} \gamma \int |E|^2 dx. \end{aligned}$$

Thus

$$\iint |v|^2 f dv dx + \gamma \int |E|^2 dx = \text{const}.$$

□

1.4. Moment propagation. We have a bound of kinetic energy even for $\gamma = -1$.

Lemma 1.2 ($L^{5/3}$ estimate of ρ).

- (1) $\|\rho(t)\|_{L_x^{5/3}} \lesssim \iint |v|^2 f \, dv \, dx$.
- (2) $\iint |v|^2 f \, dv \, dx \lesssim 1$.

Proof.

(1) Note

$$\begin{aligned} \rho(t, x) &= \int f(t, x, v) \, dv \leq \int_{|v| < R} f \, dv + \frac{1}{R^2} \int_{|v| \geq R} |v|^2 f \, dv \\ &\lesssim R^3 + R^{-2} \int |v|^2 f \, dv. \end{aligned}$$

Set $R^3 = R^{-2} \int |v|^2 f \, dv$ to get

$$\rho(t, x)^{5/3} \lesssim \int |v|^2 f \, dv.$$

(2) By the Hardy-Littlewood-Sobolev inequality,

$$\frac{1}{p} + \frac{\alpha}{d} = \frac{1}{q}$$

for $p = 2$, $d = 3$, and $\alpha = 1$ implies $q = 6/5$, hence the bound of $\|E(t)\|_2$

$$\|E(t)\|_2 \simeq \left\| \frac{1}{|x|^{d-\alpha}} * \rho(t, x) \right\|_{L_x^2} \lesssim \|\rho(t)\|_{6/5}.$$

So, interpolation with Hölder's inequality gives

$$\|E(t)\|_2 \lesssim \|\rho\|_1^{7/12} \|\rho\|_{5/3}^{5/12} \simeq \|\rho\|_{5/3}^{5/12}.$$

Thus (1) gives

$$\iint |v|^2 f \, dv \, dx = C + \|E(t)\|_2^2 \lesssim C + \left(\iint |v|^2 f \, dv \, dx \right)^{1/2},$$

so the kinetic energy $\iint f \, dv \, dx$ is bounded. As a corollary, $\|\rho\|_{5/3}$ is also bounded. \square

Lemma 1.3 (L^∞ -estimate of E). For $0 \leq \frac{1}{p} < \frac{\alpha}{d} < \frac{1}{q} \leq 1$,

$$\left\| \frac{1}{|x|^{d-\alpha}} * \rho(t, x) \right\|_{L_x^\infty}^{\frac{1}{q} - \frac{1}{p}} \lesssim \|\rho(t)\|_p^{\frac{1}{q} - \frac{\alpha}{d}} \|\rho(t)\|_q^{\frac{\alpha}{d} - \frac{1}{p}}.$$

Proof. Fix time t . For $(d - \alpha)p < d < (d - \alpha)q$,

$$\begin{aligned}
\left| \frac{1}{|x|^{d-\alpha}} * \rho(t, x) \right| &\leq \int_{|x-y| < R} \frac{\rho(t, y)}{|x-y|^{d-\alpha}} dy + \int_{|x-y| \geq R} \frac{\rho(t, y)}{|x-y|^{d-\alpha}} dy \\
&\leq \|\rho\|_{p'} \left(\int_{|y| < R} \frac{dy}{|y|^{(d-\alpha)p}} \right)^{1/p} + \|\rho\|_{q'} \left(\int_{|y| \geq R} \frac{dy}{|y|^{(d-\alpha)q}} \right)^{1/q} \\
&\simeq \|\rho\|_{p'} \left(\int_0^R r^{d-1-(d-\alpha)p} dr \right)^{1/p} + \|\rho\|_{q'} \left(\int_R^\infty r^{d-1-(d-\alpha)q} dr \right)^{1/q} \\
&\simeq \|\rho\|_{p'} R^{\frac{d}{p}-d+\alpha} + \|\rho\|_{q'} R^{\frac{d}{q}-d+\alpha}.
\end{aligned}$$

By choosing R such that $\|\rho\|_{p'} R^{\frac{d}{p}-d+\alpha} = \|\rho\|_{q'} R^{\frac{d}{q}-d+\alpha}$, we get

$$\|E(t)\|_\infty \lesssim \|\rho(t, x)\|_{p'}^{\frac{1-\frac{\alpha}{d}-\frac{1}{q}}{\frac{1}{p}-\frac{1}{q}}} \|\rho(t, x)\|_{q'}^{\frac{\frac{1}{p}-1+\frac{\alpha}{d}}{\frac{1}{p}-\frac{1}{q}}},$$

hence the inequality by interchanging p and q with their conjugates. \square

2. LOCAL EXISTENCE

2.1. Approximate solution. Suppose $f_0 \in C_c^1$. With initial term f_0 , inductively define f_{n+1} by letting the electric field E be known:

$$\begin{cases} \partial_t f_{n+1} + v \cdot \nabla_x f_{n+1} + \gamma E_n \cdot \nabla_v f_{n+1} = 0, \\ E_n(t, x) = -\nabla_x \Phi_n, \\ \Phi_n(t, x) = (-\Delta_x)^{-1} \rho_n, \\ \rho_n(t, x) = \int f_n dv, \\ f_{n+1}(0, x, v) = f_0(x, v). \end{cases}$$

Characteristics.

Then, the equation becomes a linear PDE for f_{n+1} .

Definition 2.1.

$$Q_n(t) = \sup\{|v| : f(s, x, v) \neq 0, s \in [0, t]\}$$

2.2. A priori estimates of ρ_n and E_n .

Lemma 2.1.

- (1) For all $t \in [0, \infty]$, $\|\rho_n(t)\|_\infty \lesssim Q_n^3(t)$ and $\|E_n(t)\|_\infty \lesssim Q_n^2(t)$
- (2) For sufficiently small $t < c(f_0)$, $Q_n(t) \lesssim 1$ independent on n .

Proof.

(1) Since

$$\|\rho_n(t)\|_\infty \leq Q^3(t) \|f_0\|_\infty \lesssim Q_n^3(t),$$

a rough estimate for $\|E\|_\infty$ gives

$$\|E_n(t)\|_\infty \leq \|\rho_n(t)\|_\infty^{2/3} \|\rho_n(t)\|_1^{1/3} \lesssim Q_n^2(t)$$

(2) Let $c = c(f_0)$ be a constant such that $\|E_n(t)\| \leq cQ^2(t)$. We claim that

$$Q_n(t) \leq \frac{Q_0}{1 - cQ_0 t}$$

for all n . Easily checked for $n = 0$; $Q_0(t) = Q_0 \leq \frac{Q_0}{1 - cQ_0 t}$.

Assume $Q_n(t) \leq \frac{Q_0}{1 - cQ_0 t}$. Then,

$$\begin{aligned} |V_{n+1}(s; 0, x, v)| &\leq |v| + \int_0^s |E_n(s; 0, x, v)| ds \\ &\leq Q_0 + c \int_0^s Q_n^2(s) ds \end{aligned}$$

implies

$$\begin{aligned} Q_{n+1}(t) &\leq Q_0 + c \int_0^t Q_n^2(s) ds \\ &\leq Q_0 + c \int_0^t \left(\frac{Q_0}{1 - cQ_0 s} \right)^2 ds = \frac{Q_0}{1 - cQ_0 t}. \end{aligned}$$

By induction, $Q_n(t) \lesssim 1$ for all n and $t \in [0, T]$, where $T < (cQ_0)^{-1}$. \square

2.3. A priori estimates of $\nabla_x \rho_n$ and $\nabla_x E_n$.

2.4. Convergence of f_n .

(1)

$$\|f_{n+1}(t) - f_n(t)\|_\infty \lesssim \int_0^t \|E_n(s) - E_{n-1}(s)\|_\infty ds.$$

(2)

$$\|f_{n+1}(t) - f_n(t)\|_\infty \lesssim \int_0^t \|f_n(s) - f_{n-1}(s)\|_\infty^{1/3} ds.$$

Therefore, f_n converges uniformly.

Proof.

(1) Take T such that

$$T < \min\{\|\nabla_x E_n(t)\|_\infty^{-1}, 1\} \quad \text{and} \quad Q_n(t) \lesssim 1$$

for all n and $t < T$. This can be done thanks for the local a priori bound for $\nabla_x E_n$.

The C^1 regularity of f_0 lets

$$\begin{aligned} &|f_{n+1}(t, x, v) - f_n(t, x, v)| \\ &= |f_0(X_{n+1}(0; t, x, v), V_{n+1}(0; t, x, v)) - f_0(X_n(0; t, x, v), V_n(0; t, x, v))| \\ &\lesssim |X_{n+1}(0; t, x, v) - X_n(0; t, x, v)| + |V_{n+1}(0; t, x, v) - V_n(0; t, x, v)|. \end{aligned}$$

Because

$$\begin{aligned} X_n(s; t, x, v) &= x - \int_s^t V_n(s'; t, x, v) ds', \\ V_n(s; t, x, v) &= v - \int_s^t E_{n-1}(s', X_n(s; t, x, v)) ds', \end{aligned}$$

we have

$$\begin{aligned} &|V_{n+1}(s; t, x, v) - V_n(s; t, x, v)| \\ &\leq \int_0^t |E_n(s, X_{n+1}(s; t, x, v)) - E_{n-1}(s, X_n(s; t, x, v))| ds \\ &\leq \int_0^t |E_n(s, X_{n+1}) - E_n(s, X_n)| + |E_n(s, X_n) - E_{n-1}(s, X_n)| ds \\ &\leq T \sup_{s \in [0, t]} \|\nabla_x E_n(s)\|_\infty |X_{n+1}(s) - X_n(s)| + \int_0^t \|E_n(s) - E_{n-1}(s)\|_\infty ds \end{aligned}$$

and

$$\begin{aligned} |X_{n+1}(s; t, x, v) - X_n(s; t, x, v)| &\leq \int_0^t |V_{n+1}(s; t, x, v) - V_n(s; t, x, v)| ds \\ &\leq T \sup_{s \in [0, t]} |V_{n+1}(s) - V_n(s)|. \end{aligned}$$

Since T is taken such that

$$T < \min\{\|\nabla E_n(t)\|_\infty^{-1}, 1\},$$

we get by combining the above two inequality,

$$\begin{aligned} &\sup_{s \in [0, t]} |X_{n+1}(s; t, x, v) - X_n(s; t, x, v)| + |V_{n+1}(s; t, x, v) - V_n(s; t, x, v)| \\ &\lesssim \int_0^t \|E_n(s) - E_{n-1}(s)\|_\infty ds. \end{aligned}$$

(2) Notice that

$$\|E_n(t) - E_{n-1}(t)\|_\infty \lesssim \|\rho_n(t) - \rho_{n-1}(t)\|_1^{1/3} \|\rho_n(t) - \rho_{n-1}(t)\|_\infty^{2/3}.$$

For L^∞ -norm,

$$\|\rho_n(t) - \rho_{n-1}(t)\| \lesssim Q_n^3(t) + Q_{n-1}^3(t) \lesssim 1.$$

For L^1 -norm, since the support of f_n, f_{n-1} is bounded in both directions x, v ,

$$\|\rho_n(t) - \rho_{n-1}(t)\|_1 \leq \|f_n(t) - f_{n-1}(t)\|_1 \lesssim \|f_n(t) - f_{n-1}(t)\|_\infty.$$

□

2.5. Uniqueness.

3. GLOBAL EXISTENCE

3.1. Prolongation criterion.

Theorem (Schaeffer, 1991). *Let $f_0 \in C_{c,x,v}^1$ and $f_0 \geq 0$. Then, the Cauchy problem for the VP system has a unique C^1 global solution.*

Definition 3.1. For a local solution f ,

$$Q(t) := 1 + \sup\{|v| : f(s, x, v) \neq 0 \text{ for some } s \in [0, t], x \in \mathbb{R}_x^3\}.$$

Decompose $[t - \Delta, t] \times \mathbb{R}_x^3 \times \mathbb{R}_v^3$ as

$$\begin{aligned} U &= \left\{ (s, x, v) : |v - \widehat{V}(t)| \geq P, \quad |y - \widehat{X}(s)| \geq r \right\}, \\ B &= \left\{ (s, x, v) : |v - \widehat{V}(t)| \geq P, \quad |v| \geq P \right\} \setminus U, \\ G &= \left\{ (s, x, v) : |v - \widehat{V}(t)| < P \quad \text{or} \quad |v| < P \right\}. \end{aligned}$$

(We can let $U \mapsto U \cap \{|v| \geq P\}$ to make the decomposition disjoint.) Later we choose

$$P = Q^{4/11}, \quad r = R \max\{|v|^{-3}, |v - \widehat{V}(t)|^{-3}\}, \quad R = Q^{16/33} \log^{1/2} Q.$$

Also, later we choose $\Delta \cdot \sup_{s \leq t} \|E(s)\|_\infty < \frac{P}{4}$.

3.2. Some observations. Our goal is to obtain a priori estimate like

$$\|E(t)\|_\infty \lesssim Q(t)^a \quad \text{for some } a < 1.$$

Since the force field E measures the maximal rate of changes in velocity, the estimate can be read very roughly as

$$Q'(t) \lesssim Q(t)^a,$$

which lead its polynomial growth.

So we need to bound the Riesz potential E . The following observation suggests a lower bound of relative velocity.

Claim. *Fix t, x, v . If $|v - \widehat{V}(t)| \geq P$, then*

$$|y - \widehat{X}(s)| \geq \frac{1}{4} |v - \widehat{V}(t)| |s - s_0|$$

for some $s_0 \in [t - \Delta, t]$, where $\Delta \cdot \sup_{s \leq t} \|E(s)\|_\infty < \frac{P}{4}$.

Proof. Since $\Delta \|E(s)\|_\infty < \frac{P}{4}$, we have

$$|v - w| < \frac{P}{4} \quad \text{and} \quad |\widehat{V}(t) - \widehat{V}(s)| < \frac{P}{4}.$$

The condition $|v - \widehat{V}(t)| \geq P$ implies

$$\frac{1}{2} |v - \widehat{V}(t)| \leq |v - \widehat{V}(t)| - \frac{P}{4} - \frac{P}{4} < |w - \widehat{V}(s)|.$$

Let $Z(s) := y - \widehat{X}(s)$. Then,

$$\begin{aligned} Z'(s) &= w - \widehat{V}(s), \\ Z''(s) &= \gamma[E(s, y, w) - E(s, \widehat{X}(s), \widehat{V}(s))]. \end{aligned}$$

Let $s_0 \in [t - \Delta, t]$ minimize $s \mapsto |Z(s)|$ and expand Z as

$$Z(s) = Z(s_0) + Z'(s_0)(s - s_0) + \frac{Z''(\sigma)}{2}(s - s_0)^2$$

for some σ between s and s_0 . Then,

$$|Z(s_0) + Z'(s_0)(s - s_0)| \geq |Z'(s_0)(s - s_0)| \geq \frac{1}{2}|v - \widehat{V}(t)||s - s_0|$$

and

$$\begin{aligned} \left| \frac{Z''(\sigma)}{2}(s - s_0)^2 \right| &\leq \|E(t)\|_\infty (s - s_0)^2 \leq \|E(t)\|_\infty \Delta |s - s_0| \\ &\leq \frac{P}{4}|s - s_0| \leq \frac{1}{4}|v - \widehat{V}(t)||s - s_0| \end{aligned}$$

proves

$$|y - \widehat{X}(s)| = |Z(s)| \geq \frac{1}{4}|v - \widehat{V}(t)||s - s_0|. \quad \square$$

We introduce time averaging to use the above lower bound.

Claim. Fix t, x, v . If $|v - \widehat{V}(t)| \geq P$, then

$$\int_{t-\Delta}^t \frac{1}{|y - \widehat{X}(s)|^2} \chi_A(s) ds \lesssim \frac{r^{-1}}{|v - \widehat{V}(t)|},$$

where $A = \{s : |y - \widehat{X}(s)| \geq r\}$.

Proof. Since $|y - \widehat{X}(s)| \geq \frac{1}{4}|v - \widehat{V}(t)||s - s_0|$,

$$\begin{aligned} \int_{t-\Delta}^t \frac{1}{|y - \widehat{X}(s)|^2} \chi_A(s) ds &\leq 16 \int_{t-\Delta}^t \frac{1}{|v - \widehat{V}(t)|^2 |s - s_0|^2} \chi_A(s) ds \\ &\leq 32 \int_r^\infty \frac{1}{|v - \widehat{V}(t)|^3 |s - s_0|^2} d(|v - \widehat{V}(t)||s - s_0|) \\ &= 32 \frac{r^{-1}}{|v - \widehat{V}(t)|}. \quad \square \end{aligned}$$

3.3. Divide and conquer.

3.3.1. *Ugly set estimate.* Therefore, if we let $r^{-1} \simeq \min\{|v|^3, |v - \widehat{V}(t)|^3\}$, then

$$\int_{t-\Delta}^t \frac{1}{|y - \widehat{X}(s)|^2} \chi_A(s) ds \lesssim |v|^2$$

so that we have

$$\iiint_U \frac{f(s, y, w)}{|y - \widehat{X}(s)|^2} dw dy ds \lesssim R^{-1} \int |v|^2 f(t, x, v) dv dx \lesssim R^{-1}$$

when

$$U \subset \{(s, x, v) : |v - \widehat{V}(t)| \geq P, \quad |y - \widehat{X}(s)| \geq R \max\{|v|^{-3}, |v - \widehat{V}(t)|^{-3}\}\}.$$

3.3.2. *Bad set estimate.* Consider U^c . We need to control the union of two regions

$$|y - \widehat{X}(s)| < R|v|^{-3} \quad \text{and} \quad |y - \widehat{X}(s)| < R|v - \widehat{V}(t)|^{-3}.$$

Without any conditions, the integration of fundamental solution with respect to y gives

$$\int_{|y - \widehat{X}(s)| < r} \frac{1}{|y - \widehat{X}(s)|^2} dy \simeq r.$$

Claim. If $|v| \geq P$ and $|v - \widehat{V}(t)| \geq P$, then

$$\int_{U^c} \frac{1}{|y - \widehat{X}(s)|^2} dy \lesssim \max\{|w|^{-3}, |w - \widehat{V}(s)|^{-3}\}$$

for $s \in [t - \Delta, t]$.

Proof. It follows from

$$|w| \simeq |v|, \quad |w - \widehat{V}(s)| \simeq |v - \widehat{V}(t)|$$

for $|v| \geq P$ and $|v - \widehat{V}(t)| \geq P$. □

3.3.3. *Good set estimate.*

3.4. Polynomial decay.

Lemma 3.1. *Along the time of existence we have*

$$\|E(t)\|_{L_x^\infty} \lesssim Q(t)^{4/3}.$$

Proof. Note that we have

$$\|E\|_\infty \lesssim \|\rho\|_\infty^{4/9} \|\rho\|_{5/3}^{5/9}.$$

Since the velocity support of f is bounded by finite $Q(t)$,

$$\rho(t, x) = \int_{|v| < Q(t)} f(t, x, v) dv \lesssim Q(t)^3 \|f_0(x)\|_{L_v^\infty} \lesssim Q(t)^3,$$

so

$$\|E(t)\|_{L_x^\infty} \lesssim \|\rho(t)\|_{L_x^\infty}^{4/9} \lesssim Q(t)^{4/3}. \quad \square$$