

Cardiac Advantages of **NT-proBNP** in Type 2 Diabetic Patients^{1, 2, 3}

2형 당뇨병 환자에게 HbA1c 검사 시 NT-proBNP도 함께 고려해주세요!



HbA1c



FPG



2-hour Postprandial Glucose



NT-proBNP

Grand

2형 당뇨병 환자는 심부전의 위험이 1.51배 증가¹



2형 당뇨병 환자는 심부전 입원 위험이 33% 증가⁴



ADA Diabetes Care 2024⁵



- 10.39a: Adults with diabetes are at increased risk for the development of asymptomatic cardiac structural or functional abnormalities (stage B heart failure) or symptomatic (stage C) heart failure. Consider screening adults with diabetes by

measuring a natriuretic peptide (B-type natriuretic peptide [BNP] or N-terminal pro-BNP [NT-proBNP])


to facilitate the prevention of stage C heart failure

- 10.39b: In asymptomatic individuals with diabetes and abnormal natriuretic peptide levels, echocardiography is recommended to identify stage B heart failure. A


NT-proBNP는 심장 사건 발생 6개월 전에 심부전 위험에 노출된 사람을 식별하는 데 도움^{6, 7, 8}



NT-proBNP는 2형 당뇨병 환자의 심혈관계 위험과 가장 높은 상관관계를 나타내는 바이오마커⁹



2형 당뇨병 환자의 심부전 증가 위험과 관련된 위험 요인¹⁰



Reference 1. Diabetes Fact Sheet in Korea 2022 2. Abrar Ahmad et al. Nature-Communication Medicine 4, 2024;11 3. Verma Set al. Curr Opin Cardiol. 2019;34:578-583; 4. Gerstein HC et al. Arch Intern Med 2008;168:1699-704. 5. American Diabetes Association Professional Practice Committee. Diabetes Care 2024;47(Supplement_1):S179-S218 6. Wolsk E et al. Circulation. 2017;136:1560-1562. 7. Magnusson M et al. Diabetes Care. 2004;27:1929-35; 8. Gallagher J et al. Cardiac Failure Review. 2017;3:83-85. 9. Abrar Ahmad et al. Nature-Communication Medicine 4, :11 (2024) 10. Verma S et al. Curr Opin Cardiol. 2019;34:578-583;

