

Recreating Young's Double Slit Experiment

Ed Lipchus*

Department of Natural Science, Hampshire College, Amherst, MA 01002

Danika Luntz-Martin[†]

Department of Physics, Smith College, Northampton, MA 01063

(Dated: December 9, 2014)

Abstract

I. INTRODUCTION

The double slit experiment performed by Thomas Young was seminal because it demonstrated the wave nature of light.

II. METHODS

We used the TeachSpin Two-Slit Interference, One Photon at a Time (TWS1-B) which is an apparatus designed to perform Young's double slit experiment. The apparatus has two light sources, a 670 nm diode laser and a light bulb with a removable green light filter. There are four slit holders spaced throughout the length of the apparatus, see Figure ?? . The first slit, the source slit is Additionally we used the TeachSpin Pulse Counter / Interval Timer (PCIT1)




FIG. 1. The experimental set-up that we used.

III. RESULTS

IV. ANALYSIS

V. DISCUSSION

VI. CONCLUSION

* ejl13@hampshire.edu

† dluntzma@smith.edu

¹ Jonathan F. Reichert, *TeachSpin Instruction Manuals: Two-Slit Interference, One Photon at a Time (TWS1 - B), Pulse Counter / Interval Timer (PCIT1)* Rev. 1.0, (2013)