

Introductory Electricity

Christian I. Cardozo-Aviles, David I. Mayo, and Phong T. Vo

Massachusetts Institute of Technology

HSSP Summer 2014

June 30, 2014

PROBLEM SET 1

This problem set is due **July 13th, 2014** if you would like to receive feedback on your work. The exercises below vary in difficulty. Some are straightforward computational exercises intended to hone essential skills accompanying concepts in the previous lectures. Others may require more critical thinking to develop a deeper appreciation for the topics introduced. If you are unable to make progress on any particular problem and would like to obtain some hints before the solutions are released, please feel free to email us with your request.

1 Lecture Summary:

In lecture on July 6th, 2014, we discussed the basics of *electrostatics* and some mathematical preliminaries.

1. Suppose we have functions $f(x)$ and $g(x)$ and constants C and D ; then the derivative of the sum of these two functions is

$$\frac{d[Cf(x) + Dg(x)]}{dx} = C \frac{df(x)}{dx} + D \frac{dg(x)}{dx}.$$

Likewise, the integral is

$$\int [Cf(x) + Dg(x)]dx = C \int f(x)dx + D \int g(x)dx.$$

2. If we have a function of the form $f(x) = x^n$ for some number n , then the derivative of this function is

$$\frac{d(x^n)}{dx} = nx^{n-1},$$

and its integral is

$$\int x^n dx = \frac{x^{n+1}}{n+1} + A,$$

for an arbitrary constant A if $n \neq -1$, and is

$$\int x^{-1} dx = \ln |x| + A,$$

for the case $n = -1$.

3. Given two vectors in component-form, $\mathbf{v} = \langle v_x, v_y, v_z \rangle = v_x \hat{i} + v_y \hat{j} + v_z \hat{k}$ and $\mathbf{w} = \langle w_x, w_y, w_z \rangle = w_x \hat{i} + w_y \hat{j} + w_z \hat{k}$, we can perform the following operations:
 - (a) Addition: $\mathbf{v} + \mathbf{w} = \langle v_x + w_x, v_y + w_y, v_z + w_z \rangle$,
 - (b) Subtraction: $\mathbf{v} - \mathbf{w} = \langle v_x - w_x, v_y - w_y, v_z - w_z \rangle$,
 - (c) Scalar multiplication: $c\mathbf{v} = \langle cv_x, cv_y, cv_z \rangle$ for some scalar c ,
 - (d) Norm: $\|\mathbf{v}\| = \sqrt{v_x^2 + v_y^2 + v_z^2}$,
 - (e) Inner (dot) product: $\mathbf{v} \cdot \mathbf{w} = v_x w_x + v_y w_y + v_z w_z = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta$, where θ is the smallest angle between the vectors,
 - (f) Vector (cross) product: $\mathbf{v} \times \mathbf{w} = \langle v_y w_z - v_z w_y, v_z w_x - v_x w_z, v_x w_y - v_y w_x \rangle$, and $\|\mathbf{v} \times \mathbf{w}\| = \|\mathbf{v}\| \|\mathbf{w}\| \sin \theta$, where θ is the smallest angle between the vectors,
 - (g) Unit vector: $\hat{\mathbf{v}} = \frac{\mathbf{v}}{\|\mathbf{v}\|}$.
4. There are two flavors of charges: plus (+) and minus (-). We take axiomatically that like charges repel and opposite charges attract. Charges are quantized to integral multiples of a third of the elementary charge $e = 1.60217657 \times 10^{-19}$ C.

5. Given two charges q_1 and q_2 , the force between them is given by *Coulomb's Law*

$$\mathbf{F} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2} \hat{\mathbf{r}} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^3} \mathbf{r},$$

where $\hat{\mathbf{r}}$ is the unit radial vector that joins the location of the two charges, r is the distance between the two charges, and $\epsilon_0 = 8.85418782 \times 10^{-12}$ F/m is the vacuum permittivity.

6. Coulomb's Law obeys the *superposition principle*. That is, the force on a charge due to the presence of many charges is simply the sum of the forces between that charge and each of the other charges individually. Suppose that we have a collection of N charges $\{q_1, q_2, q_3, \dots, q_N\}$. Then the force on q_1 is simply

$$\mathbf{F} = \mathbf{F}_2 + \mathbf{F}_3 + \dots + \mathbf{F}_N,$$

where the subscript i denotes the force between q_1 and q_i for $i = 2, 3, 4, \dots, N$.

2 References:

For the convenience of the readers, we provide links to third-party websites below. We are *not* responsible for any content contained in those websites. Please view them at your own discretion.

1. For a basic introduction to vectors (with some computational aides), please visit <http://hyperphysics.phy-astr.gsu.edu/hbase/vect.html>.
2. Those interested in learning more about differentiation and integration can consult any calculus textbook for more information. For instance, chapters 2 and 5 of *Calculus* by Gilbert Strang are relevant to our purpose. This textbook is free at <http://ocw.mit.edu/resources/res-18-001-calculus-online-textbook-spring-2005/textbook/>.
3. For a nice conceptual coverage of Coulomb's Law, please see sections 17.1 and 17.2 of *Physics* by Raymond A. Serway and Jerry S. Faughn.

4. For some practice with Coulomb's Law, please see http://dev.physicslab.org/Document.aspx?doctype=3&filename=Electrostatics_AdvancedCoulombsLawProblems.xml
5. To review Thomson's discovery of the electron, please visit http://www-outreach.phy.cam.ac.uk/camphy/electron/electron1_1.htm.
6. Those interested in learning more about the different atomic models are encouraged to search online for more information. The models we explored in class include the plum-pudding model, Rutherford's planetary model, Bohr's atomic model, and the Schrodinger's electron cloud model.

3 Exercises:

Exercise 1 – Differentiation and Integration

Find the derivative and integral of the following functions with respect to the specified independent variable. All letters appearing in these functions, other than the independent variable, are assumed to be constants.

1. $f(r) = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$
2. $f(x) = -kx$
3. $f(r) = \frac{l^2}{2mr^2} + \frac{GmM}{r}$

Exercise 2 – Working with Vectors, Part I

Given the following vectors, $\mathbf{a} = \langle 1, 0, 1 \rangle$ and $\mathbf{b} = \langle 2, 4, 3 \rangle$, find the following quantities:

1. $\|\mathbf{a}\|$ and $\|\mathbf{b}\|$
2. $\mathbf{a} + \mathbf{b}$
3. $\mathbf{a} - 6\mathbf{b}$
4. $\mathbf{a} \cdot 2\mathbf{b}$
5. $\mathbf{a} \times \mathbf{b}$
6. Find a vector which is both perpendicular to \mathbf{a} and \mathbf{b} by whatever means seems easiest.

Exercise 3 – Working with Vectors, Part II

Suppose we are working in the three-dimensional Cartesian coordinate system. Consider three points in the space: $P = (0, 0, 2)$, $Q = (0, 3, 0)$, and $R = (a, b, c)$, where a, b , and c are unknown coordinates to be determined.

1. Find the position vectors associated with each given point. That is, denote the origin as $O = (0, 0, 0)$. We want to find the vectors $\mathbf{P} = \overrightarrow{OP}$, $\mathbf{Q} = \overrightarrow{OQ}$, and $\mathbf{R} = \overrightarrow{OR}$.
2. Find the vector \overrightarrow{PQ} and its associated unit vector.
3. Now, we are interested in finding the point R subject to the following constraints: $\mathbf{P} \times \mathbf{R} = \mathbf{Q} \times \mathbf{R} = \mathbf{0}$ and $\|\mathbf{R}\| = 5$. Do these constraints produce a unique R ? If not, what other constraints would you need to obtain uniqueness?
4. Now, we repeat the previous step with a different set of constraints: $\mathbf{P} \times \mathbf{R} = \mathbf{0}$, $\mathbf{Q} \cdot \mathbf{R} = 2$ and $\|\mathbf{R}\| = 1$. Do these constraints produce a unique R ? If not, what other constraints would you need to obtain uniqueness?

Exercise 4 – Coulomb’s Law at Work

Consider two charges $q_1 = -e$ and $q_2 = e$ with masses m_1 and m_2 respectively.

1. Find the acceleration on m_1 as a function of the separation distance r .
2. Suppose the two masses are initially separated by a distance $r_0 = 5.29 \times 10^{-11}$ m. Find the acceleration on m_1 here.
3. Now, imagine that m_2 is much much larger than m_1 that it remains essentially at rest throughout the entire interaction. If the acceleration on m_1 remains the same as that when it is at a distance r_0 from m_2 , how much time does it take for m_1 to collide with m_2 ? Of course, this approximation is vastly inaccurate because the acceleration gets larger and larger as r goes to zero. However, it does serve to provide an upper limit for the collision time.

Exercise 5 – The Sum of Forces

Let $q_1 = q_2 = e$ be placed at $(0, 2)$ and $(0, -2)$ respectively. These two charges are pinned in place. A third charge $q_3 = -2e$ is placed originally at $(x_0, 0)$ for some x_0 .

1. Find the force, magnitude and direction, on q_3 as a function of the location of q_3 .
2. Qualitatively describe the motion of q_3 if it were free to move according to Newton’s laws.
3. Find the acceleration of q_3 as a function of its position using Newton’s second law.
4. Solve for the position of q_3 as a function of time. *Hint:* The acceleration is the second derivative of (i.e. the derivative of the derivative of) x with respect to time. So, you should arrive at an expression

$$\frac{d^2x}{dt^2} = -\omega^2x,$$

for some constant ω that you will determine. This *differential equation* has solutions

$$x(t) = A \cos(\omega t) + B \sin(\omega t),$$

for arbitrary constants A and B that are determined by initial conditions. In our problem, the initial conditions are that the charge start out at rest and is initially at x_0 on the x -axis. This reduces the solution to

$$x(t) = x_0 \cos(\omega t).$$

Does this solution confirm your intuition in part (b)?

Exercise 6 – Balancing charges

Consider a collection of four charges placed at the vertex of a square with sides of 2 m. Suppose that the charges are $q_1 = e$, $q_2 = -2e$, $q_3 = 3q$, and $q_4 = -4e$ arranged clockwise starting with the upper left vertex. Suppose that these four charges are somehow pinned down to where they are so that they cannot move by interacting with the other charges. We want to trap a fifth charge $q_5 = e$ inside the square. Where must q_5 be placed? *Hint:* This problem is not conceptually difficult (we believe), but it is computational cumbersome. So please do not become discouraged if you run into messy algebra. As a suggestion, it is crucial that you pay attention to the signs of the forces. If you cannot solve the algebra, it is sufficient to arrive at a set of equations that would, through some algebra (which is not the point of this course), solve for the location of q_5 .




Figure 1: The Balmer series of the hydrogen spectrum. This image is taken from http://chemwiki.ucdavis.edu/Under_Construction/chem1/Atoms_and_the_Periodic_Table/The_Bohr_Atom.

Exercise 7 – Spectrum of the Hydrogen

In class, we stated that the energy levels E_n of the hydrogen atom are quantized according to

$$E_n = \frac{-2.17896 \times 10^{-18} \text{ J}}{n^2}$$

for $n = 1, 2, 3, \dots$. Thus, an electron that falls from an initial energy level with quantum number n_i to a final energy level with quantum number n_f releases a photon of light with energy corresponding to the difference in energy between the two levels (the energy that the electron lost). Recall that $E = h\frac{c}{\lambda}$, where $h = 6.626 \times 10^{-34} \text{ Js}$ is Planck's constant, $c = 299,792,458 \text{ m/s}$ and λ is the wavelength of the emitted light. Calculate the wavelength of the emitted light for $n_f = 2$ and $n_i = 3, 4, \dots, 7$. These spectral lines are known as the *Balmer Series*. Compare your result with the data in Fig. 1. Agreement of the Bohr's model with experimental spectroscopic data was one of the early triumphs of the old quantum model.