Mercury / Vulcan and the id–iv Ladder

a Newtonian, measurement-only audit

Axiom accepted for the experiment: Vulcan is bound to Mercury as Charon to Pluto, always anti-Earth.
Every consequence computed from JPL DE421 and spacecraft/radar measurements; no relativity, no myth.

Deliverables: 6 Godot videos · 9 figures · full metric tables · refutation ledger
star_learning/solar_system_explorer/docs/MERCURY_VULCAN_LADDER_20261010/

2026-10-10

Mercury / Vulcan and the id–iv Ladder — a Newtonian, measurement-only audit

Date: 2026-10-10 · Workspace: star_learning/solar_system_explorer/docs/MERCURY_VULCAN_LADDER_20261010/
Tools: tools/vulcan/bodies.py, vulcan_derivation.py, ladder_analysis.py, ladder_null.py, build_report.py; Godot recorder game/tools/record_vulcan.gd; encoder tools/vulcan/encode_vulcan.sh.
Ephemeris: JPL DE421 via Skyfield. No relativistic term is used anywhere in this report. No myth is admitted as evidence. Every claim below is a measurement, a Newtonian derivation from measurements, or is labelled as the user's axiom.

0. Axioms accepted for the experiment, and the rules

- Axiom A (user): Vulcan is bound to Mercury the way Charon is bound to Pluto, and sits ALWAYS on the far side of Mercury from Earth — Config A: Earth → Moon → Mercury → Vulcan.
- Axiom B (user, provisional): when the Moon lies between Earth and Mercury the order may invert — Config B: Earth → Moon → Vulcan → Mercury — and this configuration is rare.
- Planet X: a placeholder for a pattern break demanded by the ladder; nothing else is assumed about it.
- Rule 1: Newtonian mechanics only (Kepler III, Hill sphere, Roche limit, barycentre, parallax). Rule 2: measured numbers from spacecraft/radar/helioseismology are admitted as measurements, not as theory. Rule 3: every test states what observation would overturn it.
- The Zecliptic: the workspace rule defines it as the ecliptic with Sol's equator 7.25° off it. The user asked for angles 'from the Zecliptic (Solar equator)'. Both frames are therefore tabulated: inclination to the ecliptic, to the solar equator, and to the invariable plane.

1. Data layer — every measurable metric

data/bodies_metrics.{json,csv,md} holds 43 records: Sol, the eight planets, Pluto/Charon and its four small moons, Luna, Phobos/Deimos, the Galilean moons, seven Saturnian moons, five Uranian, Triton/Proteus/Nereid, Ceres/Vesta/Pallas/Hygiea, the asteroid belt (AB) and the heliobarrier (HB). Per body: mass, GM, radii, density, a, e, i (three frames), node, sidereal and synodic periods, rotation, obliquity, J2, magnetic dipole, albedos, surface gravity, escape speed, irradiance, Hill radius, orbital and spin angular momentum, moon count, total moon mass, ring presence/mass/extent, Roche limit, barycentre offset, synchronous-orbit radius. Internal checks pass: $T^2/a^3 = 1.000 \pm 0.001$ for every planet; Charon's Kepler period from (a, M) is 6.385 d against the measured 6.387 d; Luna's is 27.28 d against 27.32 d; Mars' synchronous orbit comes out at 20,428 km (literature 20,428 km); the Earth–Luna barycentre at 4,671 km (literature 4,671 km).

Body	Mass kg	a AU	T d	Synodic d	Rot d	T/rot	Obliq °	i_ecl °	i_SolEq °	i_inv °	Moons	Rings	Ring kg	Hill km
Mercury	3.3e+23	0.3871	87.969	115.9	58.65	1.5	0.034	7.005	3.38	6.344	0	no	0	175,284
Venus	4.87e+24	0.7233	224.7	583.9	-243	0.9246	177.4	3.395	3.856	2.195	0	no	0	1,004,233
Earth	5.97e+24	1	365.26	0	0.9973	366.3	23.44	0	7.25	1.579	1	no	0	1,471,428
Mars	6.42e+23	1.524	686.98	779.9	1.026	669.6	25.19	1.85	5.649	1.68	2	no	0	982,738
Jupiter	1.9e+27	5.203	4,333	398.9	0.4135	10,477	3.13	1.304	6.09	0.327	95	yes	1e+13	50,562,816
Saturn	5.68e+26	9.537	10,759	378.1	0.444	24,232	26.73	2.485	5.504	0.93	274	yes	1.54e+19	61,645,286
Uranus	8.68e+25	19.19	30,688	369.7	-0.7183	42,722	97.77	0.772	6.478	1.028	28	yes	1e+16	66,768,642
Neptune	1.02e+26	30.07	60,182	367.5	0.6713	89,657	28.32	1.77	6.43	0.726	16	yes	1e+15	1.15e+08
Pluto	1.3e+22	39.48	90,560	366.7	-6.387	14,178	122.5	17.16	11.9	15.58	5	no	0	5,756,371

Readings from the table that matter for what follows: Mercury's spin:orbit ratio is exactly 1.500 (3:2). Mercury is the planet whose orbit lies closest to the solar equator (3.38°); Earth, by definition of the ecliptic, is the farthest of the eight (7.25°). Venus (Ω 76.68°) and Uranus (Ω 74.01°) have their nodes within 2° of the solar-equator node (75.76°). Saturn's ring mass is 2.7×10^{-8} of Saturn; Mars' moons are 1.9×10^{-8} of Mars — the same order, which the final chain exploits.

Equinox geometry (user: 'Equinox is the best time to judge angles'). The heliographic latitude of the disc centre, B0 = asin(sin i☉ · sin(λ☉ − Ω☉)), is -7.026° at the March equinox and 7.026° at the September equinox — 97 % of the full 7.25° tilt — and 0° near June 6 / December 7. So the equinoxes are indeed near-optimal for reading the Sun's tilt; the true extremes fall about two weeks earlier (≈ March 6, September 8).

2. Mercury / Vulcan — the four Vulcans the analogies produce

Pluto/Charon template (New Horizons): q = M_C/M_P = 0.1217, R_C/R_P = 0.51, a = 16.49 R_P = 19,591 km, period 6.3872 d = Pluto's spin = Charon's spin, barycentre 2,126 km from Pluto's centre, i.e. outside Pluto (R_P = 1,188 km). Mercury: M = 3.3011e+23 kg (GM known to 10^{-6} from MESSENGER), R = 2,439.7 km, spin 58.646 d, orbit 87.969 d, Hill radius 175,284 km at perihelion (220,658 km at mean distance).

Variant	q	M_V kg	R_V km	a km	a/R_M	Kepler T d	a/Hill	Bound?	Mercury reflex km	Barycentre outside Mercury?
V1 Charon-scaled	0.122	4.02e+22	1,244	40,223	16.5	3.732	0.229	yes	4,365	yes
V2 Spin-locked (58.646 d)	0.122	4.02e+22	1,244	252,366	103	58.65	1.44	NO	27,385	yes
V3 Earth-synchronous (365.26 d)	0.122	4.02e+22	1,244	854,287	350	365.3	4.87	NO	92,700	yes
V4 Chain-scaled (Phobos-class)	2.26e-08	7.47e+15	9.834	40,223	16.5	3.952	0.229	yes	0.0009107	no

- V1 Charon-scaled: q=M_Charon/M_Pluto, R_V=0.510 R_M, a=16.49 R_M (Pluto/Charon ratios applied to Mercury)
- V2 Spin-locked: Mutual tidal lock like Pluto/Charon: orbital period = Mercury's measured sidereal spin 58.646 d; a from Kepler III
- V3 Earth-synchronous: Strict anti-Earth lock: the Earth->Mercury direction turns once per sidereal year on average; a from Kepler III
- V4 Chain-scaled: From 'Saturn/rings is to Mercury/Vulcan is to Mars/moons': q = sqrt(q_SaturnRings * q_MarsMoons); rho = Phobos; a = 16.49 R_M




Figure 1 — Separation of each Vulcan variant against Mercury's Hill sphere. V2 and V3 lie outside it.

3. Test 1 — Kepler III and the Hill sphere

A Pluto/Charon-style mutual lock requires the companion's orbital period to equal the primary's spin. Mercury's spin is 58.646 d; the Kepler radius for that period about Mercury (+ q) is 252,366 km = 1.44 × the perihelion Hill radius: unbound. A Vulcan that keeps station anti-Earth must turn with the Earth→Mercury line, which turns once per sidereal year on average; the Kepler radius for 365.26 d is 854,287 km = 4.87 × Hill: unbound by a factor five. Only V1 and V4 (a = 16.5 R_M = 40,223 km) are bound, and their free period is 3.7–4.0 d, not 58.6 d and not 365 d.
What would overturn this: a measured Mercury spin of ≈ 3.7 d, or a measured Mercury mass ≈ 125 × larger (which the MESSENGER orbit excludes at the 10^{-6} level).

4. Test 2 — the anti-Earth lock, hour by hour from DE421

The Earth→Mercury unit vector u(t) was computed at 1-hour cadence for 2026–2027 from DE421. Its angular rate ω = |u̇| gives the instantaneous period a locked Vulcan must have: mean 286.8 d, range 164.1–14,915 d (the line nearly stalls at the elongation turn-arounds). The radius at which Mercury's gravity would supply that rate, a_ff = (GM/ω²)^(1/3), is never less than 482,370 km = 2.75 × Hill; the fraction of time it falls inside the Hill sphere is 0%.

Variant	Mercury gravity at a (m/s²)	centripetal needed (m/s²)	tangential needed, max (m/s²)	solar tide at a (m/s²)	gravity ÷ needed
V1 Charon-scaled	0.0136	3.1e-06	1.86e-05	5.78e-05	4.39e+03
V2 Spin-locked (58.646 d)	0.000346	1.95e-05	0.000117	0.000363	17.8
V3 Earth-synchronous (365.26 d)	3.02e-05	6.58e-05	0.000396	0.00123	0.458

Two independent failures. (i) At V1's radius gravity exceeds the needed centripetal acceleration 4,400-fold: a free body there orbits in 3.7 d; to hold it anti-Earth something must push outward at 1.4 × 10^{-2} m/s² continuously. (ii) The lock also needs a TANGENTIAL acceleration (up to 1.9 × 10^{-5} m/s² at V1, 4 × 10^{-4} at V3) because the Earth→Mercury line turns non-uniformly; no central force — gravity of Mercury, of the Sun, or any inverse-square field — can supply a tangential component. The axiom therefore requires a non-gravitational, continuously steered force. This is the sharpest result of the audit: an anti-Earth-locked Vulcan is dynamically impossible at ANY mass, including V4.
What would overturn this: a demonstration that the Earth→Mercury line turns uniformly (it does not — see Figure 2), or a bound orbit with a_ff inside the Hill sphere (none exists in 2026–27).



Figure 2 — Elongation, implied Vulcan period and free-fall radius ÷ Hill radius, 2026–2027. The free-fall radius never drops to the bound limit.

5. Test 3 — Mercury's own spin

Mercury's 58.646-d sidereal rotation (radar, 1965; MESSENGER libration) is exactly 2/3 of its 87.969-d orbit. A Charon-like companion at 16.5 R_M would, by Pluto/Charon's own template, have despun Mercury to the 3.73-d mutual-lock period. The 3:2 state is instead the Newtonian end-state of solar tides on an $e = 0.2056$ orbit. Status: the Charon analogy fails Mercury's measured spin. What would overturn this: a new radar/ranging spin period near 3.7 d.

6. Test 4 — Mercury's reflex wobble against ranging

Any companion makes Mercury orbit the pair's barycentre at $a \cdot q/(1+q)$. V1: 4,365 km (1.79 R_M — the barycentre is outside Mercury, exactly as for Pluto). V2: 27,385 km. V3: 92,700 km. V4: 0.911 m. Mercury's heliocentric orbit is tied to metre-class ranging (Mariner 10, MESSENGER, radar since 1964); a periodic displacement of thousands of km with a 3.7-d, 58.6-d or 365-d period is absent. V1–V3 exceed the 10-m precision by factors of 4.4e+05, 2.7e+06, 9.3e+06; V4 sits below it (0.091×). What would overturn this: a published Mercury residual of $\geq 10^3$ km with the stated period.




Figure 3 — Reflex displacement of Mercury for each variant against ranging precision and Mercury's radius.

7. Test 5 — can Vulcan stay hidden? Earth, the lunar far side, L2

Config A hides Vulcan behind Mercury's disc only if its angular radius $R_V/(D+a)$ plus the parallax shift $B\cdot a/(D(D+a))$ for an observer displaced by baseline B stays inside Mercury's angular radius R_M/D. Fraction of 2026–27 during which Vulcan is fully hidden:

Variant	Earth centre	Earth limb (6,378 km)	Lunar far-side camera (384,400 km)	Sun–Earth L2 (1.5 Gm)	Baseline needed (lunar distances)	Config B silhouette (fraction of disc area)
V1 Charon-scaled	1.00	1.00	1.00	1.00	12	0.26
V2 Spin-locked (58.646 d)	1.00	1.00	1.00	0.00	1.9	0.26
V3 Earth-synchronous (365.26 d)	1.00	1.00	0.00	0.00	0.58	0.26
V4 Chain-scaled (Phobos-class)	1.00	1.00	1.00	1.00	25	0.00

The user's instinct — a camera on the far side of the Moon would settle it — is right for V3 (always visible from the Moon) and partly right for V2 (visible about 40 % of the time); for V1 the 384,400-km baseline moves Vulcan only 0.18″ against a 1.8″ limb margin, and a baseline of 12 lunar distances (≈ 4.7 Gm) is needed. V4 (10 km) can hide from everything inside 25 lunar distances. Note what this buys: hiddenness is a geometric statement about where a thing could be, not evidence that it is there; Tests 1–4 are unaffected by it.




Figure 4 — Hidden fraction versus observer baseline; the Moon and L2 marked.

8. Config B — the near-side inversion and the dates on which it is testable

If Vulcan moves in front of Mercury (Earth → Moon → Vulcan → Mercury) its disc of angular radius R_V/(D−a) is silhouetted on Mercury's bright disc: for V1 a dark circle covering 26% of Mercury's area — a 10-cm telescope sees that. Lunar occultations of Mercury are the natural test dates. From DE421, Moon–Mercury conjunctions closer than 1.3° in 2026–2030 (green = geocentric occultation; the rest are occultations only from parts of Earth):

UTC	min sep °	occultation	Mercury elongation °	Earth–Mercury AU	Mercury radius ″
2026-02-18 23:10	0.119	yes	18.1	0.96	3.5
2027-01-08 05:18	0.475	partial Earth	4.46	1.41	2.39
2027-02-08 03:55	0.319	partial Earth	16.7	0.823	4.09
2027-03-06 06:32	1.11	partial Earth	24.2	0.759	4.43
2027-08-01 15:58	0.287	partial Earth	10.8	1.24	2.72
2027-12-28 16:35	0.696	partial Earth	9.82	1.36	2.47
2028-01-27 16:41	0.844	partial Earth	11.7	0.725	4.64
2028-02-23 00:05	0.202	yes	26.4	0.884	3.8
2028-09-20 05:22	1.13	partial Earth	21.1	0.738	4.56
2029-02-11 04:22	0.969	partial Earth	25.7	1.02	3.29
2029-09-09 08:15	0.746	partial Earth	13.2	0.652	5.16
2030-04-04 15:01	0.884	partial Earth	19.1	0.877	3.84
2030-08-29 01:53	0.751	partial Earth	4.48	0.624	5.39

The next geocentric event is 2028-02-23 00:05 UTC (elongation 26.4°, evening sky), which is the one animated in video/occultation_2028.mp4. No Mercury transit of the Sun occurs before 2032-11-13. What would overturn Config B: a resolved image of Mercury on any of these dates showing a clean disc. What would confirm it: a 25 %-area dark bite that is not a known albedo feature and that moves with the lunar geometry rather than with Mercury's rotation.




Figure 5 — Moon–Mercury conjunction calendar 2026–2030.

9. The record

- Le Verrier's Vulcan (1859): 1859: intra-Mercurial planet proposed to explain 38"/cy (later 43"); Lescarbault 'transit' 1859; never confirmed; searched at eclipses 1878 (Watson, Swift) — no body > ~a few hundred km exists inside Mercury's orbit (SOHO/LASCO era limits).
- Mariner 10, March 1974: UV excess 1974-03 attributed to a Mercury moon; identified as the star 31 Crateris.
- 243 Ida / Dactyl: 243 Ida's moon Dactyl: imaged by Galileo 1993-08-28, found in downlinked frames 1994-02-17 (Ann Harch); announced March 1994. First confirmed asteroid moon — not an April 1 release. The user's question — April Fool's or not — resolves to 'not': discovery in downlinked frames on 1994-02-17, announced in March 1994.
- MESSENGER: MESSENGER (2011-2015) performed dedicated satellite searches; none found. The published size limit of that search is the one number in this report I have not verified from the primary paper; it bounds V4 and is flagged in §14.

10. Verdict on Vulcan, variant by variant

Variant	Kepler/Hill	Lock kinematics	Mercury spin	Reflex vs ranging	Hiddenness (Earth)	Status
V1 Charon-scaled	bound (ok)	FAIL ×4,400 + tangential	FAIL (58.6 d ≠ 3.7 d)	FAIL $\times 4\times10^{5}$	hidden	refuted 3 ways
V2 Spin-locked	FAIL 1.44 × Hill	FAIL ×18 + tangential	(assumed)	FAIL $\times 3\times10^{6}$	hidden from Earth; 40 % visible from Moon	refuted 3 ways
V3 Earth-synchronous	FAIL 4.9 × Hill	gravity 0.5 × needed; tangential	FAIL	FAIL $\times 9\times10^{6}$	visible from Moon	refuted 4 ways
V4 Chain-scaled (10 km)	bound (ok)	FAIL as a LOCK; fine as a free moon	n/a (too small to lock)	passes (0.9 m)	hidden	not refuted as a free moon; refuted as a locked body

Summary: a Vulcan that is always anti-Earth cannot exist at any mass under Newtonian gravity — the lock needs a steered non-gravitational force. A small free moon of Mercury (V4-class, ~10 km) is not excluded by anything computed here; it is bounded only by direct imaging searches (Mariner 10, MESSENGER, BepiColombo from 2026), and a free moon does not satisfy Axiom A.

11. Mercury's perihelion without relativity

The measured anomalous advance (observed minus Newtonian planetary perturbations) is 42.98 ± 0.04″/century. The two Newtonian candidates are (a) an inner mass (Le Verrier's Vulcan, a ring) and (b) solar oblateness J2. For (b): the rate per century is $3\pi\, J2\, (R_\odot/p)^2 \cdot (36525/T)$. Fitting Mercury needs J2 = 0.000338, which is 1,538 × the helioseismic value 2.2×10^{-7} (a measurement of the Sun's shape, independent of planetary dynamics). The same J2 then predicts:

Body	measured anomaly ″/cy	± ″/cy	J2-fit prediction ″/cy	contribution of real J2 ″/cy
mercury	42.98	0.04	43	0.028
venus	8.62	0.4	4.42	0.0029
earth	3.84	0.4	1.42	0.00093
mars	1.35	0.1	0.332	0.00022
icarus	10.05	1	10.9	0.0071

The J2 route scales as $a^{-3.5}$ and fails Venus, Earth and Mars by factors of 2–4; it passes Icarus only because Icarus' eccentricity inflates both laws similarly. The data follow $a^{-2.5}$. For (a): an inner body or ring massive enough (≈ Mercury's mass inside 0.2 AU) would transit the Sun monthly and appear at mag ≈ 0–2 within 10° of the Sun; SOHO/LASCO has watched that region continuously since 1996 and sees nothing larger than a few hundred km. Neither Newtonian source survives measurement. This report, by instruction, does not go further than that sentence.




Figure 6 — Measured anomalous advances versus the solar-J2 fit; $a^{-2.5}$ and $a^{-3.5}$ guide lines.

12. The id–iv ladder, tested

Each 'A is to B is to C' was read three ways across seven metrics (mass, radius, heliocentric a, heliocentric T, rotation, density, local a): H1 geometric (A/B = B/C within the line), H2 mirror-equal (id A/B = iv A/B), H3 mirror-inverse (id A/B = iv B/A — the 'same backwards' reading). Residuals are in dex; |res| < 0.30 (a factor of two) counts as holding, < 0.10 as strong. Of 139 evaluable tests on the seven complete lines, 12 hold (8.6%); 4 strongly. A permutation null — 4,000 random ladders of the same shape drawn from the same entities — holds 10.7% of the time on average (5–95 %: 5.7%–16.4%); the probability of a random ladder matching or beating the user's rate is p = 0.73. The ladder does not beat chance on these metrics. Mass never holds on any line.

id	iv	metric	reading	residual dex	
Venus : Sol : Jupiter	Saturn : Jupiter : Neptune	rotation	H1_iv	+0.241	
Venus : Sol : Jupiter	Saturn : Jupiter : Neptune	density	H1_iv	-0.194	
Venus : Sol : Jupiter	Saturn : Jupiter : Neptune	density	H3_mirror_inv	+0.285	
Sol : Jupiter : Venus	Neptune : Jupiter : Saturn	radius	H1_id	-0.065	strong
Sol : Jupiter : Venus	Neptune : Jupiter : Saturn	rotation	H1_iv	+0.241	
Sol : Jupiter : Venus	Neptune : Jupiter : Saturn	density	H1_iv	-0.194	
Sol : Jupiter : Venus	Neptune : Jupiter : Saturn	density	H2_mirror_eq	-0.066	strong
Sol : Jupiter : Venus	Neptune : Jupiter : Saturn	density	H3_mirror_inv	+0.118	
Asteroid Belt : Mars : Mercury/Vulcan	Heliobarrier : Neptune : Pluto/Charon	density	H1_id	-0.120	
Earth : Luna : Mercury/Vulcan	Uranus : Uranus rings : Pluto/Charon	a (helio)	H2_mirror_eq	+0.000	strong
Earth : Luna : Mercury/Vulcan	Uranus : Uranus rings : Pluto/Charon	a (helio)	H3_mirror_inv	+0.000	strong
Earth : Luna : Mercury/Vulcan	Uranus : Uranus rings : Pluto/Charon	density	H2_mirror_eq	+0.113	

What does hold is informative: Sol : Jupiter : Venus is a near-exact geometric progression in RADIUS (695,700 : 69,911 : 6,052 — ratios 9.95 and 11.55); the density mirror Sol:Jupiter:Venus ↔ Neptune:Jupiter:Saturn holds at 0.07 dex; Earth:Luna ↔ Uranus:rings is exact in heliocentric a because satellites share their primary's orbit (a tautology, flagged as such). The singletons AB : HB : Sol and Sol : HB : AB are boundary statements — HB has no mass and Sol has no orbit; in radius the proportion misses by 6.7 dex.




Figure 7 — Residual heat-map, all lines × metrics × readings. Bold cells hold.

13. Planet X

The line 'iv Planet X is to Saturn is to Neptune' mirrors 'id Mars is to Jupiter is to Sol'. Solving for X in each reading and each metric:

reading	a AU	T yr	mass M⊕	radius km	V mag at opposition (p=0.5)	beyond Neptune?
H1_geometric	3.025	5.266	528	137,721	-6.9	no
H2_mirror_eq	2.793	4.671	0.0322	2,823	1.1	no
H3_mirror_inv	32.56	185.8	2.81e+05	1,201,079	-0.5	yes

H1 puts 1.7 Jupiter masses at 3.0 AU; H3 puts 0.85 solar masses at 32.6 AU; both would dominate the sky and the ephemerides and are excluded by any night's observation. H2 — the mirror reading the user's 'same backwards' phrasing points to — puts 0.032 M⊕ (0.3 Mars) at 2.79 AU, i.e. in the asteroid belt. The belt's dynamically measured total mass (from its perturbation of Mars, Pitjeva/EPM) is 2.4×10^{21} kg = 0.0004 M⊕: the ladder's Planet X exceeds what is there by a factor of 80, as one body (V ≈ 1.1 — nothing of the kind exists) or as dispersed mass (the dynamical total excludes it). The 'pattern break' the ladder requires is therefore real and already measured: the belt's mass deficit. (The minimum-mass solar nebula puts the shortfall nearer 10^3; the ladder's 80 is a milder version of the same statement.) The Kepler residuals in the table are 0.00 by construction — the inputs obey Kepler, so ratio-built outputs do too; that column tests nothing and is kept only to say so. Titius–Bode's next slot, 77.2 AU, is empty (the Kuiper cliff is at ≈ 48 AU); the independent Planet-Nine estimate (6.2 M⊕, a ≈ 380 AU, Brown & Batygin 2021) is unrelated to anything the ladder produces.




Figure 8 — The three Planet-X readings on the a–T plane with the planets, Titius–Bode slot and Planet-Nine estimate.

14. The final chain

Sol/Jupiter : Pluto/Charon : Earth/Luna : Saturn/rings : Mercury/Vulcan : Mars/moons, read as secondary/primary mass ratio q, separation in primary radii, and secondary period:

pair	q	a / R_primary	secondary period d
Sol/Jupiter	0.000954	1,119	4,333
Pluto/Charon	0.122	16.49	6.387
Earth/Luna	0.0123	60.34	27.32
Saturn rings	2.71e-08	1.778	0.3934
Mercury/Vulcan	?	?	?
Mars moons	1.89e-08	2.766	0.3189

In q the chain is not monotone (9.5e-4, 0.122, 0.0123, 2.7e-8, ?, 1.9e-8). Read as three mirror pairs — (Sol/Jupiter ↔ Pluto/Charon), (Earth/Luna ↔ Saturn/rings), (Mercury/Vulcan ↔ Mars/moons) — the last pair fixes Vulcan at q ≈ 2.26e-08: M_V ≈ 7.47e+15 kg, R ≈ 10 km at Phobos density, and a ≈ 2.22 R_M = 5,411 km (inside Mercury's fluid Roche limit of 8,533 km, outside the rigid limit of 4,378 km — Phobos' own situation about Mars). Its Kepler period there is 4.7 h; its angular separation from Mercury at 0.6 AU is 12″ at most, at V ≈ 12 beside a V ≈ 0 planet. This is V4. It survives every dynamical test in this report; it does not satisfy the anti-Earth axiom, and it is the object a BepiColombo satellite search would have to miss.




Figure 9 — Secondary/primary mass ratio along the final chain; Mercury/Vulcan interpolated from its neighbours.

15. Refutation ledger

Claim	Test	Measured	Status	What would flip it
Vulcan is Charon-like about Mercury (V1)	Mercury spin = lock period	58.646 d vs 3.73 d	refuted	spin ≈ 3.7 d
— same	reflex wobble	< 10 m vs 4,365 km	refuted	4,000-km periodic residual
Vulcan is always anti-Earth (any mass)	required vs available acceleration	needs outward 1.4e-2 + tangential 1.9e-5 m/s²	refuted (dynamics)	uniform turning of the Earth–Mercury line
Vulcan spin-locked to Mercury's 58.6 d (V2)	Hill sphere	a = 1.44 × Hill	refuted	Mercury mass ×3
Vulcan turns once a year with Earth (V3)	Hill sphere	a = 4.9 × Hill	refuted	—
Config B silhouette at occultations	imaging on listed dates	not yet observed (2028-02-23 next)	open, dated	clean disc on 2028-02-23
Far-side lunar camera reveals Vulcan	parallax	V3 yes, V2 40 %, V1/V4 no	computed	—
Mercury's 43″ from solar J2	J2 needed 3.4e-4 vs 2.2e-7; Venus/Earth/Mars	fails 3 of 5 bodies	refuted	helioseismic J2 revision ×1,500
Mercury's 43″ from an inner Vulcan/ring	SOHO/LASCO 1996–	nothing > few 100 km	refuted	—
Ladder proportions beat chance	permutation null	8.6 % vs 10.7 % random, p = 0.73	not supported	new metric set where holds exceed null
Planet X exists as the ladder defines it (H2)	belt mass / sky survey	0.0004 M⊕ present vs 0.032 needed	refuted as a body; real as a deficit	—
V4 Phobos-class moon of Mercury	imaging search limits	unverified limit (flag)	open	MESSENGER/BepiColombo limit < 10 km

16. Open items (not resolved here)

- MESSENGER's published satellite-search size limit (bounds V4). I did not read the primary paper in this session; the number is flagged, not asserted.
- Mariner 10's satellite limit (commonly quoted ≈ 5 km) — same status.
- Ring masses of Jupiter, Uranus and Neptune are order-of-magnitude estimates; Saturn's ($1.54 \pm 0.49 \times 10^{19}$ kg) is the only measured one.
- Saturn's moon count (274) moves with every MPC batch; the physics does not.
- Heliobarrier shape: Voyager/IBEX fix the nose (121.6 AU, λ 255.7°, β +5.1°); the tail length (comet vs croissant vs bubble) is unmeasured and drawn in the orrery as a model.
- Lunar occultation times are geocentric; topocentric circumstances for a given site need the observer's coordinates.

17. Addendum (2026-10-10, later): the relaxed axiom, Luna from Mercury, and the frame

Frame declaration. Everything in this addendum is computed in the Zecliptic frame as the author uses it — the solar equator (i = 7.25°, Ω = 75.76°): north is Sol's rotation pole, east is Sol's rotation sense. Earth-frame quantities appear in exactly one place (§17.3) and for one reason: the telescopes, the eyes and the photometric record are on Earth. No ecliptic or equatorial (RA/Dec) coordinates are used anywhere here.

17.1 The relaxed axiom

The axiom was loosened: Vulcan need not sit exactly on the Earth–Mercury line. It may be 'beside' Mercury (claimed invisible), between Luna and Mercury (claimed invisible), or 'just off' the far limb — 'the slightest occlusion'. The loosening is honest and it changes the test. A body that is only usually behind Mercury is a body on a free Kepler orbit, and a free orbit has three observable consequences: (i) a maximum angular separation from Mercury; (ii) a brightness of its own; (iii) mutual events — Vulcan passing in front of or behind Mercury dims the combined light, exactly the signal that fixed Charon's size in 1985–1990.

17.2 Luna from Mercury — the author's retrograde claim, tested

Claim: 'Luna would appear as in retrograde to Mercury as it passes.' DE421, hourly, 2026–2027, solar-equator frame. Two senses were measured. Against the stars, Luna's apparent motion from Mercury is dominated by Earth's: Earth is an outer planet from Mercury and retrogrades around every inferior conjunction (115.9 d), 18.4 % of the time; Luna rides along (18.6 %). Relative to Earth's disc — the sense in which Luna circles Earth — the result is sharp: on the half of its orbit that faces Mercury, Luna moves WEST 97.9 % of the time; on the far half it moves west 2.4 % of the time. Earth, at the centre of Luna's orbit, sees every part of it move east. Mercury, outside the orbit, sees the near half move west and the far half east. The orbit is unchanged and prograde; the observer moved from the centre to the outside.


The author's claim holds, and its mechanism is a mirror, not a reversal.

The mirror of the lunar occultation: when Earth sees Luna cover Mercury, Mercury sees Luna transit Earth's disc. Five such transits occur in 2026–2027 (all westward across Earth). The 2028-02-23 00:05 UTC event, from Mercury: Luna crosses Earth's 0.884-AU-distant disc moving WEST, minimum separation 2.2″ from Earth's centre, Earth's solar latitude B0 = -7.09°
(near its extreme — the tilt is fully visible on that date), Mercury's B = -3.12°.


Transit of Earth by Luna, seen from Mercury (UTC)	duration (h)	Luna's motion across Earth	rate (″/day)	Earth radius (″)
2026-02-18 22:00 → 01:00	4	WEST	-119	9.16
2027-01-08 04:00 → 07:00	4	WEST	-69.5	6.23
2027-02-08 03:00 → 05:00	3	WEST	-140	10.7
2027-08-01 15:00 → 17:00	3	WEST	-90.3	7.09
2027-12-28 16:00 → 18:00	3	WEST	-73.5	6.46



Luna from Mercury, 2026–27, solar-equator frame. Top: Earth and Luna against the stars (Earth's retrograde loops at each inferior conjunction). Middle: Luna's motion relative to Earth's disc — red shading marks the Mercury-facing half, where it runs west. Bottom: Luna–Earth separation vs Earth's angular radius; the dips below the blue line are transits.



One lunation seen from Mercury (Feb 2028), relative to Earth. Red: Mercury-facing half, moving west; blue: far half, moving east.
The ellipse is Luna's orbit seen nearly edge-on from inside Earth's orbit.



2028-02-23 from Mercury: Luna crosses Earth's disc westward. Earth sees the same event as Luna moving east over Mercury.

17.3 'Beside is invisible' — tested (Earth frame, by reason: the instruments are on Earth)

For each Vulcan variant on a free Kepler orbit about Mercury, the maximum angular separation from Mercury as seen from Earth over 2026–2027, the brightness relative to Mercury (same albedo: area ratio), the depth of a mutual event, and the fraction of an edge-on orbit spent within 1,000 km projected of Mercury's centre ('merged').

variant	free period (d)	max sep. from Mercury (″)	fainter than Mercury (mag)	mutual-event depth (mag)	merged fraction of orbit	resolvable by
V1 Charon-scaled	3.73	38–100	1.46	0.251	0.0158	small telescope (>2 arcsec)
V2 Spin-locked (58.646 d)	58.6	240–627	1.46	0.251	0.00252	naked eye (>60 arcsec)
V3 Earth-synchronous (365.26 d)	365	812–2123	1.46	0.251	0.000745	naked eye (>60 arcsec)
V4 Chain-scaled (Phobos-class)	3.95	38–100	12	1.76e-05	0.0158	small telescope (>2 arcsec)

Reading. 'Beside' is not invisible. A Charon-scaled Vulcan (V1) on a free orbit reaches 38–100″ from Mercury: 1.5 magnitudes fainter than Mercury and resolvable in any telescope since 1610 (Galileo's phases of Mercury; Hevelius; Schröter; Schiaparelli's 1881–89 maps with a 22 cm refractor; Antoniadi at Meudon). V2 and V3 reach naked-eye separations (4–35′). V1 also produces mutual events of 0.25 mag every 1.87 d in edge-on seasons; Mercury's whole-disc photometry (Mallama 2002, 2017; SOHO/LASCO, STEREO) is good to 0.01–0.02 mag.
'The slightest occlusion' is therefore the strongest refutation of V1–V3, not the weakest. For a body to be beside Mercury and truly merged with it from Earth, it must stay within ~1,000 km projected of Mercury's centre — inside Mercury (R = 2,440 km). The only Vulcan the
relaxed axiom still allows is V4: a ~10 km body, 12 mag fainter than Mercury, mutual-event depth 17 micro-mag — undetectable from Earth, bounded only by spacecraft imaging (MESSENGER / BepiColombo), as before.

17.4 Solar equinoxes — the angle-judging epochs in this frame

Earth's March/September equinoxes are when Earth's solar latitude |B0| is largest (7.03°) — the tilt between the ecliptic and the Zecliptic is then fully visible, which is why the author's 'equinox is the best time to judge angles' holds. The complementary epochs are the SOLAR equinoxes: when a body crosses Sol's equatorial plane (B = 0). Both are tabulated from DE421.

body	crosses Sol's equator (UTC)	direction	solar-equatorial longitude (°)
Earth	2026-06-06 18:58	ascending	180
Earth	2026-12-08 06:30	descending	360
Earth	2027-06-06 23:01	ascending	180
Earth	2027-12-08 13:45	descending	360
Earth	2028-06-06 04:39	ascending	180
Earth	2028-12-07 19:44	descending	360
Mercury	2026-01-29 08:53	ascending	251
Mercury	2026-03-03 12:02	descending	73.2
Mercury	2026-04-27 07:13	ascending	251
Mercury	2026-05-30 12:05	descending	73.3
Mercury	2026-07-24 05:39	ascending	251
Mercury	2026-08-26 12:03	descending	73.4
Mercury	2026-10-20 04:09	ascending	251
Mercury	2026-11-22 11:55	descending	73.3
Mercury	2027-01-16 02:44	ascending	251

Earth equinox	Earth B0 (°)	Mercury B (°)	Luna B (°)
2026-03-20	-7.02	-3.07	-7.02
2026-09-22	7.07	-3.31	7.06
2027-03-20	-7.04	-3.22	-7.04
2027-09-23	7.04	-2.32	7.04
2028-03-20	-7.02	-1.91	-7.01
2028-09-22	7.04	-0.0754	7.04

Mercury's nodes on Sol's equator sit at solar-equatorial longitudes 251.3° (ascending) and 73.3° (descending); Earth's at 180.0° and 0.0° (by construction of the frame). Mercury crosses the solar equator every 44 days; Earth every 183 days (≈ 6 June ascending, ≈ 8 December descending).

17.5 '42 Pis in a year'

Luna's orbits per year, counted against the stars: 365.25636 / 27.321661 = 13.3687. Times π: 41.999. So 42/π = 13.3690 reproduces Luna's star-fixed month count to 0.002 %. Counting against Sol instead gives 38.9, so the mnemonic belongs to the star-fixed month — Luna against the stars, which is also the count Mercury would make. It is a four-significant-figure coincidence with no mechanism
offered; as a mnemonic it is exact enough to use.

17.5b Charon is 180 Pis (added 2026-10-11)

Pis = orbits per year, counted against the stars, × π; the year's frame declared per row. Charon (period 6.387221 d) makes 57.185 orbits per Earth year; × π = 179.65, i.e. **180 Pis** to -0.19 %. The number it lands beside is 180/π = 57.296, one radian in degrees: Charon orbits Pluto once per Earth year for every degree in a radian. Pluto, locked to Charon, rotates 180 Pis a year as well. Pluto's year is so long that Charon's
star-fixed and Sol-fixed counts differ by 0.007 %, so the frame ambiguity that mattered for Luna (42 vs 38.9) does not arise.
Per Pluto year Charon is 44,542 Pis — nothing round. Charon's orbit plane (Pluto's equator) is tilted 119.1° to the Zecliptic.
Other round hits in the full moon table at the 1 % level: Titan 72, Hydra 30, Kerberos 36 — expected at about that rate by chance for a table of this size.
Five and seven Pis, in every frame where a Mercury moon is physical, fall outside the survival annulus (§19), which lives at 24–370 Pis per Mercury year;
read as a clock from 2026-10-11, 5 Pis of Earth years is 1.59 yr (May 2028) and 7 Pis is 2.23 yr (end of 2028). The Saturn→Uranus→Neptune angle sums,
the Uranus (-1.75°) and Venus node coincidences with the solar equator's node, and the Mercury ↔ Pluto/Charon mirror (3:2 spin-orbit vs 3:2 Pluto:Neptune) are in
PIS_42_180_20261011.md with data/pis.json (tools/vulcan/pis.py).

17.6 Added deliverables

- Video video/luna_from_mercury.mp4 (1,440 frames, 48 s): Luna transiting Earth's disc as seen from Mercury, 2028-02-23 ± 6 h, solar-equator frame, with a Zecliptic top-down inset (Mercury-facing half of Luna's orbit in red) and the frame declaration on screen. Contact sheet video/luna_from_mercury_sheet.png.
- Images img/luna_01_from_mercury_rates.png, img/luna_02_lunation_from_mercury.png, img/luna_03_transit_of_earth_2028.png.
- Data data/luna_from_mercury.json (fractions, transits, 'beside' table, solar equinoxes, 42/π), data/ephem_luna_from_mercury.json (1-min DE421 export in solar-equator coordinates driving the Godot shot).
- Code tools/vulcan/luna_from_mercury.py; shot LunaFromMercury in game/tools/record_vulcan.gd.

18. Deliverables

- Videos (video/*.mp4, 1280×600, 30 fps): lock_orrery (Zecliptic view, Config A over 2026), hill_frame (locked vs free Vulcan, two scales, 180 d), occultation_2028 (2028-02-23 geocentric event, Config A vs B insets), binaries (Pluto/Charon vs Mercury/Vulcan to scale about their barycentres), ladder_orrery (log-radius system with AB, HB, Planet X; each id/iv line highlighted with its residual), equinox_zecliptic (orbit planes against the solar equator at the September 2026 equinox, then a date sweep to March 2027), luna_from_mercury (2028-02-23 ±6 h from Mercury: Luna transiting Earth westward, with Zecliptic inset). Contact sheets video/*_sheet.png.
- Images (img/): nine analysis figures listed above plus luna_01..03 from the addendum.

• Data (data/): bodies_metrics.{json,csv,md}; vulcan_derivation.json; ladder_analysis.json; ladder_table.md; ladder_null.json; lunar_occultations_mercury_2026_2030.csv; ephem_2026_2027_6h.json and ephem_occultation_closeup.json (DE421 exports driving the Godot shots).

• Code (tools/vulcan/, game/tools/record_vulcan.gd): rerunnable end to end: python3 bodies.py && python3 vulcan_derivation.py && python3 ladder_analysis.py && python3 ladder_null.py && DISPLAY=:1 godot --path game -s res://tools/record_vulcan.gd && bash tools/vulcan/encode_vulcan.sh && python3 build_report.py.

• Addendum §19 (elimination pass): data/v4_annulus.json, data/perihelion_newtonian.json, data/ladder_oos.json; img/v4_01_tidal_floor.png, img/v4_02_hill_stability.png, img/perihelion_01_hall.png, img/perihelion_02_budget.png, img/ladder_oos_01_recurrence.png, img/ladder_oos_02_null.png; tools/vulcan/v4_annulus.py, perihelion_newtonian.py, ladder_oos.py; BEPICOLOMBO_OBSERVATION_REQUEST_20261010.md; GAMEPLAN_ELIMINATION_20261010.md; REFLECTION_ELIMINATION_META_20261010.md.

19. Addendum (2026-10-10, later still): the elimination pass

GAMEPLAN_ELIMINATION_20261010.md set the order: (a) the V4 annulus; (b) the three remaining Newtonian causes of the 43″; (c) the MESSENGER / Mariner 10 / BepiColombo record; (d) the ladder out of sample; (e) an observation request; (f) this addendum. Frame: solar equator (Zecliptic) throughout, except where a line says Earth and why. No relativity is used anywhere in this section.

19.1 Row 6 — where a free satellite of Mercury can live (V4 annulus)

Tides: $da/dt = -3(k_2/Q)(m/M)(R/a)^5 \cdot n \cdot a$ with $k_2 = 0.45$ (MESSENGER), integrated over 4.5 Gyr, gives the radius below which a body of a given size cannot have
survived from formation. Q = 100: 10 km body → 13,254 km; 1 km → 4,580 km; 100 m → 1,582 km. The chain-interpolated V4 (9.8 km at 5,411 km, §14) spirals into Mercury in 7–42 Myr (Q = 50–300): it cannot be primordial where the chain puts it. **Row 5 of the board is eliminated as a primordial body.**

Sun: 112 test particles on circular orbits from 8,000 to 260,000 km, prograde and retrograde, two phases each, leapfrog at 200 s for 100 Mercury years
with the Sun on Mercury's e = 0.2056 orbit and Mercury's J_2. Fast-ejection edge: prograde 81,473 km (canonical 0.5 R_Hill at perihelion = 87,641 km),
retrograde 119,950–136,457 km. Radiation pressure: a_rp/g < 1e-9 for R ≥ 100 m anywhere in the annulus.

edge	km	period (d)	from Earth at 0.9 AU (″)
tidal floor, 10 km body	13,254	0.75	20
tidal floor, 1 km body	4,580	0.15	7
prograde ceiling (this run / canonical)	81,473 / 87,641	11.4 / 12.7	125 / 134
retrograde ceiling	136,457	24.7	209
Hill radius at perihelion	175,282	—	269




Figure 19a — Tidal floor a_min(R, Q) with the canonical prograde ceiling; the shaded band is where a 4.5 Gyr-old satellite can be. The white point is the chain's V4: below every floor.




Figure 19b — Fate of the 112 test particles after 100 Mercury years. First integration was wrong (indirect term sign); the bug was found because every particle died in 463 days, including those at 8,000 km, which is impossible. Ledgered.

The record: Mariner 10's 1974 'moon' (UV source detached from Mercury, 4 km/s) was the star 31 Crateris (Stratford 1980). MESSENGER ran dedicated
satellite-search campaigns in 2011 and 2013 (MDIS WAC; JHUAPL 2012-03-21, 2013-07-18) and found none; its 2011 orbit (2,640–17,633 km) crossed the inner
annulus. The published size limit of those campaigns and the 2007 ground-based search (Planet. Space Sci. 55, 2037) were **not retrievable today** and are
flagged as unread. BepiColombo enters Mercury orbit 2026-11-21 (MPO 480 × 1,500 km; Mio 590 × 11,640 km from 2026-12-09/10); from inside the annulus a 20 m body is
V ≈ 4–9. BEPICOLOMBO_OBSERVATION_REQUEST_20261010.md states the frame, cadence, depth, kinematic filter and closure conditions. **Row 6 stays open, confined,
with a dated instrument named.**

19.2 Row 10 — the three remaining Newtonian causes of the 43″

Hall's law F ∝ r^−(2+δ) advances every apse by πδ per orbit. δ = 1.597e-07 reproduces Mercury's 42.98″/cy and then predicts Venus 16.8″ (measured 8.62), Earth 10.4″ (3.84), Mars 5.5″ (1.35) and a lunar perigee excess of
1.38″/yr against a lunar-laser-ranging limit of ~1e-4″/yr (1e+04 σ). Brown rejected it on the Moon in 1903; LLR buries it.

Exterior ring. $\dot{\omega} = (3/4)\cdot n\cdot(m_r/M_\odot)\cdot(a/b)^3$. At 1 AU the ring would need 0.61 M⊕ — 4e+07 × the zodiacal cloud (1e17 kg); at 0.6 AU 0.13 M⊕. Mass *inside* Mercury's orbit regresses the apse (apsidal angle < π) and cannot help. The asteroid belt's real contribution is 0.0014″/cy.

Perturbing-mass error. $\dot{\omega} = (n/4)(m'/M_\odot)\alpha^2\, b^{(1)}_{3/2}(\alpha)$ per planet, Laplace coefficient by quadrature. Budget reproduced: Venus 286.2″ (Clemence 277.8), Earth 96.6 (90.0),
Jupiter 160.1 (153.6), Saturn 7.6 (7.3), Mars 2.4 (2.5): total 553″ vs 531 (first-order secular theory; 4 % high, as expected without second-order terms).
To supply 43″ Venus would need a 15 % mass error (GM known to 1e-9: ×2e+08); Jupiter 27 % (×3e+07).

candidate	tuned to Mercury	then predicts	measured	status
solar J_2 (§11)	J_2 = 3.4e-4 fit	Venus 4.4″, Earth 1.4″	8.6, 3.8 (and J_2 = 2.2e-7 measured, ×1,538)	eliminated
intra-Mercurial planet/ring (§2–10)	mass inside 0.39 AU	regression, not advance; visible at transit/eclipse	none seen since 1859	eliminated
Hall exponent	δ = 1.60e-07	Moon perigee +1.38″/yr	< 1e-4″/yr (LLR)	eliminated
exterior zodiacal ring	0.61 M⊕ at 1 AU	—	1e17 kg (2e-8 M⊕)	eliminated
planetary-mass error	Venus +15 % / Jupiter +27 %	—	1e-9 / 1e-8 relative	eliminated




Figure 19c — Hall's law tuned to Mercury against the measured anomalous advances of Venus, Earth, Mars.




Figure 19d — The Newtonian planetary budget reproduced from Laplace coefficients, against Clemence (1947), with the 43″ line.

No Newtonian cause of Mercury's 43″/century survives measurement: solar J2 (report §), inner planet/ring (report §), Hall's exponent, exterior zodiacal mass, and perturbing-mass error are each excluded by more than two orders of magnitude. The residual stands as an unexplained measurement under the rules of this audit. This statement is a receipt; it is also an arrow, recorded and not followed (see GAMEPLAN §3, REFLECTION §2).

19.3 Row 11 — the ladder out of sample

Lines free of Sol-only constructs (no belts, Luna, rings, Vulcan, Planet X): 1, 3, 5. Roles mapped to five other hierarchical systems by two rules
fixed before the numbers were looked at — ordinal by distance from the primary, and rank by mass (size where mass is unknown). Same four hypotheses, same
0.30-dex hold. Null: 4,000 random role assignments per system per rule. Sol itself on these lines: 8/54.

system	mapping	holds	rate	random-mapping mean	p (random ≥ rule)
Saturn moons	ordinal	10/54	0.19	0.17	0.405
Saturn moons	massrank	11/53	0.21	0.17	0.259
Jupiter moons	ordinal	12/54	0.22	0.18	0.257
Jupiter moons	massrank	19/54	0.35	0.18	0.025
TRAPPIST-1	ordinal	0/15	0.00	0.19	1.000
TRAPPIST-1	massrank	6/37	0.16	0.19	0.697
Kepler-90	ordinal	0/15	0.00	0.08	1.000
Kepler-90	massrank	2/15	0.13	0.09	0.215
HD 10180	ordinal	0/5	0.00	0.10	1.000
HD 10180	massrank	0/14	0.00	0.10	1.000

1 of 10 mapping tests beat random at p < 0.05 (Jupiter's moons by mass rank, p = 0.025) — the number expected at 5 % from ten tries.
Of the 8 proportions that hold in the Sol system on these lines, 19/42 recur out of sample; excluding density, 6/16.
Density is a degenerate metric: 81 % of all pairwise density ratios among Saturn's moons and 100 % among TRAPPIST-1's planets are already within 0.30 dex of unity, so a density 'hold' says nothing about a ladder. **Row 11: the ladder does not transfer. Eliminated as a law; retained as a mnemonic.**




Figure 19e — Each Sol-system hold re-tested in five systems under two mappings; bold cells recur.





Figure 19f — Rule-based mapping hold-rate against random mappings, per system.

19.4 Board after the pass


row	claim	before	after	closed by
5	chain-interpolated V4 (9.8 km, 5,411 km)	open	**eliminated** (as primordial)	tidal decay 7–42 Myr, k_2 measured
6	any free satellite of Mercury (V4 class)	open	open, **confined** to 13,000–136,000 km; dated instrument named	BepiColombo 2026-11-21 →; MESSENGER null (limit unread)
7	Config B 2028-02-23 (Vulcan on near side)	open	open (dated)	lunar occultation 2028-02-23 00:05 UTC
10	Newtonian 43″	open	**no Newtonian cause survives**	Hall/LLR; ring/zodiacal mass; masses/GM
11	ladder as law	open (in-sample p = 0.73)	**eliminated as law**	out-of-sample 1/10 at p < 0.05; recurrence 6/16 ex-density

19.5 Ledger of the chronicler's own errors in this pass

- v4_annulus.py, first run: sign of the indirect term in the solar tide ($-rs/|rs|^3$ for $+rs/|rs|^3$). Symptom: 112/112 particles 'escaped' within 463 d including those at 8,000 km. Caught by the impossibility, fixed, assertion added so it cannot recur silently.
- Perihelion budget 4 % high relative to Clemence: first-order secular theory only. Not corrected; stated.
- Two literature items (MESSENGER campaign size limit; 2007 P&SS ground search) not retrievable with today's tools. Stated as unread rather than paraphrased.




video/lock_orrery.mp4 — six evenly spaced frames




video/hill_frame.mp4 — six evenly spaced frames




video/occultation_2028.mp4 — six evenly spaced frames



video/binaries.mp4 — six evenly spaced frames



video/ladder_orrery.mp4 — six evenly spaced frames



video/equinox_zecliptic.mp4 — six evenly spaced frames