

L -functions for all meromorphic $\mathrm{SL}_2(\mathbb{Z})$ modular forms, via manifestly finite contours

David A. McGady

Abstract

We give a single finite, regulator-free functional realizing in one object the period polynomials (at integer s) and the completed L -functions (at general s) of meromorphic modular forms on $\mathrm{SL}_2(\mathbb{Z})$, with poles anywhere: at the cusp and in the interior. It is a two-segment contour integral with an explicit Hurwitz-zeta T -kernel, built from any finite base point $\tau_0 \in \mathbb{H}$ and manifestly finite. For weakly holomorphic forms it is independent of τ_0 and of the path, reproducing at $\tau_0 = i$ the special L -values of Bringmann–Fricke–Kent [1]; its integer- s values give the period polynomial, Brown’s quasi-periods [2] in closed form. For forms with interior poles the value depends only on the homotopy class of the contour — fixed up to the residues it crosses — and a natural choice of class, the one prescription a pole on the line $\mathrm{im} \tau = 1$ requires, renders it canonical. The resulting closed form generalizes [3]: a regularized cusp sum plus one polylog integral per interior pole. The construction introduces no new objects, unifying these periods and L -values through a single all- s kernel, and every formula is checked numerically [4].

1 A finite L -integral

Definition 1.1 (S - and T -segments). *Let $\tau_0 \in \mathbb{H}$ and $f \in F_k$, let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ denote the generators of $\Gamma = \mathrm{SL}_2(\mathbb{Z})$, and let $f^{-1}(\infty)$ be the locus of all poles of $f \in \mathbb{H}$ (inclusive of the cusp of $\mathrm{SL}_2(\mathbb{Z})$). The S - and T -segments at τ_0 are smooth paths in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$,*

$$\gamma_{\tau_0}^S : -1/\tau_0 \longrightarrow \tau_0, \quad \gamma_{\tau_0}^T : \tau_0 - 1 \longrightarrow \tau_0, \quad (1)$$

running from $S^{-1}\tau_0 = -1/\tau_0$ and $T^{-1}\tau_0 = \tau_0 - 1$ to τ_0 .

Avoiding poles is automatic for $f \in M_k^!$, as long as the base-point τ_0 , is not at a cusp of $\mathrm{SL}_2(\mathbb{Z})$, i.e. there is only the trivial homotopy class. When $f \in F_k$, and f has poles at finite and non-zero imaginary heights, integration contours fall into nontrivial homotopy classes.

Definition 1.2 (Hurwitz T -kernel). *Let $\zeta(s, x) = \sum_{m \geq 0} (m + x)^{-s}$ as a function $\zeta : \mathbb{C}_s \times (\mathbb{C} \setminus \{0\})_x \rightarrow \mathbb{C}$ denote the Hurwitz zeta function (meromorphic in s , holomorphic in $x \in \mathbb{H}$). We define Hurwitz T -kernel as*

$$\tilde{k}_T(\tau, s) := \zeta(1 - s, \tau + 1) - e^{i\pi(s-1)} \zeta(1 - (k - s), \tau + 1). \quad (2)$$

Definition 1.3 (L -integral). *Let $f \in F_k$, $\tau_0 \in \mathbb{H}$, and $s \in \mathbb{C}$. Further, let $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$ be the S - and T -segments at τ_0 , as in Definition 1.1, and let $\tilde{k}_T$ be as defined in Definition 1.2. The L -integral of f is*

$$L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) := e^{-\pi i s/2} \left[\int_{\gamma_{\tau_0}^S} f(\tau) \tau^{s-1} d\tau + \int_{\gamma_{\tau_0}^T} f(\tau) \tilde{k}_T(\tau, s) d\tau \right]. \quad (3)$$

We use this L -integral to define a generalized L -function:

Definition 1.4 (*L*-function). Let $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ be as defined in 1.3. We define the generalized *L*-function of $f \in F_k$ to be,

$$L(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T). \quad (4)$$

Having defined the *L*-integral and *L*-function, we now prove both are manifestly finite throughout:

Lemma 1.5 (Finiteness). For $f \in F_k$, both integrals in Definition 1.3 are finite for every $s \in \mathbb{C}$, every $\tau_0 \in \mathbb{H}$, and every choice of paths $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$.

Proof. Both segments are compact and lie in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, so f is continuous on them; τ^{s-1} is continuous on $\mathbb{H}$, and $\tilde{k}_T(\tau, s)$ is continuous there since $\zeta(\nu, \tau + 1)$ is holomorphic in $\tau \in \mathbb{H}$ (Definition 1.2). A continuous integrand on a compact path is integrable, for every $s \in \mathbb{C}$. $\square$

We now turn to path-dependence and τ_0 -independence of the *L*-integral and the *L*-function of $f \in F_k$. This holds for all $f \in M_k^!$:

Proposition 1.6 (Basepoint and path independence). For $f \in M_k^!$ the value $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ is independent of $\tau_0 \in \mathbb{H}$ and of the paths: with no interior poles, $\mathbb{H}$ is simply connected, so all admissible contours are homotopic. We write $L^*(f, s)$ for the common value and $L(f, s) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s)$.

Proof. With no interior poles the integrands $f(\tau)\tau^{s-1}$ and $f(\tau)\tilde{k}_T(\tau, s)$ are holomorphic on $\mathbb{H}$, which is simply connected; by Cauchy's theorem each integral is unchanged under any deformation fixing its endpoints, hence depends on those endpoints alone. This is path-independence.

For independence of τ_0 we differentiate the two integrals and watch them cancel. The *T*-segment runs from $T^{-1}\tau_0 = \tau_0 - 1$ to τ_0 — two endpoints a unit apart, moving together with τ_0 — so by Leibniz's rule and the *T*-periodicity $f(\tau_0 - 1) = f(\tau_0)$,

$$\partial_{\tau_0} \int_{\gamma^T} f \tilde{k}_T = f(\tau_0) \tilde{k}_T(\tau_0, s) - f(\tau_0 - 1) \tilde{k}_T(\tau_0 - 1, s) = f(\tau_0) \delta_T \tilde{k}_T(\tau_0 - 1, s), \quad (5)$$

where $\delta_T \tilde{k}_T(\tau, s) := \tilde{k}_T(\tau + 1, s) - \tilde{k}_T(\tau, s)$. This is the whole role of δ_T : the unit gap between the endpoints turns the ∂_{τ_0} boundary term into a unit shift of the kernel. That shift is elementary — the kernel was built for it — since $\zeta(\nu, x + 1) - \zeta(\nu, x) = -x^{-\nu}$ gives, on both terms at $x = \tau + 1$,

$$\delta_T \tilde{k}_T(\tau, s) = -(\tau + 1)^{s-1} + e^{i\pi(s-1)}(\tau + 1)^{k-1-s} \quad (6)$$

$$\implies \partial_{\tau_0} \int_{\gamma^T} f \tilde{k}_T = f(\tau_0) [-\tau_0^{s-1} + e^{i\pi(s-1)}\tau_0^{k-1-s}]. \quad (7)$$

The *S*-segment runs $-1/\tau_0 \rightarrow \tau_0$; its lower endpoint moves with $\partial_{\tau_0}(-1/\tau_0) = \tau_0^{-2}$, so by Leibniz, the modularity $f(-1/\tau_0) = \tau_0^k f(\tau_0)$, and the phase $(-1/\tau_0)^{s-1} = e^{i\pi(s-1)}\tau_0^{1-s}$,

$$\partial_{\tau_0} \int_{\gamma^S} f \tau^{s-1} = f(\tau_0)\tau_0^{s-1} - f(-1/\tau_0)(-1/\tau_0)^{s-1}\tau_0^{-2} = f(\tau_0)[\tau_0^{s-1} - e^{i\pi(s-1)}\tau_0^{k-1-s}]. \quad (8)$$

The two brackets are negatives of each other, so $\partial_{\tau_0} L^* = e^{-\pi i s/2}(\partial_{\tau_0} \int_{\gamma^S} + \partial_{\tau_0} \int_{\gamma^T}) = 0$. $\square$

Remark 1.7 (Classical *L*-function contour). Let $f \in S_k$. As $\tau_0 \rightarrow i\infty$, the *S*-segment maps onto the standard *L*-function contour, $[0, i\infty]$. Simultaneously, as $f(it) \sim e^{-2\pi t}$ as $t \rightarrow \infty$, the *T*-segment vanishes exponentially. Thus, in this limit, $L^*(f, s)$ exactly maps onto the classical *L*-function integral [14]:

$$\lim_{\tau_0 \rightarrow i\infty} L^*(f, s) = \int_0^\infty t^{s-1} f(it) dt. \quad (9)$$

2 L -functions for weak modular forms

Proposition 2.1 (Hurwitz-Gamma). *For $n \neq 0$ and $\nu \in \mathbb{C}$,*

$$\int_i^{i+1} e^{2\pi i n u} \zeta(\nu, u) du = e^{i\pi(1-\nu)/2} (2\pi n)^{\nu-1} \Gamma(1-\nu, 2\pi n). \quad (10)$$

Proof. Write $\zeta(\nu, u) = \sum_{m \geq 0} (m+u)^{-\nu}$ and shift $u \mapsto u+m$ in the m -th term; as $e^{2\pi i n(u+m)} = e^{2\pi i n u}$ for integer m , the unit segments stack into one ray,

$$\int_i^{i+1} e^{2\pi i n u} \zeta(\nu, u) du = \sum_{m \geq 0} \int_{i+m}^{i+1+m} e^{2\pi i n v} v^{-\nu} dv = \int_i^{i+\infty} e^{2\pi i n v} v^{-\nu} dv. \quad (11)$$

For $n > 0$ the integrand is holomorphic in the first quadrant and decays as $\text{im } v \rightarrow \infty$ (since $|e^{2\pi i n v}| = e^{-2\pi n \text{im } v}$), so the horizontal ray rotates to the vertical $[i, i\infty)$. With $v = it$ ($t \geq 1$; then $dv = i dt$, $e^{2\pi i n v} = e^{-2\pi n t}$, $v^{-\nu} = e^{-\pi i \nu/2} t^{-\nu}$) and $w = 2\pi n t$,

$$\int_i^{i+\infty} e^{2\pi i n v} v^{-\nu} dv = e^{i\pi(1-\nu)/2} \int_1^\infty e^{-2\pi n t} t^{-\nu} dt = e^{i\pi(1-\nu)/2} (2\pi n)^{\nu-1} \Gamma(1-\nu, 2\pi n). \quad (12)$$

For $n < 0$ the integrand instead grows as $\text{im } v \rightarrow \infty$, so the closing arc survives and the rotation is illegal; but both sides are analytic in n and agree for $n > 0$, so the identity continues to $n < 0$, with $\Gamma(1-\nu, 2\pi n)$ on its principal branch. At $n = 0$ the prefactor $(2\pi n)^{\nu-1}$ is singular — that mode, the constant Fourier coefficient, is handled separately. $\square$

Theorem 2.2 (L -function for weakly holomorphic forms). *For $f \in M_k^!$, we may set $\tau_0 = i$, and prove*

$$L^*(f, s) = -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right). \quad (13)$$

This L -integral satisfies the functional equation $L^(f, s) = i^k L^*(f, k-s)$.*

Proof. By path independence and base-point independence from Proposition 1.6, we may freely set $\tau_0 = i$. Here, the S -segment degenerates ($-1/i = i$), and the L -integral simplifies dramatically:

$$L^*(f, s) = e^{-\pi i s/2} \int_{i-1}^i f \tilde{k}_T d\tau. \quad (14)$$

We then expand f in terms of its q -series, $\sum_n c_f(n) e^{2\pi i n \tau}$, and substitute $u = \tau + 1$, and note that $e^{2\pi i n(u-1)} = e^{2\pi i n u}$. Here, each $n \neq 0$ mode is handled by Proposition 2.1. The first kernel term with $\nu = 1-s \leftrightarrow 1-\nu = s$ gives

$$e^{-\pi i s/2} e^{\pi i s/2} (2\pi n)^{-s} \Gamma(s, 2\pi n) = \Gamma(s, 2\pi n) / (2\pi n)^s, \quad (15)$$

and the second kernel term with $\nu = s-k+1 \leftrightarrow 1-\nu = k-s$ gives

$$-e^{-\pi i s/2} e^{i\pi(s-1)} e^{\pi i(k-s)/2} (2\pi n)^{s-k} \Gamma(k-s, 2\pi n) = i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}}. \quad (16)$$

So each $n \neq 0$ mode contributes $\Gamma(s, 2\pi n)/(2\pi n)^s + i^k \Gamma(k-s, 2\pi n)/(2\pi n)^{k-s}$. Crucially, the $n = 0$ mode contributes

$$c_f(0) e^{-\pi i s/2} \int_{i-1}^i \tilde{k}_T d\tau = -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right). \quad (17)$$

Combining these reproduces Eq. (13). Finally, the functional equation, $L^*(f, s) = i^k L^*(f, k-s)$, follows by direct inspection. $\square$

2.1 Period polynomials, periods and quasi-periods

Because $L^*(f, s)$ is canonical for every $f \in M_k^!$, the period polynomial needs no separate construction — it is the generating polynomial of the critical values.

Definition 2.3 (Period polynomial). *For $f \in S_k^!$ and $n = k - 2$, the period polynomial is*

$$r_f(X, Y) := (2\pi i)^{n+1} \sum_{\ell=0}^n (-1)^\ell \binom{n}{\ell} i^{\ell+1} L^*(f, \ell+1) X^{n-\ell} Y^\ell. \quad (18)$$

Proposition 2.4 (Period polynomial as a contour integral). *With $(X - \tau Y)^n$ in place of the L -weight τ^{s-1} , and $\mathbf{K}_T(\tau; X, Y) := \sum_{\ell=0}^n \binom{n}{\ell} (-1)^\ell \tilde{k}_T(\tau, \ell+1) X^{n-\ell} Y^\ell$ in place of $\tilde{k}_T$, the same two segments assemble r_f directly:*

$$r_f = (2\pi i)^{n+1} \left[\int_{\gamma^S} f(\tau) (X - \tau Y)^n d\tau + \int_{\gamma^T} f(\tau) \mathbf{K}_T(\tau; X, Y) d\tau \right]. \quad (19)$$

Proof. By Definition 1.3, the $X^{n-\ell} Y^\ell$ coefficient of the bracket is

$$r_f|_{X^{n-\ell} Y^\ell} = (-1)^\ell \binom{n}{\ell} \left[\int_{\gamma^S} f \tau^\ell + \int_{\gamma^T} f \tilde{k}_T(\tau, \ell+1) \right] = (-1)^\ell \binom{n}{\ell} i^{\ell+1} L^*(f, \ell+1). \quad (20)$$

This matches definition 2.3 term by term, and proves the claim. $\square$

Quasi-periods and their determinant: As an aside, the Hurwitz T-kernel in this note was originally uncovered in order to reproduce the manifestly finite computations of the quasi-periods for $\widehat{\Delta} = 1/q + \mathcal{O}(q^2) \in S_{12}^!$ originally given in [2]. In this vein, it is interesting to show collect quasi-periods computable from our L-integrals/period-polynomials for $f \in S_k^!$ (when $\dim(S_k) = 1$).

To this effect, we recall the notation of [5]. Decompose the modular weight $k = 12\lambda + k'$ where $k' \in \{0, 4, 6, 8, 10, 14\}$; thus $\lambda = \dim(S_k)$. For any λ , we may denote $f_{k,m} := 1/q^m + \mathcal{O}(q^{\lambda+1}) \in M_k^!$. When $\dim(S_k) = \lambda = 1$, we have $f_{k,m} = 1/q^m + \mathcal{O}(q^2)$. Here, define $\Delta_k := f_{k,-1} = q + \mathcal{O}(q^2) \in S_k$ and $\widehat{\Delta}_k := f_{k,1} = 1/q + \mathcal{O}(q^2) \in S_k^!$. Further, define

$$\text{pd}_k := \begin{pmatrix} \omega(\Delta_k)^+ & \omega(\Delta_k)^- \\ \eta(\widehat{\Delta}_k)^+ & \eta(\widehat{\Delta}_k)^- \end{pmatrix} \quad (21)$$

As shown in [2], the determinant of pd_k satisfies a Legendre-type relation, $\det(\text{pd}_k) \in \mathbb{Q}^\times \cdot (2\pi i)^{k-1}$; when $k = 12$ this determinant is equal to $-(k-2)! (2\pi i)^{k-1}$. We reproduce this result, and analogous rational multiples in $\mathbb{Q}$ for $k \in \{146, 18, 20, 22, 26\}$, in Table 1, below:

k	$\omega^+(f_k)$	$\omega^-(f_k)/i$	$\eta^+(\widehat{\Delta}_k)$	$\eta^-(\widehat{\Delta}_k)/i$	$\frac{-i^k \det(\text{pd}_k)}{(2\pi)^{k-1} (k-2)!}$
12	-6.89168e7	-5.58502e6	1.27202e11	1.02767e10	1
16	-8.81481e11	-1.54063e10	4.14120e13	6.83541e11	13/30
18	-5.07833e13	-5.43305e12	-2.19200e14	-3.36339e13	2/3
20	-4.22882e16	-3.73170e14	6.38640e16	-1.89823e14	17/5
22	-1.49974e18	-4.24489e16	1.43795e18	-4.41050e16	19/21
26	-1.18673e22	-3.60961e21	1.18653e22	-3.61016e21	23/15

Table 1: Periods $\omega^\pm(\Delta_k)$ and quasi-periods $\eta^\pm(\widehat{\Delta}_k)$, and the rational $(-1)^{1+k/2} \det / ((2\pi i)^{k-1} (k-2)!)$ with $\det = \omega^+ \eta^- - \omega^- \eta^+$. Our rational results for the $\det(\text{pd}_k)$ match those in [2], which presented the results for $k = 12$. We are unaware what governs the sequence of non-unity ratios in the $\det(\text{pd}_k)$ -column of this table. Full 50-digit values in [4]

3 Meromorphic modular forms

Proposition 3.1 (Basepoint and path independence within a homotopy class). *For $f \in F_k$, the value $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ is independent of τ_0 and of the path so long as both vary within a single homotopy class of admissible contours in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$. For $f \in M_k^!$ there is one class and Proposition 1.6 is recovered; the dependence between classes is the residue crossing of Corollary 3.10.*

Proof. The τ_0 -derivative of Proposition 1.6 uses only the T -periodicity $f(\tau_0 - 1) = f(\tau_0)$ and the modularity $f(-1/\tau_0) = \tau_0^k f(\tau_0)$, both valid for every $f \in F_k$, and is evaluated at the moving endpoints alone. While τ_0 stays in one chamber of $\mathbb{H} \setminus f^{-1}(\infty)$ — where the segments meet no pole — the boundary terms cancel exactly as there, so $\partial_{\tau_0} L^* = 0$; the interior poles lie off the contour and never enter. Within a fixed class the integrands $f \tau^{s-1}$ and $f \tilde{k}_T$ are holomorphic along the contours, so any two are joined by a deformation avoiding $\Gamma \cdot f^{-1}(\infty)$, and Cauchy's theorem equates their integrals. $\square$

Definition 3.2 (Polylog projection). *Let $f \in F_k$ have interior poles at the points τ_p of a set $\mathcal{P}_f$ of Γ -inequivalent representatives, and write $\text{Li}_s(z) = \sum_{n>0} z^n n^{-s}$. Since $\text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)})$ has a pole of order m at τ_p , the principal part of f there is matched by a unique polylog projection*

$$\hat{P}_{\tau_p} f(\tau) := \sum_{m>0} r_{f, \tau_p}^*(m) \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)}), \quad (22)$$

its coefficients $r_{f, \tau_p}^(m)$ being the principal-part data of f at τ_p . The projection $\hat{P} := \sum_{\text{Im } \tau_p \geq 1} \hat{P}_{\tau_p}$ removes the principal part of every pole on or above the line $\text{Im } \tau = 1$.*

Definition 3.3 (Regularized cusp coefficients). *With $\hat{P}$ as in Definition 3.2,*

$$c_{Rf}(n) := [(I - \hat{P})f]_{q^n}. \quad (23)$$

Definition 3.4 (The space $F(\infty)$). *$F(\infty)$ is the space of meromorphic g on $\mathbb{H}$ with $g(\tau) = g(\tau + 1)$, finitely many poles in any compact subset of the strip $\{\text{Im } \tau > 0, |\text{Re } \tau| \leq \frac{1}{2}\}$, and $g(\tau) = O(e^{C \text{Im } \tau})$ as $\text{Im } \tau \rightarrow \infty$ for some C . Each such g has a Fourier expansion $g = \sum_n c_g(n) q^n$.*

Then $F_k \subset F(\infty)$, and $(I - \hat{P})f \in F(\infty)$ — the polylog tails are T -periodic but not modular, so the projection leaves F_k .

Lemma 3.5 (q -series–integral commutation). *Let $g \in F(\infty)$ have no pole on or above the line $\text{Im } \tau = 1$ (e.g. $g = (I - \hat{P})f$). Then $g = \sum_n c_g(n) q^n$ converges along the T -segment at $\tau_0 = i$ and integrates term by term:*

$$\int_{\gamma_i^T} g(\tau) \tilde{k}_T(\tau, s) d\tau = \sum_n c_g(n) \int_{i-1}^i e^{2\pi i n \tau} \tilde{k}_T(\tau, s) d\tau. \quad (24)$$

A pole at $\text{Im } \tau_p \geq 1$ defeats this — there the coefficients grow at least like $n^{m-1} e^{2\pi n \text{Im } \tau_p}$ [3, Lem. 2.6].

Proof. With no pole on or above $\text{Im } \tau = 1$, the finitely many poles of g in the strip have heights < 1 ; let $h < 1$ be the largest. The Fourier series $g = \sum_n c_g(n) q^n$ converges absolutely and uniformly on compact subsets of $\{\text{Im } \tau > h\}$, in particular on the T -segment $[i-1, i]$; with $\tilde{k}_T(\tau, s)$ bounded there, the sum passes through the integral, which is the displayed identity. Conversely a pole at height $\text{Im } \tau_p \geq 1$ forces $|c_g(n)| \gtrsim n^{m-1} e^{2\pi n \text{Im } \tau_p}$ [3, Lem. 2.6], so $|c_g(n) q^n| \gtrsim n^{m-1} e^{2\pi n (\text{Im } \tau_p - 1)}$ $\not\rightarrow 0$ on $[i-1, i]$ and the series diverges. $\square$

Definition 3.6 (Polylog integral). For $N \geq 0$ and $z \in \mathbb{H}$, the polylog integral is the L -integral of a single tail $\text{Li}_{-N}(e^{2\pi i(\tau-z)})$ along the T -segment at $\tau_0 = i$ (where γ^S degenerates, since $-1/i = i$):

$$I_N(s, z) := e^{-\pi i s/2} \int_{i-1}^i \text{Li}_{-N}(e^{2\pi i(\tau-z)}) \tilde{k}_T(\tau, s) d\tau. \quad (25)$$

Lemma 3.7 (Closed form of the polylog integral). Write $\mathcal{G}(s, m) := \frac{\Gamma(s, 2\pi m)}{(2\pi m)^s} + i^k \frac{\Gamma(k-s, 2\pi m)}{(2\pi m)^{k-s}}$, the building block of Theorem 2.2 continued to $m < 0$. For the projected poles $\text{Im } z > 1$, where $|e^{2\pi i(\tau-z)}| \geq 1$ on the contour, polylog inversion $\text{Li}_{-N}(w) = (-1)^{N+1} \text{Li}_{-N}(1/w)$ resums the integral into the convergent series

$$I_N(s, z) = \delta_{N,0} \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + (-1)^{N+1} \sum_{n \geq 1} n^N e^{2\pi i n z} \mathcal{G}(s, -n). \quad (26)$$

Proof. For $\text{Im } z > 1$, $|e^{2\pi i(\tau-z)}| \geq 1$ on $[i-1, i]$, so the series of $\text{Li}_{-N}(e^{2\pi i(\tau-z)})$ diverges; the inversion rewrites it through $\text{Li}_{-N}(e^{2\pi i(z-\tau)}) = \sum_{n \geq 1} n^N e^{2\pi i n(z-\tau)}$, convergent since $|e^{2\pi i(z-\tau)}| \leq 1$. Integrating term by term against $\tilde{k}_T$, each $e^{-\pi i s/2} \int_{i-1}^i e^{-2\pi i n \tau} \tilde{k}_T d\tau$ is the building block $\mathcal{G}(s, -n)$ of Theorem 2.2 at mode $-n$, giving the displayed sum; for $N = 0$ the leftover -1 in $\text{Li}_0(w) = -1 - \text{Li}_0(1/w)$ adds $-e^{-\pi i s/2} \int_{i-1}^i \tilde{k}_T = \frac{1}{s} + \frac{i^k}{k-s}$. $\square$

Theorem 3.8 (L -function of a meromorphic modular form). For $f \in F_k$ with no pole on the edge $\text{Im } \tau = 1$, at $\tau_0 = i$,

$$L^*(f, s) = -c_{Rf}(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n \neq 0} c_{Rf}(n) \mathcal{G}(s, n) + \sum_{\text{Im } \tau_p > 1} \sum_{m > 0} r_{f, \tau_p}^*(m) I_{m-1}(s, \tau_p), \quad (27)$$

$\mathcal{G}(s, n) = \frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}}$; the regularized cusp sum (Definition 3.3, Theorem 2.2) plus one polylog integral (Lemma 3.7) per interior pole above the edge. Poles on the edge enter by the principal value of Lemma 3.9.

Proof. At $\tau_0 = i$ the S -segment degenerates, so $L^*(f, s) = e^{-\pi i s/2} \int_{i-1}^i f \tilde{k}_T d\tau$. Split $f = (I - \hat{P})f + \hat{P}f$. The regularized part has no pole on or above $\text{Im } \tau = 1$ (Definition 3.2), so by Lemma 3.5 its Fourier series integrates term by term; mode by mode this is Theorem 2.2, giving $-c_{Rf}(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n \neq 0} c_{Rf}(n) \mathcal{G}(s, n)$. The projected part $\hat{P}f = \sum_{\tau_p} \sum_{m > 0} r_{f, \tau_p}^*(m) \text{Li}_{1-m}(e^{2\pi i(\tau-\tau_p)})$ integrates to $\sum_{\text{Im } \tau_p > 1} \sum_{m > 0} r_{f, \tau_p}^*(m) I_{m-1}(s, \tau_p)$ by Lemma 3.7. Their sum is the stated formula. $\square$

Lemma 3.9 (L -function, a pole on the edge). Routing the T -path just below versus just above a pole τ_p of $f \in F_k$ at height $\text{Im } \tau_p = 1$ gives finite values differing by the residue,

$$L_{\text{below}}^*(f, s) - L_{\text{above}}^*(f, s) = 2\pi i e^{-\pi i s/2} \text{Res}_{\tau=\tau_p} [f(\tau) \tilde{k}_T(\tau, s)]. \quad (28)$$

We define the edge value as the principal value $L^* = \frac{1}{2}(L_{\text{above}}^* + L_{\text{below}}^*) = L_{\text{below}}^* - \pi i e^{-\pi i s/2} \text{Res}_{\tau=\tau_p} [f \tilde{k}_T]$. At the elliptic point $\tau_p = i$ the pole lies on the S -segment instead; its principal value, reached as $\tau_0 \rightarrow i$, is finite and basepoint-independent.

Proof. The two routings of γ^T differ by a contour encircling τ_p once, so the residue theorem gives $L_{\text{below}}^* - L_{\text{above}}^* = 2\pi i e^{-\pi i s/2} \text{Res}_{\tau=\tau_p} [f \tilde{k}_T]$, and the principal value is their mean. At $\tau_p = i$ the pole sits on the degenerate S -segment; displacing to $\tau_0 = i(1+\delta)$ keeps it on γ^S for all small $\delta > 0$, so no wall is crossed and L^* is finite and δ -independent (verified numerically [4]; a closed proof is the S -segment principal value of [3]). $\square$

Corollary 3.10 (Wall-crossing). *By Proposition 3.1, $L^*(f, s)$ is fixed within each homotopy class. Crossing a wall, where the contour sweeps past an interior pole τ_p , changes it by*

$$\Delta L^*(f, s) = 2\pi i e^{-\pi i s/2} \operatorname{Res}_{\tau=\tau_p} [f(\tau) \tilde{k}_T(\tau, s)] \quad (29)$$

when the T -segment crosses τ_p (and by the same with τ^{s-1} in place of $\tilde{k}_T$ when the S -segment does). Thus L^ is fixed by the contour up to the residues it crosses.*

Proof. The within-class constancy is Proposition 3.1. Crossing a wall, the contour sweeps once past τ_p ; the residue theorem then adds $2\pi i e^{-\pi i s/2} \operatorname{Res}_{\tau=\tau_p} [f \tilde{k}_T]$ when the T -segment crosses it, or the same with τ^{s-1} when the S -segment does. $\square$

References

- [1] K. Bringmann, K.-H. Fricke, and Z. Kent, *Special L -values and periods of weakly holomorphic modular forms*, Ramanujan J. **24** (2011), 119–139.
- [2] F. Brown, *A class of non-holomorphic modular forms III*, arXiv:1710.07912 (2017).
- [3] D. A. McGady, *L -functions for meromorphic modular forms and sum rules in conformal field theory*, arXiv:1806.09874 (2018).
- [4] *Numerical verification scripts* (companion code), 2026.
- [5] W. Duke and P. Jenkins, *On the zeros and coefficients of certain weakly holomorphic modular forms*, Pure Appl. Math. Q. **4** (2008), no. 4, 1327–1340.
- [6] R. Bruggeman, Y. Choie, and N. Diamantis, *Holomorphic automorphic forms and cohomology*, Mem. Amer. Math. Soc. **253** (2018), no. 1212; arXiv:1404.6718.
- [7] K. Bringmann, P. Guerzhoy, Z. Kent and K. Ono, *Eichler-Shimura Theory for Mock Modular Forms*, Mathematische Annalen, March 2013 355(3).
- [8] W. Kohnen and D. Zagier, *Modular forms with rational periods*, in *Modular Forms* (R. A. Rankin, ed.), Ellis Horwood, 1984, pp. 197–249.
- [9] V. Paşol and A. A. Popa, *Modular forms and period polynomials*, Proc. London Math. Soc. (3) **107** (2013), 713–743; arXiv:1202.5802.
- [10] W. Wang and H. Zhang, *Meromorphic quasi-modular forms and their L -functions*, arXiv:2202.09542 (2022).
- [11] F. Brown and T. J. Fonseca, *Single-valued periods of meromorphic modular forms and a motivic interpretation of the Gross–Zagier conjecture*, arXiv:2508.04844 (2025).
- [12] K. Bringmann, N. Diamantis, and S. Ehlen, *Regularized inner products and errors of modularity*, Int. Math. Res. Not. IMRN (2016); arXiv:1603.03056.
- [13] N. Diamantis and J. Drewitt, *Period functions associated to real-analytic modular forms*, Res. Math. Sci. **7** (2020), no. 21.
- [14] T. M. Apostol, “Modular functions and Dirichlet series in number theory,” Graduate Texts in Mathematics, London, Springer (1997).