

L-functions for all meromorphic $\mathrm{SL}_2(\mathbb{Z})$ modular forms, via manifestly finite contours

David A. McGady

Abstract

We give a single manifestly finite, regulator-free functional realizing in one object L-functions quasi-periods and period polynomials of meromorphic modular forms on $\mathrm{SL}_2(\mathbb{Z})$, with generic poles throughout the upper half-plane $\mathbb{H}$ and at the cusp of $\mathrm{SL}_2(\mathbb{Z})$. Additionally, our construction preserves holomorphicity. It is a two-segment contour integral, $L^*(f, s) = L_T^*(f, s; \gamma_{\tau_0}^T) + L_S^*(f, s; \gamma_{\tau_0}^S)$, where L_T^* and L_S^* have distinct integration kernels, and $\gamma_{\tau_0}^S$ is a path in $\mathbb{H}$ from $S(\tau_0)$ to τ_0 , and the same for $\gamma_{\tau_0}^T$. For weakly holomorphic forms our L-function is independent of τ_0 and of the path, and reproduces classical L-functions for holomorphic modular forms, reproduces the L-functions of weak modular forms in [1], which previously needed non-holomorphic regularization to be defined, and reproduces Brown's quasi-periods [2]. For properly meromorphic modular forms with poles inside the interior of $\mathbb{H}$, our L-function depends on the homotopy classes, defined in terms of pole-crossings. There is a natural choice of trivial homotopy class, which renders our L-functions canonical when the modular form has a pole in the interior of the fundamental domain, generalizing [3].

1 Introduction and preliminary concepts

Our primary goal in this note is to put forth a unifying generalization of the L-function associated to a given modular form. Our L-function naturally encompasses the classical case of L-functions associated to holomorphic modular forms [4], and seamlessly unifies Ref. [1]'s extension to L-functions of modular forms with poles at the cusp with Ref. [3]'s extension to L-functions of modular forms with poles at finite locations, exclusively away from cusps. Secondly, we show that our L-functions reproduce the numerical results for quasi-periods associated to weak cusp forms [2].

Note: Throughout this document, we refer to L-integrals and L-functions. The classical L-function-as-Dirichlet-series which has been suitably extended to the complex s plane is simply not defined when the modular form that defines the L-function has poles either at the cusp or at finite locations within the upper half-plane. So, if there is ever a doubt, we state clearly: *any L-function in this note is defined by an L-integral*. The focus of this note is to define L-integrals for any integer weight $\mathrm{SL}_2(\mathbb{Z})$ modular form which may have poles at cusps or away from cusps.

1.1 Modular forms

Write $\mathrm{Re}(z)$ and $\mathrm{Im}(z)$ for the real and imaginary parts of a complex number $z \in \mathbb{C}$. Let $\mathbb{H} := \{\tau \in \mathbb{C} \mid \mathrm{Im}(\tau) > 0\}$ denote the upper half-plane. $\mathrm{SL}_2(\mathbb{Z})$ is the group generated by $T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $S := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

There is a natural $\mathrm{SL}_2(\mathbb{Z})$ action on functions $f : \mathbb{H} \rightarrow \mathbb{C}$, graded by integers k : $(f|_k \gamma)(\tau) := (c\tau + d)^{-k} f(\gamma \cdot \tau)$ for all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$. Similarly, $\gamma \cdot \tau := \frac{a\tau + b}{c\tau + d}$.

In this vein, a *holomorphic modular form* of weight k for $\mathrm{SL}_2(\mathbb{Z})$ is a holomorphic function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that $(f|_k \gamma)(\tau) = f(\tau)$ for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ (aka $f(\tau + 1) = f(-\frac{1}{\tau})\tau^{-k} = f(\tau)$) for all

$\tau \in \mathbb{H}$, and $f(\tau) = \mathcal{O}(\tau^k)$ as $\text{Im}(\tau) \rightarrow \infty$, and which has no poles at finite locations within $\mathbb{H}$. By virtue of the periodicity $f(\tau) = f(\tau + 1)$ and holomorphicity, each holomorphic modular form has a q -series decomposition,

$$f(\tau) = \sum_n c_f(n) e^{2\pi i n \tau} = \sum_n c_f(n) q^n \quad , \quad q := e^{2\pi i \tau} \quad . \quad (1)$$

In order for f to be bounded as $\tau \rightarrow i\infty$, we conclude that $c_f(n) = 0$ for all $n < 0$, i.e. that $f = \sum_{n \geq 0} c_f(n) q^n$; for such modular forms f we write $f \in M_k$. If a modular form f additionally has a vanishing constant term, i.e. if $c_f(0) = 0$, then f is a *cusp form*, and we write $f \in S_k$. Clearly, $S_k \subset M_k$. Further, M_k is a finite-dimensional vector-space, as is S_k . Holomorphic $\text{SL}_2(\mathbb{Z})$ modular forms must have even integer weights, and $k \geq 4$ [4].

A product of modular forms of weights k_1 and k_2 is a modular form of weight $k_1 + k_2$, so $M_* := \bigoplus_k M_k$ is naturally a (graded) ring. This graded ring is freely generated by the two modular forms $E_4 \in M_4$ and $E_6 \in M_6$, the $k = 4, 6$ specializations of the *weight k Eisenstein series* E_k , which may be defined as follows:

$$E_k := \frac{1}{2} \sum_{(m,n)=1} \frac{1}{(m+n\tau)^k} = 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \quad , \quad (2)$$

where $\sum_{(m,n)=1}$ denotes summing over all relatively prime $(m, n) \in \mathbb{Z}^2$, the Bernoulli number B_n is the coefficient of $t^n/n!$ of the Laurent expansion of $t/(e^t - 1)$ about $t = 0$, and $\sigma_m(n) := \sum_{d|n} d^m$ is the Divisor function where $d|n$ means $n/d \in \mathbb{N}$. In short, $M_* = \mathbb{C}[E_4, E_6]$. We now define the *modular discriminant*:

$$\Delta := \frac{E_4^3 - E_6^2}{1728} = q + O(q^2) \in S_{12} \quad . \quad (3)$$

Holomorphicity is most canonically relaxed via allowing a modular form to have a pole at the cusp of $\text{SL}_2(\mathbb{Z})$, i.e. for a weight k modular form for $\text{SL}_2(\mathbb{Z})$ to have negative powers of q in its q -series expansion. In this case, f diverges as $\tau \rightarrow i\infty$ as $f \sim e^{-2\pi N \tau}$, where $-N$ is the largest negative number such that $c_f(n)$ is non-zero. Such a modular form f is called a *weak modular form*, and we write $f \in M_k^!$. If a weak modular form f has a vanishing coefficient of q^0 , i.e. if $c_f(0) = 0$, it is called a *weak cusp form* and we write $f \in S_k^!$. Clearly, $S_k^! \subset M_k^!$, $S_k \subset S_k^!$ and $M_k \subset M_k^!$. Two important weak cusp forms in this note are as follows:

$$J := \frac{E_4^3}{\Delta} - 744 = \frac{1}{q} + O(q) \in S_0^! \quad , \quad (4)$$

$$\widehat{\Delta} = (J^2 + 24J - 393444)\Delta := \frac{1}{q} + O(q^2) \in S_{12}^! \quad . \quad (5)$$

In this work, we further relax from even weakly holomorphic modular forms, and consider *meromorphic modular forms* of weight k which are quotients: $f = \frac{f_1}{f_2}$ where $f_i \in M_{k_i}$, the denominator f_2 is not identically zero, and $k = k_1 - k_2$. We write F_k for the vector space of meromorphic modular forms of weight k , and set $F_* := \bigoplus_k F_k$. Then F_k vanishes unless k is an even integer, and F_* may be regarded as a “graded field of fractions” of M_* .

1.2 L-functions for holomorphic modular forms and periods for cusp forms

For any $f \in S_k$, one may define the associated L-function:

$$L(f, s) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s) \stackrel{f \in S_k}{:=} \frac{(2\pi)^s}{\Gamma(s)} \int_0^\infty f(it) t^{s-1} dt = \sum_{n \geq 1} \frac{c_f(n)}{n^s} \quad . \quad (6)$$

A similar definition applies for $f \in M_k$. (The classical definition of $L^*(f, s)$ for $f \in M_k$ is the focus of Remark 3.6, below.) Similarly, one may define a period polynomial for $f \in S_k$ in terms of this L-function,

$$r_f(X, Y) := (2\pi i)^{k-1} \sum_{\ell=0}^{k-2} (-1)^\ell \binom{k-2}{\ell} i^{\ell+1} L^*(f, \ell+1) X^{k-2-\ell} Y^\ell. \quad (7)$$

This polynomial, $r_f(X, Y) \in V_{k-2} := \mathbb{C}[X, Y]_{k-2}$, has a natural action under $\mathrm{SL}_2(\mathbb{Z})$, $(P|_\gamma)(X, Y) = P(aX + bY, cX + dY)$ for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$. This period polynomial can be decomposed into a basis of highly constrained polynomials, which are annihilated by $|(1 + S)$ and $|(1 + U + U^2)[6]$, where $U := TS$ generates an order-3 cyclic subgroup of $\mathrm{SL}_2(\mathbb{Z})$. We define the *periods* of a cusp form to be coefficients of r_f within this highly constrained sub-space W of V_{k-2} , i.e. $W \subset V_{k-2}$.

Why this work? As classically computed, these L-functions and associated period polynomials (when defined in terms of the special values of the associated L-function) are not defined when f has poles either at the cusp of $\mathrm{SL}_2(\mathbb{Z})$ or within the interior of $\mathbb{H}$. As stated above, there are extant, but partial and non-overlapping, extensions of the definition of $L(f, s)$ to $f \in M_k^!$ [1] or $f \in F_k$ when f is regular at the cusp of $\mathrm{SL}_2(\mathbb{Z})$ [3]. In this note, we provide a construction that unifies these disparate extensions. We start with generic L-integrals in section 2, then specialize to $f \in M_k^!$ in section 3, and finally explore the more general context of $f \in F_k$ in section 4.

2 A finite L-integral

Definition 2.1. Let $\tau_0 \in \mathbb{H}$ and $f \in F_k$, let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ denote the generators of $\Gamma = \mathrm{SL}_2(\mathbb{Z})$, and let $f^{-1}(\infty)$ be the locus of all poles of $f \in \mathbb{H}$ (inclusive of the cusp of $\mathrm{SL}_2(\mathbb{Z})$). The S - and T -segments at τ_0 are smooth paths in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$ from $\sigma^{-1}\tau_0$ to τ_0 , for $\sigma = S$ and $\sigma = T$ respectively,

$$\gamma_{\tau_0}^S : -1/\tau_0 \longrightarrow \tau_0, \quad \gamma_{\tau_0}^T : \tau_0 - 1 \longrightarrow \tau_0. \quad (8)$$

Avoiding poles is automatic for $f \in M_k^!$, as long as the base-point τ_0 , is not at a cusp of $\mathrm{SL}_2(\mathbb{Z})$, i.e. there is only the trivial homotopy class. When $f \in F_k$, and f has poles at finite and non-zero imaginary heights, integration contours fall into nontrivial homotopy classes.

Definition 2.2. Let $\zeta(s, x) = \sum_{m \geq 0} (m + x)^{-s}$ as a function $\zeta : \mathbb{C}_s \times (\mathbb{C} \setminus \mathbb{N})_x \rightarrow \mathbb{C}$ denote the Hurwitz zeta function (meromorphic in s , holomorphic in $x \in \mathbb{H}$). We define Hurwitz T -kernel as

$$\tilde{k}_T(\tau, s) := \zeta(1 - s, \tau + 1) - e^{i\pi(s-1)} \zeta(1 - (k - s), \tau + 1). \quad (9)$$

Definition 2.3 (Generic L-integral). Let $f \in F_k$, $\tau_0 \in \mathbb{H}$, and $s \in \mathbb{C}$. Further, let $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$ be the S - and T -segments at τ_0 , as in Definition 2.1, and let $\tilde{k}_T$ be as defined in Definition 2.2. The L-integral of f is

$$L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) := L_S^*(f, s; \gamma_{\tau_0}^S) + L_T^*(f, s; \gamma_{\tau_0}^T) := e^{-\pi i s/2} \left(\int_{\gamma_{\tau_0}^S} f(\tau) \tau^{s-1} d\tau + \int_{\gamma_{\tau_0}^T} f(\tau) \tilde{k}_T(\tau, s) d\tau \right) \quad (10)$$

We use this L -integral to define a generalized L -function:

Definition 2.4 (Generic L-function). Let $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ be as defined in Definition 2.3. We define the generalized L-function of $f \in F_k$ to be,

$$L(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T). \quad (11)$$

Having defined the L-integral and L-function, we prove they are manifestly finite for $f \in F_k$:

Lemma 2.5 (Finiteness of L-integrals). *For $f \in F_k$, both $L_S^*(f, s; \gamma_{\tau_0}^S)$ and $L_T^*(f, s; \gamma_{\tau_0}^T)$ in Definition 2.3 are each, separately finite for every $s \in \mathbb{C}$, every $\tau_0 \in \mathbb{H}$, and for every path $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$.*

Proof. Both segments are compact and lie in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, so f is continuous on them; τ^{s-1} is continuous on $\mathbb{H}$, and $\tilde{k}_T(\tau, s)$ is continuous there since $\zeta(\nu, \tau + 1)$ is holomorphic in $\tau \in \mathbb{H}$ (Definition 2.2). A continuous integrand on a compact path is integrable, for every $s \in \mathbb{C}$. $\square$

We now turn to path-dependence and τ_0 -independence of the L-integral and the L-function of $f \in F_k$. This holds for all $f \in M_k^!$:

Proposition 2.6 (Basepoint & path independence for $f \in M_k^!$). *For $f \in M_k^!$ the value $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ is independent of $\tau_0 \in \mathbb{H}$ and of the paths: with no interior poles, $\mathbb{H}$ is simply connected, so all admissible contours are homotopic. We write $L^*(f, s)$ for the common value and $L(f, s) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s)$.*

Proof. With no interior poles the integrands $f(\tau)\tau^{s-1}$ and $f(\tau)\tilde{k}_T(\tau, s)$ are holomorphic on $\mathbb{H}$, which is simply connected; by Cauchy's theorem each integral is unchanged under any deformation fixing its endpoints, hence depends on those endpoints alone. This is path-independence.

For independence of τ_0 show that variations in τ_0 for the S-term and T-term in $L^*(f, s)$ cancel, i.e. that $\partial_{\tau_0} L_S^*(f, s; \gamma_{\tau_0}^S) = -\partial_{\tau_0} L_T^*(f, s; \gamma_{\tau_0}^T)$. For L_T^* , note that the T -segment runs from $T^{-1}\tau_0 = \tau_0 - 1$ to τ_0 , so by Leibniz's rule and the T -periodicity $f(\tau_0 - 1) = f(\tau_0)$,

$$\partial_{\tau_0} L_T^*(f, s; \gamma_{\tau_0}^T) = f(\tau_0) \tilde{k}_T(\tau_0, s) - f(\tau_0 - 1) \tilde{k}_T(\tau_0 - 1, s) = f(\tau_0) \delta_T \tilde{k}_T(\tau_0 - 1, s), \quad (12)$$

where $\delta_T \tilde{k}_T(\tau, s) := \tilde{k}_T(\tau + 1, s) - \tilde{k}_T(\tau, s)$. This is the whole role of δ_T : the unit gap between the endpoints turns the ∂_{τ_0} boundary term into a unit shift of the kernel. That shift is elementary, since $\zeta(\nu, x + 1) - \zeta(\nu, x) = -x^{-\nu}$ gives, on both terms at $x = \tau + 1$,

$$\delta_T \tilde{k}_T(\tau, s) = -(\tau + 1)^{s-1} + e^{i\pi(s-1)} (\tau + 1)^{k-1-s} \quad (13)$$

$$\implies \partial_{\tau_0} L_T^*(f, s; \gamma_{\tau_0}^T) = f(\tau_0) [-\tau_0^{s-1} + e^{i\pi(s-1)} \tau_0^{k-1-s}]. \quad (14)$$

The S -segment runs $-1/\tau_0 \rightarrow \tau_0$; its lower endpoint moves with $\partial_{\tau_0}(-1/\tau_0) = \tau_0^{-2}$, so by Leibniz, the modularity $f(-1/\tau_0) = \tau_0^k f(\tau_0)$, and the phase $(-1/\tau_0)^{s-1} = e^{i\pi(s-1)} \tau_0^{1-s}$,

$$\partial_{\tau_0} L_S^*(f, s; \gamma_{\tau_0}^S) = f(\tau_0) \tau_0^{s-1} - f(-1/\tau_0) (-1/\tau_0)^{s-1} \tau_0^{-2} = f(\tau_0) [\tau_0^{s-1} - e^{i\pi(s-1)} \tau_0^{k-1-s}]. \quad (15)$$

Thus τ_0 -variations are equal and opposite, $\partial_{\tau_0} L_S^* = -\partial_{\tau_0} L_T^*$. Thus $\partial_{\tau_0} L^* = \partial_{\tau_0} L_S^* + \partial_{\tau_0} L_T^* = 0$. $\square$

Remark 2.7 (Recovering the classical L-integral for $f \in S_k$). Let $f \in S_k$. As $f^{-1}(\infty)$ does not include the cusp $i\infty$, $\lim_{\tau_0 \rightarrow i\infty} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ exists. In this limit, the S -segment maps onto the standard L-function contour, $[0, i\infty]$. Simultaneously, as $f(it) \sim e^{-2\pi t}$ as $t \rightarrow \infty$, the T -segment vanishes exponentially. Thus, in this limit, $L^*(f, s)$ exactly maps onto the classical L-function integral [4]:

$$\lim_{\tau_0 \rightarrow i\infty} L^*(f, s) = \int_0^\infty t^{s-1} f(it) dt. \quad (16)$$

3 L-functions, period polynomials and poles at the cusp

Proposition 3.1. *Let $\Gamma(s, x) := \int_x^\infty e^{-t} t^{s-1} dt$ denote the incomplete Gamma function. For $n \neq 0$, $\nu \in \mathbb{C}$, and $z \in \mathbb{H}$,*

$$\int_z^{z+1} e^{2\pi i n u} \zeta(\nu, u) du = e^{i\pi(1-\nu)/2} (2\pi n)^{\nu-1} \Gamma(1-\nu, -2\pi i n z). \quad (17)$$

Proof. Write $\zeta(\nu, u) = \sum_{m \geq 0} (m+u)^{-\nu}$ and shift $u \mapsto u+m$ in the m -th term; as $e^{2\pi i n(u+m)} = e^{2\pi i n u}$ for integer m , the unit segments stack into one horizontal ray,

$$\int_z^{z+1} e^{2\pi i n u} \zeta(\nu, u) du = \sum_{m \geq 0} \int_{z+m}^{z+1+m} e^{2\pi i n v} v^{-\nu} dv = \int_z^{z+\infty} e^{2\pi i n v} v^{-\nu} dv. \quad (18)$$

Now, substitute $w = -2\pi i n v$, so that $v = iw/(2\pi n)$, $dv = i dw/(2\pi n)$, and $e^{2\pi i n v} = e^{-w}$: the lower endpoint becomes $w = -2\pi i n z$, and for $n > 0$ the ray rotates to $w \rightarrow +\infty$ along the positive reals. Since $e^{-w} w^{-\nu}$ decays in the right half-plane, the closing arc vanishes. With $v^{-\nu} = i^{-\nu} (2\pi n)^{\nu} w^{-\nu}$,

$$\int_z^{z+\infty} e^{2\pi i n v} v^{-\nu} dv = i^{1-\nu} (2\pi n)^{\nu-1} \int_{-2\pi i n z}^{\infty} e^{-w} w^{-\nu} dw = e^{i\pi(1-\nu)/2} (2\pi n)^{\nu-1} \Gamma(1-\nu, -2\pi i n z). \quad (19)$$

For $n < 0$ the ray instead grows as $\text{Im} v \rightarrow \infty$ and the rotation is illegal; but both sides are analytic in n and agree for $n > 0$, so the identity continues to $n < 0$ with Γ on its principal branch. At $n = 0$ the prefactor $(2\pi n)^{\nu-1}$ is singular; that case is handled separately. $\square$

Theorem 3.2 (The L-function for $f \in M_k^!$). *For $f \in M_k^!$, we may set $\tau_0 = i$, and prove*

$$L^*(f, s) = -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right). \quad (20)$$

Here, trivially, $L^*(f, s) = i^k L^*(f, k-s)$. The L-function is $L(f, s) := (2\pi)^s L^*(f, s)/\Gamma(s)$.

Proof. By path independence and base-point independence from Proposition 2.6, we may freely set $\tau_0 = i$. Here, $L_S^*(f, s; \gamma_{\tau_0}^S) = 0$ (the S -contour vanishes in length and there are no nontrivial homotopy classes as $f \in M_k^!$ has no poles which a finite contour can wrap), and the L-integral simplifies dramatically:

$$L^*(f, s) = L_T^*(f, s; \gamma_i^T) = e^{-\pi i s/2} \int_{i-1}^i f(\tau) \tilde{k}_T(\tau, s) d\tau. \quad (21)$$

We then expand f in terms of its q-series, $\sum_n c_f(n) e^{2\pi i n \tau}$, and substitute the u -variable of Proposition 3.1 as $u = \tau + 1$, and note that $e^{2\pi i n(u-1)} = e^{2\pi i n u}$. When $n \neq 0$, straightforward substitutions of ν, z into Proposition 3.1 suffice: $1-\nu = s$ gives the contribution of the first term in $\tilde{k}_T(\tau, s)$, and $1-\nu = k-s$ gives the second:

$$e^{-\pi i s/2} \left(e^{\pi i s/2} \frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} - e^{i\pi(s-1)} e^{\pi i(k-s)/2} \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right) = \frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}}. \quad (22)$$

We may then evaluate the $n = 0$ mode's contribution follows from noting the closed form solution for the definite integral,

$$\int_{i-1}^i \zeta(\sigma, \tau + 1) d\tau = \int_i^{i+1} \zeta(\sigma, u) du = \frac{i^{1-\sigma}}{\sigma-1}. \quad (23)$$

This result follows from noting $\partial_a(\zeta(\sigma-1, a)/(1-\sigma)) = \zeta(\sigma, a)$ and $\zeta(\sigma-1, a+1) - \zeta(\sigma-1, a) = -a^{1-\sigma}$. Applying this to the two distinct terms of $\tilde{k}_T(\tau, s)$ at $\sigma = 1-s$ and $\sigma = 1-(k-s)$ gives

$$e^{-\pi is/2} \int_{i-1}^i c_f(0) \tilde{k}_T(\tau, s) d\tau = -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right). \quad (24)$$

Combining the $n=0$ contributions with the $n \neq 0$ contributions reproduces Eq. (20). Finally, the functional equation, $L^*(f, s) = i^k L^*(f, k-s)$, follows by direct inspection. $\square$

3.1 Explicit S- and T-contours for general τ_0

Lemma 3.3 (L_T^* for $M_k^!$). For $f \in M_k^!$, and $\tau_0 \in \mathbb{H}$ away from cusps, $L_T^*(f, s; \gamma_{\tau_0}^T)$ evaluates to:

$$L_T^*(f, s; \gamma_{\tau_0}^T) = -c_f(0) \left(\frac{(\tau_0/i)^s}{s} + i^k \frac{(\tau_0/i)^{k-s}}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, -2\pi in\tau_0)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, -2\pi in\tau_0)}{(2\pi n)^{k-s}} \right). \quad (25)$$

Proof. Expand $f = \sum_n c_f(n) q^n$ on the compact segment $[\tau_0 - 1, \tau_0]$, and restrict attention to the sub-sum, $\sum_{n \neq 0}$. For each such $n \neq 0$, apply the Proposition 3.1 in $\tilde{k}_T(\tau, s) = \zeta(1-s, \tau+1) - e^{i\pi(s-1)} \zeta(s-(k-1), \tau+1)$ at $\nu = 1-s$ and $\nu = s-(k-1)$. The phases for the second term work-out identically as in the proof of Theorem 3.2.

For $n=0$, the constant $c_f(0)$ multiplies $e^{-\pi is/2} \int_{\tau_0-1}^{\tau_0} \tilde{k}_T(\tau, s) d\tau$, which we compute as follows. First, the closed-form of the Hurwitz antiderivative $\int_{\tau_0-1}^{\tau_0} \zeta(\sigma, \tau+1) d\tau = \tau_0^{1-\sigma}/(\sigma-1)$ comes from $\partial_\tau \zeta(\sigma-1, \tau+1) = -(\sigma-1)\zeta(\sigma, \tau+1)$ and the shift identity $\zeta(\sigma-1, \tau_0+1) - \zeta(\sigma-1, \tau_0) = -\tau_0^{1-\sigma}$. Finally, note that the two terms in $\tilde{k}_T$ are simply specializations of $\sigma = 1-s$ and $\sigma = s-k+1$, which yield $-e^{-\pi is/2} \tau_0^s/s = -(\tau_0/i)^s/s$ and $-e^{\pi is/2} \tau_0^{k-s}/(k-s) = -i^k (\tau_0/i)^{k-s}/(k-s)$ respectively. $\square$

Corollary 3.4 (L_T^* as $\tau_0 \rightarrow i\infty$). Let $f \in S_k^!$, where $f = \sum_{n \geq -N} c_f(n) q^n$ for some $-N \in \mathbb{Z}$. The T-contour's contribution to $L^*(f, s)$ is given by Lemma 3.3. As $\tau_0 = it$ goes to the cusp $i\infty$, the T-contour scales exactly as $f(it)$ scales: $L_T^*(f, s; \gamma_{\tau_0}^T) \sim f(it) \sim e^{2\pi Nt}$.

Proof. For the asymptotics, $\Gamma(\sigma, z) \sim z^{\sigma-1} e^{-z}$ as $|z| \rightarrow \infty$ for $|\arg z| < \frac{3\pi}{2}$ [10, §8.11] gives $|c_f(n)|$ -weighted go as $e^{-2\pi n \text{Im}(\tau_0)}$. The smallest mode n in the q-series, i.e. $n = -N$, dominates the scaling of $f(it)$ and of $L_T^*(f, s; \gamma_{it}^T)$. Combining these results yields the claimed asymptotics. $\square$

Now, exploiting τ_0 -independence of $L^*(f, s)$, its explicit value as given in Theorem 3.2, and the explicit T-contribution $L_T^*(f, s; \gamma_{\tau_0}^T)$ trivially implies clean, closed-form expressions for $L_S^*(f, s; \gamma_{\tau_0}^S)$:

Corollary 3.5 (L_S^* for $M_k^!$). For $f \in M_k^!$ and $\tau_0 \in \mathbb{H}$, $L_S^*(f, s; \gamma_{\tau_0}^S)$ evaluates to

$$L_S^*(f, s; \gamma_{\tau_0}^S) = c_f(0) \left(\frac{(\tau_0/i)^s - 1}{s} + i^k \frac{(\tau_0/i)^{k-s} - 1}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, 2\pi n) - \Gamma(s, -2\pi in\tau_0)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n) - \Gamma(k-s, -2\pi in\tau_0)}{(2\pi n)^{k-s}} \right). \quad (26)$$

Proof. Recall $L^*(f, s) - L_T^*(f, s; \gamma_{\tau_0}^T) = L_S^*(f, s; \gamma_{\tau_0}^S)$. Now, $L^*(f, s)$ for $f \in M_k^!$ is given by Theorem 3.2, and $L_T^*(f, s; \gamma_{\tau_0}^T)$ is given by Lemma 3.3. Their difference yields Eq. (26). $\square$

3.2 Classical Mellin recovery and canonical uniqueness.

As discussed in Remark 2.7, for $f \in S_k$, taking the cusp-limit $\tau_0 \rightarrow i\infty$ in Theorem 3.2 collapses the S -segment onto the classical Mellin contour $[0, i\infty]$ while via Corollary 3.4 the T -segment vanishes exponentially as $L_T^*(f, s; \gamma_{\tau_0}^T) \sim f(\tau_0) \sim e^{-2\pi\text{Im}(\tau_0)N}$ (as $f \in S_k \Leftrightarrow f = \sum_{n \geq N} c_f(n)q^n$ with $N \geq 1$). I.e., here, one recovers the classical L-integral of f without obstruction:

$$\lim_{\tau_0 \rightarrow i\infty} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) = \int_0^\infty t^{s-1} f(it) dt. \quad (27)$$

This story changes when $f = 1/q^N(1 + O(q)) \in S_k^!$ for some $N \geq 1$.

In short, when f has a polar q-series, it has a pole at the cusp of $\text{SL}_2(\mathbb{Z})$. Thus, the limit $\tau_0 \rightarrow i\infty$, which was crucial to recovering the classical Mellin contour in Remark 2.7 simply does not exist.

However, it can still be instructive to examine $L_T^*(f, s; \gamma_{\tau_0}^T)$, and $L_S^*(f, s; \gamma_{\tau_0}^S)$ as τ_0 goes towards $i\infty$. In this context, while of course the sum $L_T^*(f, s; \gamma_{\tau_0}^T) + L_S^*(f, s; \gamma_{\tau_0}^S) = L^*(f, s)$ is manifestly invariant under changes of the base-point, each of these two terms can indeed change values. Indeed, Corollary 3.4 clearly shows that as $\tau_0 \rightarrow i\infty$, when $f \in S_k^!$ has negative powers of q in its q-series, then $L_T^*(f, s; \gamma_{\tau_0}^T) \sim e^{2\pi N \text{Im}(\tau_0)}$. Discarding $\gamma_{\tau_0}^T$ therefore does violence to the result: When the S -contour is isolated from its T-partner, and $\tau_0 \rightarrow i\infty$, it diverges. This is the precise reason that [1] needed to regularize their divergent L-integral representation for properly weak cusp forms.

That said, there is one further aspect of our L-function even in the classical case of $f \in M_k$ that is worthy of remark:

Remark 3.6 (Recovering the classical L-integral for $f \in M_k$). In textbook treatments [4], when $f \in M_k$, its L-integral is defined as

$$\begin{aligned} L^*(f, s)|_{\text{cl}} &:= \int_0^\infty (f(it) - c_f(0))t^{s-1}dt = \frac{\Gamma(s)}{(2\pi)^s} \sum_{n=1}^\infty \frac{c_f(n)}{n^s} \\ &= -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n=1}^\infty c_f(n) \left(\frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right) \end{aligned} \quad (28)$$

Classically, this constant-term subtraction is needed because $\int_0^\infty t^{s-1}dt$ is a pure divergence, so the q-series cannot commute with the integration contour.

However, as is clear from our general Theorem 3.2 for L-integrals for $f \in M_k^!$, and from the specific results of the S- and T-contours in Lemma 3.3 and Corollary 3.5, our L-integral requires no such explicit subtraction of the constant term $-c_f(0)$. Thus, not only does our prescription naturally handle divergences at poles when $f \in M_k^!$ has polar terms in its q-series, but it also obviates the need to by-hand remove the constant mode from the kernel of the L-integral that is needed when defining L-functions in terms of the classical Mellin contour, $\int_0^\infty$.

3.3 Period polynomials, periods and quasi-periods

Because $L^*(f, s)$ is canonical for every $f \in S_k^!$, and because period polynomials for a given modular form can be written in terms of their L-functions evaluated at integer- s in the critical strip, they need no separate construction.

Definition 3.7 (Period polynomial for $S_k^!$). For $f \in S_k^!$ and $n = k - 2$, the period polynomial is

$$r_f(X, Y) := (2\pi i)^{n+1} \sum_{\ell=0}^n (-1)^\ell \binom{n}{\ell} i^{\ell+1} L^*(f, \ell+1) X^{n-\ell} Y^\ell. \quad (29)$$

Proposition 3.8 (Period polynomial as a contour integral). *Let $n := k - 2$. With $(X - \tau Y)^n$ in place of the L -weight τ^{s-1} , and $\mathbf{K}_T(\tau; X, Y) := \sum_{\ell=0}^n \binom{n}{\ell} (-1)^\ell \tilde{k}_T(\tau, \ell + 1) X^{n-\ell} Y^\ell$ in place of $\tilde{k}_T$, the same two segments assemble r_f directly:*

$$r_f(X, Y) = (2\pi i)^{n+1} \left[\int_{\gamma_{\tau_0}^S} f(\tau) (X - \tau Y)^n d\tau + \int_{\gamma_{\tau_0}^T} f(\tau) \mathbf{K}_T(\tau; X, Y) d\tau \right]. \quad (30)$$

Proof. By Definition 2.3, the $X^{n-\ell} Y^\ell$ coefficient of the bracket is

$$r_f|_{X^{n-\ell} Y^\ell} = (-1)^\ell \binom{n}{\ell} i^{\ell+1} [L_S^*(f, \ell + 1) + L_T^*(f, \ell + 1)] = (-1)^\ell \binom{n}{\ell} i^{\ell+1} L^*(f, \ell + 1). \quad (31)$$

This matches Definition 3.7 term by term, and proves the claim. $\square$

Having now defined the period polynomial associated to a weak cusp form $f \in S_k^!$, we now seek to find its (*quasi*-)periods. First, a number of helpful definitions.

All objects below take values in $V_n := \mathbb{C}[X, Y]_n$, $n = k - 2$, with the slash action $(P|_\gamma)(X, Y) := P(aX + bY, cX + dY)$ of §1. A map $\gamma \mapsto A_\gamma \in V_n$ is a 1-cocycle for $\mathrm{SL}_2(\mathbb{Z})$ if

$$A_{\gamma\delta} = A_\gamma|_\delta + A_\delta \quad \text{for all } \gamma, \delta \in \mathrm{SL}_2(\mathbb{Z}), \quad (32)$$

and is *parabolic* if in addition $A_T = 0$. For fixed $P \in V_n$ the map $\gamma \mapsto P|_\gamma - P$ is a *coboundary*, and satisfies (32) identically, since $(P|_\gamma - P)|_\delta + (P|_\delta - P) = P|_{\gamma\delta} - P$. Adding or subtracting a coboundary from a cocycle therefore returns a cocycle, which is what lets C_γ below be replaced by a representative $\hat{C}_\gamma$ normalised so that $\hat{C}_T = 0$.

For starters, let $f \in S_k^!$, $\tau_0 \in \mathbb{H}$ and $\gamma \in \mathrm{SL}_2(\mathbb{Z})$, and define

$$C_\gamma(f; \tau_0) := (2\pi i)^{n+1} \int_{\gamma^{-1}\tau_0}^{\tau_0} \omega_f, \quad \text{where } \omega_f := f(\tau)(X - \tau Y)^n d\tau. \quad (33)$$

Finally, it is helpful to define two slight abstractions related to $\mathbf{K}_T(\tau; X, Y)$ and to $L_T^*(f, s; \gamma_{\tau_0}^T)$:

$$\chi(\tau; X, Y) := \sum_{\ell=0}^n \binom{n}{\ell} (-1)^\ell \zeta(-\ell, \tau + 1) X^{n-\ell} Y^\ell, \quad (34)$$

$$E(f; \tau_0) := (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) \chi(\tau; X, Y) d\tau. \quad (35)$$

At non-positive integer first argument the Hurwitz zeta is elementary, $\zeta(-\ell, x) = -B_{\ell+1}(x)/(\ell + 1)$ with $B_m(x)$ the m -th Bernoulli polynomial [10, §25.11], so $\chi(\tau; X, Y)$ is a polynomial in τ of degree $n + 1$. Its role is that of a discrete antiderivative of the Eichler kernel: the Hurwitz shift $\zeta(-\ell, \tau) = \zeta(-\ell, \tau + 1) + \tau^\ell$ gives, term by term,

$$\chi(\tau - 1; X, Y) - \chi(\tau; X, Y) = (X - \tau Y)^n. \quad (36)$$

Both facts are used in the proof of Lemma 3.9.

For each τ , $\chi(\tau; \cdot, \cdot) \in V_{k-2} = \mathbb{C}[X, Y]_{k-2}$; the $\mathrm{SL}_2(\mathbb{Z})$ -slash $(P|_\gamma)(X, Y) := P(aX + bY, cX + dY)$ acts on the polynomial variables alone, holding τ fixed:

$$\chi|_\gamma(\tau; X, Y) := \chi(\tau; aX + bY, cX + dY), \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}). \quad (37)$$

Thus $|_T$ sends $(X, Y) \mapsto (X + Y, Y)$ with τ untouched. Being linear and τ -independent, $|_\gamma$ commutes with $\int d\tau$, so slashing the polynomial E agrees with slashing χ under the integral; the two occurrences of $|_\gamma$ in Lemma 3.9 denote the same operation.

Lemma 3.9. *Let $f \in S_k^!$, $\tau_0 \in \mathbb{H}$, and let ω_f , $C_\gamma(f; \tau_0)$, $\chi(\tau; X, Y)$, and $E(f; \tau_0)$ be as defined in Eqs. (33), (34) and (35). Then $\hat{C}_\gamma := C_\gamma - (E|_\gamma - E)$ is a 1-cocycle for $\mathrm{SL}_2(\mathbb{Z})$ with $\hat{C}_S = r_f$ and $\hat{C}_T = 0$.*

Proof. First, we prove $\hat{C}_\gamma$ is a cocycle. Path-additivity and covariance of ω_f give the Eichler–Shimura cocycle relation $C_{\gamma\delta} = C_\gamma|_\delta + C_\delta$ [11, 6]; as $\gamma \mapsto E|_\gamma - E$ is a coboundary, $\hat{C}$ is a cocycle.

Having established it is a cocycle, we now show that $\hat{C}_S = r_f$. By Proposition 3.8 the period polynomial splits along the two segments of Definition 2.1, and we identify each contribution in turn. The S -segment carries the kernel τ^{s-1} , so assembling it over ℓ returns $(2\pi i)^{n+1} \int_{\gamma_{\tau_0}^S} \omega_f = C_S$ exactly.

For the T -segment, at $s = \ell + 1$ Definition 2.2 reads $\tilde{k}_T(\tau, \ell + 1) = \zeta(-\ell, \tau + 1) - (-1)^\ell \zeta(-(n - \ell), \tau + 1)$; assembling $\mathbf{K}_T$ and re-indexing $\ell \mapsto n - \ell$ in the second term (with n even) gives $\mathbf{K}_T = \chi - \chi|_S$, so the T -term of r_f is $E - E|_S$. Hence

$$r_f = C_S + (E - E|_S) = C_S - (E|_S - E) = \hat{C}_S. \quad (38)$$

Neither summand in (38) vanishes in general: the S -segment runs from $-1/\tau_0$ to τ_0 and degenerates only at the fixed point $\tau_0 = i$, where $C_S = 0$ and $E|_S - E = -r_f$.

Finally, we show that $\hat{C}_T = 0$. Applying the shift identity (36) twice shows that $A(\tau) := \chi(\tau; X + Y, Y) - \chi(\tau - 1; X, Y)$ satisfies $A(\tau - 1) = A(\tau)$. Crucially, χ is a polynomial in τ , as recorded below Eq. (35), so A is a 1-periodic polynomial and hence constant: $\partial_\tau A(\tau) = 0 \Leftrightarrow A(\tau) = A_0$. Thus,

$$\begin{aligned} \hat{C}_T &= C_T - (E|_T - E) \\ &= (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) (X - \tau Y)^n d\tau - (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) [\chi|_T(\tau; X, Y) - \chi(\tau; X, Y)] d\tau \\ &= (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) \left([\chi(\tau - 1; X, Y) - \chi(\tau; X, Y)] - [\chi(\tau; X + Y, Y) - \chi(\tau; X, Y)] \right) d\tau \\ &= (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) [\chi(\tau - 1; X, Y) - \chi(\tau; X + Y, Y)] d\tau = -(2\pi i)^{n+1} \int_{\tau_0-1}^{\tau_0} f(\tau) A(\tau) d\tau \\ &= -(2\pi i)^{n+1} A_0 \int_{\tau_0-1}^{\tau_0} f(\tau) d\tau = -(2\pi i)^{n+1} A_0 c_f(0) = 0, \end{aligned} \quad (39)$$

as all elements $f \in S_k^!$ must have $c_f(0) = 0$. $\square$

Definition 3.10. $W := \{w \in \mathbb{C}[X, Y]_{k-2} : w|_{(1+S)} = w|_{(1+U+U^2)} = 0\} \subset V_n$

Lemma 3.11. *For $f \in S_k^!$, and $W \subset V_n$ from Definition 3.10, the period polynomial $r_f(X, Y) \in W$.*

Proof. Let $\hat{C}_\gamma$ be the parabolic cocycle of Lemma 3.9. Thus, $r_f = \hat{C}_S$ and $\hat{C}_T = 0$. Now, note that $\hat{C}_\gamma(f; \tau_0) = 0$ when $\gamma = \pm I$. Since $n = k - 2$ is even, $\pm I$ act trivially on V_n , so the coboundary term $E|_{\pm I} - E$ vanishes and $\hat{C}_{\pm I} = C_{\pm I}$; and $C_{\pm I}$ is a closed-contour integral of a function which has no poles in $\mathbb{H}$. Finally, direct computation, starting from the identities $-I = U^3 = S^2$ and using the cocycle law $\hat{C}_{\gamma\delta} = \hat{C}_\gamma|_\delta + \hat{C}_\delta$ and clever use of $\hat{C}_T = 0$, shows $r_f|_{(1+S)} = r_f|_{(1+U+U^2)} = 0$:

$$\hat{C}_U = \hat{C}_{TS} = \hat{C}_T|_S + \hat{C}_S = \hat{C}_S = r_f, \quad (40)$$

$$0 = \hat{C}_{S^2} = \hat{C}_S|_S + \hat{C}_S = \hat{C}_S|_{(1+S)} = r_f|_{(1+S)}, \quad (41)$$

$$0 = \hat{C}_{U^3} = \hat{C}_U|_{UU} + \hat{C}_{UU} = \hat{C}_U|_{U^2} + (\hat{C}_U|_U + \hat{C}_U) = \hat{C}_U|_{(1+U+U^2)} = r_f|_{(1+U+U^2)}. \quad (42)$$

Thus, $r_f \in W \subset V_n$, and finishes the proof. $\square$

To define periods and quasi-periods, we recall the standard fact that $\dim(W) = 2 \dim(S_k) + 1$ (see, e.g. [6]), and define the inner-product:

$$\langle P, Q \rangle := \sum_a (-1)^a \binom{n}{a}^{-1} p_a q_{n-a} . \quad (43)$$

Now for the basis elements for k where $\dim(S_k) = 1$:

Definition 3.12. Let $\dim S_k = 1$, $n := k - 2$, $p_0(X) := X^n - 1$, $B_\ell := \frac{\ell!}{2\pi i} \oint \frac{t^{-\ell}}{e^t - 1} dt$, and $B_m^0(X) := \sum_{i \neq 1} \binom{m}{i} B_i X^{m-i}$. Motivated by [6, Thm. 1'], we define:

$$W_-^{(k)}(X) := \text{pp} \left(\frac{1}{n-1} (B_{n-1}^0(X) + X^n B_{n-1}^0(\frac{1}{X})) - \frac{1}{3} (B_3^0(X) + X^n B_3^0(\frac{1}{X})) \right), \quad (44)$$

$$W_+^{(k)}(X) := \text{pp} \left(\frac{1}{2} (B_2^0(X) - X^n B_2^0(\frac{1}{X})) + \frac{1}{n} (B_n^0(X) - X^n B_n^0(\frac{1}{X})) - c p_0(X) \right), \quad (45)$$

$$v_0^{(k)}(X) := p_0(X), \quad (46)$$

where pp acts on any polynomial with rational coefficients by multiplying by all coefficient-denominators and dividing by the gcd, and $\langle P, Q \rangle$ is as in Eq. (43). The bracketed polynomial defining $W_+^{(k)}$ has coefficients $a_i = -a_{n-i}$, so the single constant $c := a_n$ eliminates its X^n and constant terms at once; this is the p_0 -component, and removing it is what makes $\langle W_+^{(k)}, p_0 \rangle = 0$ below.

Lemma 3.13. For k with $\dim S_k = 1$, the space W of Lemma 3.11 has $\dim W = 3$, and $\{W_+^{(k)}, W_-^{(k)}, v_0^{(k)}\}$ is an orthogonal basis of W under $\langle \cdot, \cdot \rangle$ from Eq. (43).

Proof. First, let $W_\pm^{(k)}(X, Y) := Y^{k-2} W_\pm^{(k)}(X/Y)$, and let $v_0^{(k)}(X, Y) := Y^{k-2} v_0^{(k)}(X/Y)$.

Membership in W is the substantive point. Theorem 1' of [6] gives the odd and even period polynomials of the cusp form $E_{k-12}\Delta$ (where $E_0 := 1$) as modified-Bernoulli combinations and places them in W ; the $W_\pm^{(k)}$ of Definition 3.12 are specializations of those formulae, $W_-^{(k)}$ odd and $W_+^{(k)}$ even under $X \mapsto -X$, so both lie in W . So does $v_0^{(k)} = p_0 = X^n - Y^n$, directly: $p_0|_S = -p_0$ since n is even, and $p_0|_{(1+U+U^2)} = 0$.

By Eichler–Shimura $\dim W = 2 \dim S_k + 1 = 3$, so it is enough that the three be independent. Each satisfies $\langle P, P \rangle \neq 0$ [9], and mutually orthogonal vectors with $\langle P, P \rangle \neq 0$ are independent: pairing $\sum_i c_i v_i = 0$ against v_j leaves $c_j \langle v_j, v_j \rangle = 0$.

For orthogonality, note first that since n is even, $X \mapsto -X$ multiplies the coefficient of $X^{n-a} Y^a$ by $(-1)^a$, so an odd polynomial has $p_a = 0$ for even a and an even one has $q_b = 0$ for odd b . As Eq. (43) pairs p_a against q_{n-a} , and a odd forces $n - a$ odd, opposite parities pair to zero: the odd $W_-^{(k)}$ is orthogonal to the even $W_+^{(k)}$ and $v_0^{(k)}$. For the remaining pair, p_0 is supported on its two extreme coefficients, so $\langle P, p_0 \rangle$ is the coefficient of Y^n in P less that of X^n ; c annihilates both of these in $W_+^{(k)}$, whence $\langle W_+^{(k)}, v_0^{(k)} \rangle = 0$. Hence $\{W_+^{(k)}, W_-^{(k)}, v_0^{(k)}\}$ is an orthogonal basis, with $\langle v_0^{(k)}, v_0^{(k)} \rangle = -2$ at every such k . $\square$

Definition 3.14 (Periods, Quasi-Periods). Let k be such that $\dim(S_k) = 1$, $n := k - 2$, $W_\pm^{(k)}(X, Y) := Y^n W_\pm^{(k)}(X/Y)$ be as in Definition 3.12, and $r_f(X, Y)$ be as in Definition 3.7. Now, define

$$\omega^\pm(f) := \frac{\langle r_f, W_\pm^{(k)} \rangle}{\langle W_\pm^{(k)}, W_\pm^{(k)} \rangle}, \quad (47)$$

where $\langle \cdot, \cdot \rangle$ is as in Eq. (43). When f is a cusp form, $\omega^\pm(f)$ are its periods; when f is a weak cusp form, $\omega^\pm(f)$ are its quasi-periods, sometimes denoted $\eta^\pm(f)$.

Now, having defined our periods and quasi-periods, we use them together with our L-functions to extract periods and quasi-periods for $f \in S_k^!$ (when $\dim(S_k) = 1$). To this effect, we recall the notation of [5]. Decomposing the modular weight $k = 12\lambda + k'$ where $k' \in \{0, 4, 6, 8, 10, 14\}$, we see $\lambda = \dim(S_k)$ when $\lambda \geq 0$. For any λ , we may denote $f_{k,m} := 1/q^m + O(q^{\lambda+1}) \in M_k^!$. When $\dim(S_k) = 1$, we have $f_{k,m} = 1/q^m + O(q^2)$. Here, define $\Delta_k := f_{k,-1} = q + O(q^2) \in S_k$ and $\hat{\Delta}_k := f_{k,1} = 1/q + O(q^2) \in S_k^!$. Further, define

$$\text{pd}_k := \begin{pmatrix} \omega^+(\Delta_k) & \omega^-(\Delta_k) \\ \eta^+(\hat{\Delta}_k) & \eta^-(\hat{\Delta}_k) \end{pmatrix}, \quad \alpha_k := \frac{(-1)^{1+k/2} \det(\text{pd}_k)}{(2\pi i)^{k-1} (k-2)!} \quad (48)$$

k	$\omega^+(\Delta_k)$	$\omega^-(\Delta_k)/i$	$\eta^+(\hat{\Delta}_k)$	$\eta^-(\hat{\Delta}_k)/i$	α_k
12	-6.89168e7	-5.58502e6	1.27202e11	1.02767e10	1
16	-8.81481e11	-1.54063e10	4.14120e13	6.83541e11	13/30
18	-5.07833e13	-5.43305e12	-2.19200e14	-3.36339e13	2/3
20	-4.22882e16	-3.73170e14	6.38640e16	-1.89823e14	17/5
22	-1.49974e18	-4.24489e16	1.43795e18	-4.41050e16	19/21
26	-1.18673e22	-3.60961e21	1.18653e22	-3.61016e21	23/15

Table 1: Periods $\omega^\pm(\Delta_k)$ and quasi-periods $\eta^\pm(\hat{\Delta}_k)$, and the rational number α_k . While $\alpha_{12} = 1$ matches Brown [2], α_k for $k \geq 16$ appear to be new, interesting, rational numbers, albeit with implicit dependence on the period-polynomial basis normalizations—in contrast to the normalization-free Petersson check above. We have not found a clear pattern for the α_k . See [9] for the scripts that extracted these figures (to 50-digit precision).

Brown–Hain [7, Prop. 5.6] show that $\det(\text{pd}_k)$ satisfies a Legendre-type relation, $\det(\text{pd}_k) \in \mathbb{Q}^\times \cdot (2\pi i)^{k-1}$. Brown [2, §8.3] numerically finds $\alpha_k = 1$; we reproduce this result at $k = 12$ and give the analogous rational multiples for $k \in \{16, 18, 20, 22, 26\}$ in Table 1, below, via the normalization of Definition 3.12.

4 L-functions, period polynomials and poles at generic locations

For the reconstruction of period polynomials from special values of L^* , which Section 3 carried out for $f \in S_k^!$, we needed a region free of poles. As it happens, and as we show below, when the L-functions defined in this section are used to build period polynomials that are naturally elements of W , choosing the base point $\tau_0 = \rho + 1 = \frac{1}{2} + \frac{\sqrt{3}}{2}i$ and taking the base contour of the reference homotopy class to lie on the arc $|\tau| = 1$ running from ρ to $\rho + 1$, along the lower boundary of the fundamental domain and its S -image, makes that pole-free region of codimension zero in the upper half-plane. Motivated by this, we derive here explicit expressions for the L-function along this reference contour, defined so as to allow similarly explicit representations in terms of the q -series coefficients of generic $f \in F_k$.

For $f \in F_k$ with poles in $\mathbb{H}$, the two segment integrals of Definition 2.3 depend on their contours; we show their sum depends only on the homotopy class of the pair.

Proposition 4.1. *For $f \in F_k$ and $\tau_0 \in \mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, the L-integral $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ depends only on the homotopy class of its two segments in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, and not on the basepoint τ_0 . At fixed τ_0 each of L_S^*, L_T^* is separately path-independent.*

Proof. At fixed τ_0 the integrands $f(\tau)\tau^{s-1}$ and $f(\tau)\tilde{k}_T(\tau, s)$ are holomorphic on $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, so Cauchy’s theorem makes each of L_S^*, L_T^* constant on the homotopy class of its own segment. Under

a motion of the basepoint, $\partial_{\tau_0} L^*$ is evaluated at the moving endpoints alone; the computation of Proposition 2.6, which uses only $f(\tau_0 - 1) = f(\tau_0)$ and $f(-1/\tau_0) = \tau_0^k f(\tau_0)$, cancels the boundary terms of L_S^* against those of L_T^* , the interior poles lying off the contours. Hence $\partial_{\tau_0} L^* = 0$ along any motion in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, while each summand may separately vary. $\square$

Definition 4.2. For $f \in F_k$ and $\tau_0 \in \mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, let $[\gamma_{\tau_0}^S], [\gamma_{\tau_0}^T]$ denote the homotopy classes of the two segments in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$.

Thus, with $\gamma_{\tau_0}^S \in [\gamma_{\tau_0}^S]$ and $\gamma_{\tau_0}^T \in [\gamma_{\tau_0}^T]$, by Proposition 4.1 the value

$$L^*(f, s; [\gamma_{\tau_0}^S], [\gamma_{\tau_0}^T]) := L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) = L_S^*(f, s; \gamma_{\tau_0}^S) + L_T^*(f, s; \gamma_{\tau_0}^T) \quad (49)$$

is independent of the chosen representative paths and independent of τ_0 (subject to natural constraints of τ_0 monodromies around singular points, which move between homotopy classes).

The independence of τ_0 is exact, and the mechanism is the T -shift of the kernel; we record it, since the treatment of a pole at an elliptic point rests on it.

Lemma 4.3. Let $f \in F_k$ and let $\tau_0, \tau'_0 \in \mathbb{H}$ be joined by a path σ meeting no pole of f , with the segments at τ'_0 obtained from those at τ_0 by appending σ and its S - and T^{-1} -images. Then $L^*(f, s; \gamma_{\tau'_0}^S, \gamma_{\tau'_0}^T) = L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ for every $s \in \mathbb{C}$.

Proof. Write Δ for the change in each segment integral. On the S -side, $\Delta L_S^* = -\int_{\sigma} f(\tau) \tau^{s-1} d\tau + \int_{S\sigma} f(\tau) \tau^{s-1} d\tau$, and substituting $\tau \mapsto S\tau$ with $f|_k S = f$ gives $f(S\tau)(S\tau)^{s-1} d(S\tau) = (-1)^{s-1} f(\tau) \tau^{k-s-1} d\tau$, so the second term is $(-1)^{s-1} \int_{\sigma} f(\tau) \tau^{k-s-1} d\tau$. On the T -side, T -periodicity of f gives $\Delta L_T^* = -\int_{\sigma} f(\tau) [\tilde{k}_T(\tau, s) - \tilde{k}_T(\tau - 1, s)] d\tau$, and $\zeta(w, \tau + 1) = \zeta(w, \tau) - \tau^{-w}$ applied to both Hurwitz terms of Definition 2.2 yields

$$\tilde{k}_T(\tau, s) - \tilde{k}_T(\tau - 1, s) = -\tau^{s-1} + e^{i\pi(s-1)} \tau^{k-s-1}. \quad (50)$$

Adding the two changes, the integrand is $-\tau^{s-1} + (-1)^{s-1} \tau^{k-s-1} + \tau^{s-1} - e^{i\pi(s-1)} \tau^{k-s-1}$, which vanishes identically since $(-1)^{s-1} = e^{i\pi(s-1)}$. The prefactor $e^{-\pi i s/2}$ multiplies through. $\square$

The hypothesis is only that σ avoids the poles: crossing one moves between homotopy classes, which is the content of Lemma 4.8 below. In particular L^* is *constant* in any parameter that slides τ_0 without crossing a pole, not merely convergent as that parameter tends to a limit.

We would like to speak of “the” L-integral, and homotopy classes split this into L-integrals for contours in different classes. We therefore fix a reference class, and we fix it on the boundary of the fundamental domain.

Definition 4.4 (Reference homotopy class for $f \in F_k$). Let γ^{arc} be the geodesic segment $\{e^{i\theta} : \frac{\pi}{3} \leq \theta \leq \frac{2\pi}{3}\}$, oriented from ρ to $\rho + 1$, and let $\tau_0 = \rho + 1$. The geodesic reference class is the class of $\hat{\gamma}_f^S = \hat{\gamma}_f^T = \gamma^{\text{arc}}$.

The choice is admissible by Definition 2.1, since $-1/(\rho + 1) = \rho$ and $(\rho + 1) - 1 = \rho$: the two segments share their endpoints and may be taken to be the same path.

It is also the only such choice. The two segments share their endpoints exactly when $S\tau_0 = T^{-1}\tau_0$, that is $-1/\tau_0 = \tau_0 - 1$, and $\tau^2 - \tau + 1$ has the single root $\rho + 1$ in $\mathbb{H}$; the same equation says τ_0 is the fixed point of $U = TS$. The segment is then the geodesic from ρ to $\rho + 1$, which is the wall between the fundamental domain and its S -image, and S carries it to itself. T admits no counterpart: being parabolic it fixes no point of $\mathbb{H}$, and it exchanges the two vertical walls rather than preserving either.

Poles on γ^{arc} never occur singly. As S acts on the unit circle by $e^{i\theta} \mapsto e^{i(\pi-\theta)}$, preserving $\frac{\pi}{3} \leq \theta \leq \frac{2\pi}{3}$, while $f^{-1}(\infty)$ is Γ -stable and $f|_k S = f$, a pole of order P at $e^{i\phi}$ on the arc is accompanied by one of the same order at $e^{i(\pi-\phi)}$, again on the arc. These S -pairs have two degenerations: the pair collapses to a single point exactly when $\phi = \frac{\pi}{2}$, and it consists of the two endpoints exactly when $\phi \in \{\frac{\pi}{3}, \frac{2\pi}{3}\}$, since $S(\rho+1) = -1/(\rho+1) = \rho$. All three cases are treated the same way, by moving the poles off the contour rather than moving the contour.

We move the contour off the poles, never the poles off the contour. Displacing a pole is not available to us: $f^{-1}(\infty)$ is $\text{SL}_2(\mathbb{Z})$ -stable, so a pole cannot be moved alone, and at an elliptic point of order n the stabiliser acts as rotation by $2\pi/n$ about a point the contour runs through, sending any displacement to the far side and pinching the contour. Displacing the base point costs nothing, by Lemma 4.3.

Definition 4.5. For $0 < \epsilon < \frac{1}{2}$ put $\tau_0(\epsilon) := (1 + \epsilon)e^{i\pi/3}$, so that $S\tau_0(\epsilon) = (1 + \epsilon)^{-1}e^{2i\pi/3}$ and $T^{-1}\tau_0(\epsilon) = \rho + \epsilon e^{i\pi/3}$. Let $\hat{\gamma}_\epsilon^S$ be the logarithmic spiral

$$\tau(\theta) = \exp\left(\log(1 + \epsilon) \frac{\pi/2 - \theta}{\pi/6} + i\theta\right), \quad \theta : \frac{2\pi}{3} \rightarrow \frac{\pi}{3}, \quad (51)$$

and let $\hat{\gamma}_\epsilon^T$ be a path from $T^{-1}\tau_0(\epsilon)$ to $S\tau_0(\epsilon)$ inside $|\tau - \rho| = O(\epsilon)$, followed by $\hat{\gamma}_\epsilon^S$. The ϵ -shifted geodesic class is the pair of their homotopy classes in $\mathbb{H} \setminus f^{-1}(\infty)$.

Three features of (51) are used below. Its $\log r$ is odd about $\theta = \frac{\pi}{2}$ and S acts by $(r, \theta) \mapsto (1/r, \pi - \theta)$, so $\hat{\gamma}_\epsilon^S$ is S -symmetric and Lemma 4.29 applies to the S^2 -path. It lies outside $|\tau| = 1$ for $\theta < \frac{\pi}{2}$ and inside for $\theta > \frac{\pi}{2}$, so an S -pair of poles on the arc is passed on opposite sides automatically, with no side chosen by hand. And it reduces to γ^{arc} as $\epsilon \rightarrow 0$, the two segments then sharing their endpoints as in Definition 4.4.

Since $S\tau_0(\epsilon)$ and $T^{-1}\tau_0(\epsilon)$ are distinct for $\epsilon \neq 0$, the connector in Definition 4.5 is not determined by its endpoints, and the classes available to it are indexed by an integer. We name that integer through the relator path, where it is what the period polynomial sees.

Definition 4.6. Let $W(f, \epsilon) \in \mathbb{Z}$ be the winding number about $\rho + 1$ of the closed path traced by $(ST)^3$ in the ϵ -shifted geodesic class, computed in the coordinate $w = (\tau - \rho - 1)/(\tau - \overline{\rho + 1})$.

By Lemma 4.8 the L-integral is an affine function of W , and by Lemma 4.36 below $W \equiv -1 \pmod{3}$, so $W = 0$ is not realised by any single contour. It is nonetheless the value at which the period polynomial lies in W , and L^* is linear, so we define the L-integral by evaluating that affine function there.

Definition 4.7. Let $f \in F_k$ have a pole on $\text{SL}_2(\mathbb{Z}) \cdot \rho$ or at i . Then $L^*(f, s)$ is the value at $W = 0$ of the affine function $W \mapsto L^*(f, s; \hat{\gamma}_\epsilon^S, \hat{\gamma}_\epsilon^T)$, and likewise for the period polynomial of Definition 3.7.

Equivalently, $L^*(f, s)$ is the weighted average of L-integrals over two realisable classes, with weights w_j fixed by $\sum_j w_j = 1$ and $\sum_j w_j W_j = 0$: at i , where W is odd, the two classes $W = \pm 1$ enter with weight $\frac{1}{2}$ each, so L^* is the mean of the two indentations; at ρ the classes $W = -1$ and $W = 2$ enter with weights $\frac{2}{3}, \frac{1}{3}$. In both cases the correction to the non-winding class is $\frac{1}{n}$ of one residue,

$$L^*(f, s) = L^*(f, s; W = -1) + \frac{2\pi i}{n} e^{-\pi i s/2} \text{Res}_{\tau=\tau_e}(f \tilde{k}_T), \quad (52)$$

the weight $\frac{1}{n}$ depending only on the order of the elliptic point, never on f or on the order of its pole. So (52) is linear in f , and is independent of ϵ by Lemma 4.3. Poles of f on γ^{arc} away from

the elliptic points need no prescription at all: the spiral of (51) already separates each S -pair, and by Lemma 4.33 either separation puts $\hat{r}_f$ in W .

With this definition we may unambiguously write

$$L^*(f, s) := L^*(f, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]) , \quad (53)$$

where for f with a pole at an elliptic point the right-hand side is read through Definition 4.7: there the value is not an integral over one homotopy class but the weighted average of two, so (53) fixes a number and not a contour. Having fixed this reference value, consider two paths $\gamma_{\tau_0}^S$ and $\gamma_{\tau_0}^T$ which may lie in any homotopy class, and let $X_S(p), X_T(p) \in \mathbb{Z}$ count the windings of $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$ about each pole $\tau_p \in f^{-1}(\infty)$ relative to the reference class.

Lemma 4.8. *Let $f \in F_k$, $r_S(f, \tau_p) := \text{Res}_{\tau=\tau_p}(f(\tau)\tau^{s-1})$ and $r_T(f, \tau_p) := \text{Res}_{\tau=\tau_p}(f(\tau)\tilde{k}_T(\tau, s))$. Then,*

$$L^*(f, s; [\gamma_{\tau_0}^S], [\gamma_{\tau_0}^T]) = L^*(f, s) + 2\pi i e^{-\pi i s/2} \sum_{\tau_p \in f^{-1}(\infty)} \left[X_S(\tau_p) r_S(f, \tau_p) + X_T(\tau_p) r_T(f, \tau_p) \right]. \quad (54)$$

Proof. Two pairs whose winding data differ by one unit at a single pole p , on one segment, bound a loop encircling p once; the residue theorem adds $2\pi i$ times the residue of that segment's integrand, $f(\tau)\tau^{s-1}$ for γ^S or $f(\tau)\tilde{k}_T(\tau, s)$ for γ^T , and the prefactor $e^{-\pi i s/2}$ of Definition 2.3 multiplies through. Summing over crossings gives (54). $\square$

In particular two contours passing on opposite sides of a pole differ by its residue, and nothing else. For a pole on the arc away from the elliptic points this is the whole of the ambiguity.

Proposition 4.9. *Let $f \in F_k$ have poles on γ^{arc} , none of them at i , ρ or $\rho + 1$. Write $\gamma_{\pm}^{\text{arc}}(p)$ for γ^{arc} indented near p so as to pass outside, respectively inside, $|\tau| = 1$. Then $L^*(f, s)$ is defined on either contour, and replacing $\gamma_+^{\text{arc}}(p)$ by $\gamma_-^{\text{arc}}(p)$ at one pole changes it by*

$$-2\pi i e^{-\pi i s/2} [r_S(f, p) + r_T(f, p)]. \quad (55)$$

The spiral $\hat{\gamma}_\epsilon^S$ of (51) realises one such contour for every $\epsilon > 0$, passing outside at $\theta < \frac{\pi}{2}$ and inside at $\theta > \frac{\pi}{2}$, so each S -pair is separated without a side being chosen by hand.

Proof. Each indented contour keeps p at positive distance, so both segment integrals converge and L^* is defined on either; the spiral does the same for every $\epsilon > 0$, and the sides it takes are as stated because $\log r$ in (51) is positive for $\theta < \frac{\pi}{2}$ and negative for $\theta > \frac{\pi}{2}$.

For the difference, $\gamma_+^{\text{arc}}(p)$ and $\gamma_-^{\text{arc}}(p)$ agree off a neighbourhood of p and together bound a loop encircling p once. Along γ^{arc} the direction of travel is θ decreasing, so it is $-ie^{i\theta}$ up to a positive factor, while the outward radial direction is $e^{i\theta} = i \cdot (-ie^{i\theta})$: outward lies to the left of travel. The sliver between the two indentations therefore lies to the right of $\gamma_+^{\text{arc}}(p)$ traversed forward, and the loop encircles p negatively. This is the winding datum $X_S(p) = X_T(p) = -1$ of Lemma 4.8, which gives (55). $\square$

The exclusion of i , ρ and $\rho + 1$ is not a technicality: there the stabiliser orbit meets γ^{arc} on both sides at once, no single-sided indentation exists, and the winding of Definition 4.6 cannot be set to zero by any one contour.

This provides the basics for the finite L-functions of $f \in F_k$ in this class. We wish, however, to present them in terms of their q -series coefficients, as was done for $f \in M_k^1$ in Section 3. The obstruction to doing so directly is not a technical one. For $f \in M_k^1$ the series $\sum_n c_f(n)q^n$ converges

on all of $\mathbb{H}$, as f is holomorphic there, and term-by-term evaluation is legitimate throughout. A pole at finite height ends this. The expansion about the cusp then represents f only above the highest pole, on $\text{Im } \tau > y_f := \sup\{\text{Im } \tau_p\}$, with coefficients growing as $|c_f(n)| \gtrsim n^{m-1} e^{2\pi n \text{Im } \tau_p}$ for a pole of order m [3]. The arc has height varying between $\frac{\sqrt{3}}{2}$ and 1, so the operative threshold is its lowest point $\frac{\sqrt{3}}{2}$: as soon as f has a pole at or above that height the q -series diverges somewhere on the contour, and there is then no series to integrate term by term, rather than a slowly convergent one to be handled with care. To do this, we must introduce the polylogarithmic projections of Ref. [3].

4.1 Explicit formulae for $L^*(f, s)$ on the arc

We lay out the starting definitions and facts.

Definition 4.10. $F(\infty)$ is the space of meromorphic functions $g : \mathbb{H} \rightarrow \mathbb{C}$ with $g(\tau + 1) = g(\tau)$, finitely many poles in each compact subset of the strip $|\text{Re } \tau| \leq \frac{1}{2}$, and bounded pole order at $q = 0$.

Definition 4.11. Let $f \in F_k$ have a pole of order m_p at $\tau_p \in \mathbb{H}$. On a punctured ε -ball about τ_p small enough to exclude the other poles of f , we have

$$f(\tau) = \sum_{m=1}^{m_p} \frac{r_{f,\tau_p}^*(m)}{(\tau - \tau_p)^m} + g(\tau), \quad r_{f,\tau_p}^*(m) = \frac{1}{2\pi i} \oint_{|\tau - \tau_p|=\varepsilon} f(\tau) (\tau - \tau_p)^{m-1} d\tau, \quad (56)$$

with g holomorphic at τ_p . We call $r_{f,\tau_p}^*(1), \dots, r_{f,\tau_p}^*(m_p)$ the pole weights of f at τ_p ; the first is the residue, and the last is nonzero, which is what fixes the order. The polylog projection of f at τ_p is

$$\hat{P}_{\tau_p} f(\tau) := \sum_{m=1}^{m_p} r_{f,\tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)}), \quad \text{Li}_s(x) := \sum_{n \geq 1} \frac{x^n}{n^s}. \quad (57)$$

The composite projector is the one place where the contour enters. Its threshold is set so that $(1 - \hat{P})f$ admits a q -series everywhere it is used, which is a statement about a half-plane: a q -series converges exactly on $\text{Im } \tau > h$ with h the height of the highest surviving pole. The operative height is the lowest point of γ^{arc} , namely $\frac{\sqrt{3}}{2}$, since the mode sums below are controlled by $e^{-2\pi n \text{Im } \tau}$ and the arc descends to that height at its endpoints.

Definition 4.12. For $f \in F_k$ with T -inequivalent finite-order poles $\{\tau_p\}$ of orders $\{m_p\}$, set

$$\hat{P}^{\text{geo}} := \sum_{\text{Im } \tau_p \geq \frac{\sqrt{3}}{2}} \hat{P}_{\tau_p}, \quad (58)$$

with $\hat{P}_{\tau_p}$ as in Definition 4.11, and write $f = \hat{P}^{\text{geo}} f + (I - \hat{P}^{\text{geo}})f$.

Lemma 4.13. The normalized block $\frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)})$ has principal part exactly $(\tau - \tau_p)^{-m}$, so $\hat{P}_{\tau_p} f$ matches the principal part of f and $(I - \hat{P}_{\tau_p})f$ is holomorphic at τ_p .

Proof. By the small- t expansion $\text{Li}_{1-m}(e^{-t}) = (m-1)! t^{-m} + O(t^0)$ with $t = -2\pi i(\tau - \tau_p)$, the block has a pure pole $(\tau - \tau_p)^{-m} + O((\tau - \tau_p)^0)$ at τ_p , with no lower-order poles. Summing against the Laurent coefficients $r_{f,\tau_p}^*(m)$ reproduces the principal part of f term by term, leaving $(I - \hat{P}_{\tau_p})f$ regular at τ_p . $\square$

Definition 4.14. For $f \in F_k$ and $\hat{P}^{\text{geo}}$ as in Definition 4.12, write

$$(I - \hat{P}^{\text{geo}})f(\tau) = \sum_n c_{Rf}(n) q^n, \quad q = e^{2\pi i \tau}. \quad (59)$$

Lemma 4.15. $F_k \subset F(\infty)$ and $(I - \hat{P}^{\text{geo}})f \in F(\infty)$. Every pole of $(I - \hat{P}^{\text{geo}})f$ has $\text{Im } \tau < \frac{\sqrt{3}}{2}$, with a maximal height $y_b < \frac{\sqrt{3}}{2}$ (set $y_b = 0$ if $(I - \hat{P}^{\text{geo}})f$ is holomorphic).

Proof. A weakly holomorphic form is T -periodic, meromorphic, and $O(e^{C\text{Im}\tau})$ at the cusp, so $F_k \subset F(\infty)$; the projected polylogs $\text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)})$ share these properties, so $(I - \hat{P}^{\text{geo}})f \in F(\infty)$ (and leaves F_k , being non-modular). By Lemma 4.13, $\hat{P}^{\text{geo}}$ cancels the principal part of every pole with $\text{Im } \tau \geq \frac{\sqrt{3}}{2}$, so the poles of $(I - \hat{P}^{\text{geo}})f$ lie strictly below $\frac{\sqrt{3}}{2}$; finite in each compact set and accumulating only at $\text{Im } \tau = 0$, they have a maximal height $y_b < \frac{\sqrt{3}}{2}$. $\square$

The Fourier coefficients of a T -periodic meromorphic function are controlled by its highest pole: $|c_{Rf}(n)| \leq e^{2\pi n(y_b + \varepsilon)}$ for every $\varepsilon > 0$ and all large n [3]. Since $y_b < \frac{\sqrt{3}}{2}$, the series $\sum_n c_{Rf}(n) q^n$ converges geometrically down to $\text{Im } \tau = \frac{\sqrt{3}}{2}$, at base $e^{-2\pi(\frac{\sqrt{3}}{2} - y_b - \varepsilon)}$, and so on γ^{arc} itself. This is the point of the projection: convergence merely for large $\text{Im } \tau$ would be automatic and of no use, as the contour descends to height $\frac{\sqrt{3}}{2}$.

One further integral of f along the contour will be needed, both for the residue of L^* at $s = 0$ and, later, for period polynomials.

Definition 4.16. For $f \in F_k$, set $\Phi(f) := \int_{\hat{\gamma}_f^T} f(\tau) d\tau$, the integral of f along the T -segment of the contour defining $L^*(f, s)$.

Φ is not an independent choice: it is what the parabolicity condition $\hat{C}_T = 0$ measures, and $\hat{C}_T$ is assembled from the T -transport, so Φ follows the contour. Deforming the segments replaces γ^{arc} in the integral by whatever $\hat{\gamma}_f^T$ then is, with no further stipulation.

Since $\hat{\gamma}_f^T$ runs from $T^{-1}\tau_0$ to τ_0 , and $e^{2\pi i n \tau}$ takes equal values at those two endpoints, the modes $n \neq 0$ of the q -series of f cancel between them. Hence $\Phi(f) = c_f(0)$ whenever that series converges along the segment, which is to say whenever no pole of f lies above $\hat{\gamma}_f^T$; a pole below is harmless. Otherwise $\Phi(f)$ is $c_f(0)$ less $2\pi i$ times the residues of the poles in the strip between the segment and the cusp. In particular $\Phi(E_k) = 1$, and $\Phi(f) = 0$ for $f \in S_k$.

The relations satisfied by the pole weights at an elliptic point are contour-free, so they hold here verbatim; we record them now, as both the block integral below and the period polynomials constructed later use them.

Lemma 4.17. Let τ_e be i or ρ , of order $n = 2$ or $n = 3$, let $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be the generator S or ST of its stabiliser in $\text{SL}_2(\mathbb{Z})$, and set $\zeta := c\tau_e + d$, so that $c = 1$ and $\zeta = e^{i\pi/n}$. Let $f \in F_k$ have a pole of order P at τ_e with pole weights $r_{f, \tau_e}^*(m)$. Modularity is equivalent to one relation for each principal-part order μ ,

$$\sum_{m=\mu}^P r_{f, \tau_e}^*(m) \binom{m}{\mu} \zeta^m = \zeta^k \sum_{m=\mu}^P r_{f, \tau_e}^*(m) \binom{k}{m-\mu} \zeta^{-m}, \quad \forall \mu = 1, \dots, P, \quad (60)$$

and the leading relation is $(\zeta^{2P} - \zeta^k) r_{f, \tau_e}^*(P) = 0$, so that $2P \equiv k \pmod{2n}$.

Proof. Write $\tau = \tau_e + \varepsilon$. Since $\det \gamma = 1$ and $\gamma\tau_e = \tau_e$,

$$\gamma\tau - \tau_e = \frac{\tau - \tau_e}{(c\tau + d)(c\tau_e + d)} = \frac{\varepsilon}{\zeta(\zeta + \varepsilon)}, \quad (c\tau + d)^k = \zeta^k(1 + \varepsilon/\zeta)^k, \quad (61)$$

using $c = 1$. Hence $(\gamma\tau - \tau_e)^{-m} = \zeta^{2m}\varepsilon^{-m}(1 + \varepsilon/\zeta)^m$, and matching the coefficient of $\varepsilon^{-\mu}$ in $f(\gamma\tau) = (c\tau + d)^k f(\tau)$ through $(1 + \varepsilon/\zeta)^a = \sum_l \binom{a}{l} \zeta^{-l} \varepsilon^l$ gives $\sum_{m \geq \mu} r^*(m) \zeta^{2m} \binom{m}{\mu} \zeta^{-(m-\mu)} = \zeta^k \sum_{m \geq \mu} r^*(m) \binom{k}{m-\mu} \zeta^{-(m-\mu)}$; dividing by ζ^μ yields (60). At $\mu = P$ only $m = P$ survives on each side. $\square$

At $n = 2$ the order condition forces P odd when $k \equiv 2 \pmod{4}$ and P even when $4 \mid k$; at $n = 3$ it reads $2P \equiv k \pmod{6}$.

Each pole with $\text{Im } \tau_p \geq \frac{\sqrt{3}}{2}$ contributes one projected polylog to $L^*(f, s)$, whose L-integral we now record.

Definition 4.18. For a pole of order $N + 1$ projected at z , let $\lambda_{N,z}$ denote the corresponding block of (57), and set $I_N^{\text{geo}}(s, z) := L^*(\lambda_{N,z}, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T])$, evaluated in the geodesic reference class of Definition 4.4.

The blocks need not be integrated on the arc. Both kernels τ^{s-1} and $\tilde{k}_T(\tau, s)$ are holomorphic on all of $\mathbb{H}$, so the only obstruction to deforming is the block's own poles at $z + \mathbb{Z}$, and one may evaluate on the horizontal chord from ρ to $\rho + 1$ instead. Let Λ be the open lens between the two, bounded above by γ^{arc} and below by the chord, so that $\Lambda \subset \{|\tau| < 1, \text{Im } \tau > \frac{\sqrt{3}}{2}, |\text{Re } \tau| < \frac{1}{2}\}$; at most one member of $z + \mathbb{Z}$ lies in it. If z itself lies on γ^{arc} the contour is indented past it as in Proposition 4.9. In either situation set $\varepsilon_z := 1$ when z lies between the contour and the chord, that is when $z \in \Lambda$, or when $z \in \gamma^{\text{arc}}$ and the indentation is the outward one $\gamma_+^{\text{arc}}(z)$; and $\varepsilon_z := 0$ otherwise. The deformation to the chord costs one residue exactly when $\varepsilon_z = 1$.

Lemma 4.19. Let $\text{Im } z > \frac{\sqrt{3}}{2}$, set $\tau_0 = \rho + 1$, let $\gamma^*(s, x) := \sum_{\ell \geq 0} \frac{(-x)^\ell}{\ell! (\ell + s)}$, entire in x , read empty products below as 1, and set $(2\pi n)^w := e^{w(\log 2\pi |n| + i\pi)}$ for $n < 0$. With, for $n \in \mathbb{Z} \setminus \{0\}$,

$$G_{s,k}^{\text{geo}}(n) := e^{-\pi i s/2} \left[\tau_0^s \gamma^*(s, -2\pi i n \tau_0) - \rho^s \gamma^*(s, -2\pi i n \rho) \right] + \frac{\Gamma(s, -2\pi i n \tau_0)}{(2\pi n)^s} + i^k \frac{\Gamma(k - s, -2\pi i n \tau_0)}{(2\pi n)^{k-s}}, \quad (62)$$

$$G_{s,k}^{\text{geo}}(0) := e^{-\pi i s/2} \frac{\tau_0^s - \rho^s}{s} - \frac{(\tau_0/i)^s}{s} - i^k \frac{(\tau_0/i)^{k-s}}{k-s}, \quad (63)$$

$$X_N(s, z) := \frac{e^{-\pi i s/2}}{(2\pi i)^N} \left[(-1)^N z^{s-1-N} \prod_{j=0}^{N-1} (s-1-j) + \zeta(1-s+N, z+1) \prod_{j=0}^{N-1} (1-s+j) - e^{i\pi(s-1)} \zeta(1-k+s+N, z+1) \prod_{j=0}^{N-1} (1-k+s+j) \right], \quad (64)$$

we have

$$I_N^{\text{geo}}(s, z) = -\delta_{N,0} G_{s,k}^{\text{geo}}(0) + (-1)^{N+1} \sum_{n \geq 1} n^N e^{2\pi i n z} G_{s,k}^{\text{geo}}(-n) + \varepsilon_z X_N(s, z), \quad (65)$$

the sum converging at rate $e^{-2\pi(\text{Im } z - \frac{\sqrt{3}}{2})}$.

Proof. Both kernels are holomorphic on $\mathbb{H}$, and the only poles of $\lambda_{N,z}$ are at $z + \mathbb{Z}$, so Cauchy's theorem moves both segment integrals to the chord at the cost of whatever residues lie between the two paths; by the paragraph above that is $\{z\}$ when $\varepsilon_z = 1$ and nothing otherwise. The chord is not a convenience: along γ^{arc} itself $\text{Im } \tau$ ranges over $[\frac{\sqrt{3}}{2}, 1]$, so for $\frac{\sqrt{3}}{2} < \text{Im } z < 1$ the comparison below would change sign partway along, whereas on the chord $\text{Im } \tau \equiv \frac{\sqrt{3}}{2}$ and it is a single global inequality.

On the chord $|e^{2\pi i(\tau-z)}| > 1$, so the defining series of Li_{-N} diverges and inversion $\text{Li}_{-N}(w) = (-1)^{N+1} \text{Li}_{-N}(1/w) - \delta_{N,0}$ rewrites $\lambda_{N,z}$ as $(-1)^{N+1} \sum_{n \geq 1} n^N e^{2\pi i n z} q^{-n} - \delta_{N,0}$, uniformly convergent there since $\text{Im } z > \frac{\sqrt{3}}{2}$. Integrating term by term, $G_{s,k}^{\text{geo}}(-n) = L^*(q^{-n}, s)$ in this class: the last two terms of (62) are L_T^* by Lemma 3.3 at τ_0 , and the first is $L_S^* = e^{-\pi i s/2} \int_{\rho}^{\tau_0} e^{-2\pi i n \tau} \tau^{s-1} d\tau$, which on expanding $e^{-2\pi i n \tau}$ and integrating $\tau^{\ell+s-1}$ term by term gives $e^{-\pi i s/2} \sum_{\ell \geq 0} \frac{(-2\pi i n)^\ell}{\ell! (\ell+s)} [\tau_0^{\ell+s} - \rho^{\ell+s}]$, the first bracket of (62). Likewise $G_{s,k}^{\text{geo}}(0) = L^*(1, s)$. The rate follows from $|e^{2\pi i n z}| = e^{-2\pi n \text{Im } z}$ against $|G_{s,k}^{\text{geo}}(-n)| \sim e^{\pi \sqrt{3} n}$.

For the crossing term put $u = \tau - z$ and $v = 2\pi i u + t$. Then $\sum_{N \geq 0} \text{Li}_{-N}(e^{2\pi i u}) \frac{t^N}{N!} = \frac{e^v}{1-e^v} = -\sum_{m \geq 0} \frac{B_m(-v)^m}{m! v}$, of which only the $m = 0$ term $-1/v$ carries negative powers of u ; expanding $-1/(2\pi i u + t) = -\sum_{r \geq 0} (-1)^r t^r (2\pi i u)^{-r-1}$ and reading off $t^N/N!$ gives

$$\lambda_{N,z}(\tau) = \frac{(-1)^{N+1} N!}{(2\pi i(\tau - z))^{N+1}} + O(1), \quad (66)$$

a lone singular term, so $\text{Res}_{\tau=z}(\lambda_{N,z} \kappa) = (-1)^{N+1} \kappa^{(N)}(z)/(2\pi i)^{N+1}$ for any κ holomorphic at z . Both paths run $\rho \rightarrow \rho + 1$ with the arc above, so together they bound Λ clockwise and the deformation contributes $-2\pi i$ times that residue for each kernel. With $\partial_z^N z^{s-1} = z^{s-1-N} \prod_{j=0}^{N-1} (s-1-j)$ and $\partial_z^N \zeta(w, z+1) = (-1)^N \zeta(w+N, z+1) \prod_{j=0}^{N-1} (w+j)$ applied to both terms of k_T , this is $X_N(s, z)$ of (64). $\square$

Corollary 3.5 is deliberately not used here: it folds the S -segment about i using modularity, and q^n is only T -periodic, so its S -integral must be taken directly. Nor does that segment drop out, as it would for a base point at which the two segments coincide: on the arc it contributes the first bracket of (62), and the threshold governing the convergence rate stands at $\frac{\sqrt{3}}{2}$, in step with Definition 4.12.

The S -term is written with γ^* because that form is branch-free: γ^* is single-valued, so the only multivalued factors are the explicit τ_0^s and ρ^s , and those are unambiguous since $\arg \tau \in [\frac{\pi}{3}, \frac{2\pi}{3}]$ along the chord. It may equally be written with upper incomplete gammas, provided the sheet relation is carried. Under $u = 2\pi i n \tau$ the chord's image runs from $\arg u = \frac{7\pi}{6}$ at ρ to $\frac{5\pi}{6}$ at τ_0 , so the ρ endpoint sits one turn off principal; since $\gamma(s, x) = x^s \gamma^*(s, x)$ with γ^* single-valued, a shift of $\arg x$ by 2π gives

$$\Gamma^{\text{cont}}(s, x) = e^{2\pi i s} \Gamma(s, x) + (1 - e^{2\pi i s}) \Gamma(s), \quad (67)$$

and hence

$$S_n(s) = (2\pi i n)^{-s} \left[e^{2\pi i s} \Gamma(s, 2\pi i n \rho) + (1 - e^{2\pi i s}) \Gamma(s) - \Gamma(s, 2\pi i n \tau_0) \right]. \quad (68)$$

Both the sheet factor and the additive constant are trivial at integer s , which is all that Definition 3.7 requires. The T -term needs neither, since by Lemma 3.3 only τ_0 appears there, at $\arg(2\pi i n \tau_0) = \frac{5\pi}{6}$, the ρ endpoint having been absorbed by the periodicity of $\tilde{k}_T$. It does need the branch of $(2\pi n)^w$ at

$n < 0$ fixed above, which is the one making $\arg(2\pi n) + \arg(\tau_0/i) = \frac{5\pi}{6}$; the opposite choice differs by $e^{2\pi i s}$, and so is invisible at integer s and wrong elsewhere. In practice the γ^* series loses about $2.7n$ digits to cancellation, so (68) is preferable if large n is needed.

On the threshold line itself the roles of the two series are exchanged. A pole at $\text{Im } z = \frac{\sqrt{3}}{2}$ lies below the arc, which at abscissa $\text{Re } z$ has height $\sqrt{1 - (\text{Re } z)^2} > \frac{\sqrt{3}}{2}$, so it is the defining series of Li_{-N} that applies rather than its inversion. That series converges absolutely for $\text{Im } z < \frac{\sqrt{3}}{2}$ and is exactly marginal at $\frac{\sqrt{3}}{2}$, since the arc attains that height at its two endpoints. Write $(a)_{(\ell)} := a(a-1)\cdots(a-\ell+1)$ for the falling factorial, with $(a)_{(0)} := 1$.

Lemma 4.20. *Let $\text{Im } z = \frac{\sqrt{3}}{2}$, and either $z \not\equiv \rho \pmod{\mathbb{Z}}$ or $c_N(s, k) = 0$, and set*

$$c_\ell(s, k) := (s-1)_{(\ell)} \left[-i i^\ell e^{-\pi i s/2} (\tau_0^{s-1-\ell} - \rho^{s-1-\ell}) + (-i\tau_0)^{s-1-\ell} \right] + i^k (k-s-1)_{(\ell)} (-i\tau_0)^{k-s-1-\ell}. \quad (69)$$

Then for every $L \geq N$,

$$I_N^{\text{geo}}(s, z) = \sum_{\ell=0}^L \frac{c_\ell(s, k)}{(2\pi)^{\ell+1}} \text{Li}_{\ell+1-N}(e^{2\pi i(\tau_0-z)}) + R_{N,L}^{\text{geo}}(s, z), \quad (70)$$

where

$$R_{N,L}^{\text{geo}}(s, z) := \sum_{n \geq 1} n^N e^{-2\pi i n z} \left[G_{s,k}^{\text{geo}}(n) - e^{2\pi i n \rho} \sum_{\ell=0}^L \frac{c_\ell(s, k)}{(2\pi n)^{\ell+1}} \right] \quad (71)$$

converges absolutely, its terms being $O(n^{N-L-2})$.

Proof. For $\text{Im } z < \frac{\sqrt{3}}{2}$ the series $\lambda_{N,z} = \sum_{n \geq 1} n^N e^{-2\pi i n z} q^n$ converges uniformly on γ^{arc} , both kernels are bounded there, and term-by-term integration gives $I_N^{\text{geo}}(s, z) = \sum_{n \geq 1} n^N e^{-2\pi i n z} G_{s,k}^{\text{geo}}(n)$, absolutely convergent.

Repeated integration by parts of the S -integral gives $\int_\rho^{\tau_0} e^{2\pi i n \tau} \tau^{s-1} d\tau \sim \sum_{\ell \geq 0} \frac{(-1)^\ell (s-1)_{(\ell)}}{(2\pi i n)^{\ell+1}} [e^{2\pi i n \tau} \tau^{s-1-\ell}]_\rho^{\tau_0}$, while $\Gamma(a, X) \sim X^{a-1} e^{-X} \sum_{\ell \geq 0} (a-1)_{(\ell)} X^{-\ell}$ at $X = -2\pi i n \tau_0 = 2\pi n(-i\tau_0)$ handles the two incomplete gammas. Since $e^{2\pi i n \tau_0} = e^{2\pi i n \rho}$ by T -periodicity, both endpoints carry the same exponential and, using $(-1)^\ell / i^{\ell+1} = -i i^\ell$, $G_{s,k}^{\text{geo}}(n) = e^{2\pi i n \rho} [\sum_{\ell=0}^L c_\ell(s, k) (2\pi n)^{-\ell-1} + O(n^{-L-2})]$ with c_ℓ as in (69).

Splitting the sum along this expansion, the peeled part is $\sum_{\ell=0}^L \frac{c_\ell}{(2\pi)^{\ell+1}} \sum_{n \geq 1} n^{N-\ell-1} e^{2\pi i n(\rho-z)}$, and $e^{2\pi i(\rho-z)} = e^{2\pi i(\tau_0-z)}$, giving the polylogarithms of (70); the remainder is (71), whose terms are $O(n^{N-L-2})$ because $|e^{-2\pi i n z} e^{2\pi i n \rho}| = 1$ on the threshold line. Both sides of (70) are holomorphic in z wherever z avoids $\gamma^{\text{arc}} + \mathbb{Z}$ and $e^{2\pi i(\tau_0-z)} \neq 1$ — the left because the contour integral of Definition 4.18 then has no singular integrand, the right because $R_{N,L}^{\text{geo}}$ converges locally uniformly — so the identity, proved on $\text{Im } z < \frac{\sqrt{3}}{2}$, persists onto the threshold line away from $\rho + \mathbb{Z}$.

At $z \equiv \rho$ the argument $e^{2\pi i(\tau_0-z)}$ equals 1 and the $\ell = N$ term carries $\text{Li}_1(1)$, so the statement needs $c_N = 0$. That coefficient is exactly the endpoint mismatch: by the computation above, c_0 is a non-zero multiple of

$$[\rho^{s-1} + \tilde{k}_T(\rho, s)] - [\tau_0^{s-1} + \tilde{k}_T(\tau_0, s)], \quad (72)$$

which by the shift identity (50) reads $\rho^{s-1} = e^{i\pi(s-1)} \tau_0^{k-s-1}$ and so, with $\rho = e^{2\pi i/3}$ and $\tau_0 = e^{i\pi/3}$, vanishes for every s precisely when $k \equiv 2 \pmod{6}$. This is not a side condition to be hoped for. A pole at ρ has $P \equiv 1 \pmod{n}$ by Lemma 4.17, and at $P = 1$ — the only order for which the block

itself, rather than the winding prescription of Definition 4.7, is what one needs — that forces $k \equiv 2 \pmod{6}$ and so $c_0 = 0$. The divergence one might have expected at ρ is absent exactly when the theory calls on it. $\square$

The blocks assemble into the projection at a pole by linearity alone.

Lemma 4.21. *Let $f \in F_k$ have a pole of order m_p at τ_p with $\text{Im } \tau_p > \frac{\sqrt{3}}{2}$, and pole weights $r_{f,\tau_p}^*(m)$ as in Definition 4.11. Then*

$$L^*(\widehat{P}_{\tau_p} f, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]) = \sum_{m=1}^{m_p} r_{f,\tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} I_{m-1}^{\text{geo}}(s, \tau_p). \quad (73)$$

Proof. By (57) the projection $\widehat{P}_{\tau_p} f$ is the finite combination $\sum_m r_{f,\tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} \lambda_{m-1,\tau_p}$ of the blocks of Definition 4.18. Both segment integrals of Definition 2.3 are linear in the integrand and the combination is finite, so L^* distributes over it. $\square$

What survives every such projection carries a q -series on the whole contour, and may be integrated mode by mode.

Lemma 4.22. *Let $f \in F_k$, write $Rf := (I - \widehat{P}^{\text{geo}})f$, and let $c_{Rf}(n)$ be its Fourier coefficients as in Definition 4.14. Then*

$$L^*(Rf, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]) = c_{Rf}(0) G_{s,k}^{\text{geo}}(0) + \sum_{n \neq 0} c_{Rf}(n) G_{s,k}^{\text{geo}}(n), \quad (74)$$

the sum converging absolutely.

Proof. By Lemma 4.15 every pole of Rf has height at most $y_b < \frac{\sqrt{3}}{2}$, so for any $\varepsilon > 0$ with $y_b + \varepsilon < \frac{\sqrt{3}}{2}$ one has $|c_{Rf}(n)| \leq e^{2\pi n(y_b + \varepsilon)}$ for all large n , while only finitely many $n < 0$ occur, Rf being meromorphic at the cusp. On γ^{arc} one has $\text{Im } \tau \geq \frac{\sqrt{3}}{2}$, so those terms are bounded by $e^{-2\pi n(\frac{\sqrt{3}}{2} - y_b - \varepsilon)}$ and $\sum_n c_{Rf}(n) q^n$ converges uniformly on the contour. Both kernels are continuous, hence bounded, on that compact set, so each segment integral may be taken term by term. The mode q^n with $n \neq 0$ contributes $L^*(q^n, s) = G_{s,k}^{\text{geo}}(n)$, by the computation in the proof of Lemma 4.19 read at general n , and $q^0 = 1$ contributes $G_{s,k}^{\text{geo}}(0)$ of (63). $\square$

Before assembling, we record the functional equation. It holds for every $f \in F_k$, not only for the weakly holomorphic ones, and for the same reason the arc was chosen: γ^{arc} is S -stable. The T -segment needs no property of the path at all.

Lemma 4.23. *Let k be even and $f \in F_k$. In the geodesic reference class,*

$$L^*(f, s) = i^k L^*(f, k - s) \quad (75)$$

for every $s \in \mathbb{C}$, and the same holds for each segment separately.

Proof. For the S -segment put $I(s) := \int_{\hat{\gamma}_f^S} f(\tau) \tau^{s-1} d\tau$ and substitute $\tau \mapsto -1/\tau$. Since $f|_k S = f$ gives $f(-1/u) = u^k f(u)$, and $d\tau = du/u^2$,

$$I(s) = (-1)^{s-1} \int_{S\hat{\gamma}_f^S} f(u) u^{k-s-1} du.$$

By Lemma 4.29 the arc is carried by S onto itself with reversed orientation, so $S\hat{\gamma}_f^S = -\hat{\gamma}_f^S$ and $I(s) = -(-1)^{s-1}I(k-s)$. Multiplying by the prefactor $e^{-\pi is/2}$ of Definition 2.3 and using $(-1)^{s-1} = e^{i\pi(s-1)}$, the phases collect to $-e^{i\pi(k/2-1)} = e^{i\pi k/2}$, which is i^k .

For the T -segment no property of the path is used. Reading Definition 2.2 at $k-s$ and factoring out $-e^{i\pi(k-s-1)}$,

$$\tilde{k}_T(\tau, k-s) = -e^{i\pi(k-s-1)} \tilde{k}_T(\tau, s), \quad (76)$$

the two Hurwitz terms exchanging places; the identification of the second with the first requires $e^{-i\pi(k-s-1)} = e^{i\pi(s-1)}$, that is $e^{-i\pi(k-2)} = 1$, which is where k even enters. Hence $L_T^*(f, k-s) = i^k L_T^*(f, s)$, i.e. $L_T^*(f, s) = i^{-k} L_T^*(f, s')$ at $s' = k-s$. Finally $i^k = i^{-k} = (-1)^{k/2}$ for even k , so the two segments carry the same factor and (75) follows for the sum. $\square$

The two halves now assemble: every ingredient on the right is either a Laurent coefficient of f at one of its own poles, a Fourier coefficient of what survives projection, or one of the two explicit kernels I_N^{geo} , $G_{s,k}^{\text{geo}}$.

Theorem 4.24. *Let $f \in F_k$ and let $\{\tau_p\}$ be its T -inequivalent poles with $\text{Im } \tau_p \geq \frac{\sqrt{3}}{2}$, of orders m_p . Then the set $\{\tau_p\}$ is finite and, in the geodesic reference class,*

$$L^*(f, s) = \sum_p \sum_{m=1}^{m_p} r_{f, \tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} I_{m-1}^{\text{geo}}(s, \tau_p) + c_{Rf}(0) G_{s,k}^{\text{geo}}(0) + \sum_{n \neq 0} c_{Rf}(n) G_{s,k}^{\text{geo}}(n), \quad (77)$$

with $r_{f, \tau_p}^*(m)$ the pole weights of Definition 4.11, $c_{Rf}(n)$ the Fourier coefficients of Definition 4.14, and $I_N^{\text{geo}}(s, z)$ given by (65) for $\text{Im } z > \frac{\sqrt{3}}{2}$ and by (70) for $\text{Im } z = \frac{\sqrt{3}}{2}$. Both sides are entire in s away from the pole of $G_{s,k}^{\text{geo}}(0)$, the sum over n converges absolutely, and both sides satisfy

$$L^*(f, s) = i^k L^*(f, k-s). \quad (78)$$

If some τ_p lies on γ^{arc} the identity holds on either indentation of Proposition 4.9, the two sides changing together by (55); if some τ_p is elliptic, both sides are read at $W = 0$ through Definition 4.7, each block $I_{m-1}^{\text{geo}}(s, \tau_p)$ along with the left-hand side. No pole configuration is excluded.

Proof. Finiteness: f is meromorphic at the cusp, so it has no poles above some height, and its poles are locally finite in the strip $|\text{Re } \tau| \leq \frac{1}{2}$; the region cut out by $\text{Im } \tau \geq \frac{\sqrt{3}}{2}$ therefore contains finitely many of them modulo T . Definition 4.12 splits $f = \hat{P}^{\text{geo}} f + Rf$ with $\hat{P}^{\text{geo}} = \sum_p \hat{P}_{\tau_p}$ over exactly that finite set. Both segment integrals of Definition 2.3 are linear in the integrand, and the splitting is a finite sum, so L^* distributes over it. Lemma 4.21 evaluates each $L^*(\hat{P}_{\tau_p} f, s)$ as the inner double sum, and Lemma 4.22 evaluates $L^*(Rf, s)$ as the remaining two terms, with the convergence asserted there. The description of I_N^{geo} is Lemma 4.19 off the threshold line and Lemma 4.20 on it, the two agreeing where both apply. Equation (78) is Lemma 4.23, which applies to f itself and so to the left-hand side; the right-hand side inherits it, being equal to the left.

For a pole on γ^{arc} the blocks are integrated on an indented contour, and the ε_z bookkeeping preceding Lemma 4.19 records which indentation was taken; changing it changes the left-hand side by (55) and the corresponding block by the same residue, by Proposition 4.9. For an elliptic pole no single indentation exists, and Definition 4.7 supplies the value as the weighted average of two admissible classes; (77) holds in each of them, hence in the average, both sides being affine in the winding of Definition 4.6. $\square$

The formula is explicit in the sense that matters for computation: the pole weights are contour integrals of f about its own poles, the $c_{Rf}(n)$ are Fourier coefficients of an explicitly constructed T -periodic function, and the two kernels are incomplete gammas and polylogarithms. Nothing is left implicit except the arithmetic of f itself.

Lemma 4.19 assumes $\text{Im } z > \frac{\sqrt{3}}{2}$ strictly, so it says nothing on the threshold line itself, where the chord meets the pole and neither the defining series of Li_{-N} nor its inversion converges. That line is treated separately below, away from ρ and at it.

4.2 Relations to the L-functions of [3]

We pause to connect the L-functions of this section to the definition in Ref. [3].

Compared term by term, the two look different: the $1/s$ residue of [3] is proportional to $c_f(0)$ alone, whereas ours receives a contribution from every simple pole above the contour. The difference is one of reference class and not of object. Ref. [3] anchors its contour implicitly at $\tau_0 \rightarrow i\infty$, which was legitimate in its setting because poles at the cusp were excluded there, so that the limit is smooth; the present note anchors at $\tau_0 = \rho + 1$ on the boundary of the fundamental domain, by Definition 4.4. Moving between the two crosses pole orbits, and every discrepancy between the two definitions is the corresponding sum of residues.

Remark 4.25. The $1/s$ residue is carried entirely by the T -segment. L_S^* is a finite integral of $f \tau^{s-1}$ and is entire in s , while $\tilde{k}_T$ has a simple pole at $s = 0$ whose coefficient is the integral of f along that segment, the constant Fourier mode of f in the strip containing it. In the geodesic class that integral is $\Phi(f)$ of Definition 4.16, so $\text{Res}_{s=0} L^*(f, s) = -\Phi(f)$, whereas Ref. [3] anchors its contour at the cusp, above every pole, and carries $-c_f(0)$. The two differ by the residues lying in the strip between the contours, which is Lemma 4.8. The quantity that must be subtracted to reach W is therefore the same object as the departure from the residue convention of [3], and not a second normalisation imposed on top of it.

Remark 4.26. The mechanism is already visible in (63), before any assembly. Its S -part $e^{-\pi i s/2}(\tau_0^s - \rho^s)/s$ is finite at $s = 0$, since $\tau_0^s - \rho^s$ vanishes like $s \log(\tau_0/\rho)$, while $-(\tau_0/i)^s/s$ carries the pole. The two classes sit at opposite ends of one deformation: the contour of [3] lies above every pole, making the residue bookkeeping trivial and the pole-free region empty, while γ^{arc} lies below every pole orbit not on it, making the pole-free region of codimension zero and the residue bookkeeping maximal.

That deformation is the whole of the relation between the two definitions, and it needs no matching of q -series coefficients.

Proposition 4.27. *Let $f \in F_k$, let $\hat{\gamma}^{T\uparrow}$ be the T -segment deformed to pass above every pole of f with the S -segment left on γ^{arc} , and write $L^{*\uparrow}(f, s) := L^*(f, s; [\hat{\gamma}_f^S], [\hat{\gamma}^{T\uparrow}])$. Then $L^{*\uparrow}$ is the L-function of [3], and*

$$L^*(f, s) = L^{*\uparrow}(f, s) - 2\pi i e^{-\pi i s/2} \sum_{\tau_p} X_T(\tau_p) r_T(f, \tau_p), \quad (79)$$

the sum running over the pole orbits of f lying between γ^{arc} and $\hat{\gamma}^{T\uparrow}$, with X_T their crossing numbers and r_T as in Lemma 4.8.

Proof. Only the T -segment moves, so $X_S \equiv 0$ and Lemma 4.8 gives (79) directly. For the identification, L_S^* is entire in s , while the $s = 0$ pole of $\tilde{k}_T$ has coefficient $\int_{\hat{\gamma}_T} f d\tau$; on $\hat{\gamma}^{T\uparrow}$ that integral is the constant Fourier mode at the cusp, so $\text{Res}_{s=0} L^* = -c_f(0)$, which is the residue convention of [3], and by Lemma 4.3 nothing further depends on where above the poles the segment is placed. $\square$

One might then ask why the arc is used at all, since the raised class has the simpler residue bookkeeping. The answer is that the period polynomial does not survive there. Raising $\hat{\gamma}^T$ alone makes $\hat{\gamma}^S \neq \hat{\gamma}^T$, and Lemma 4.33 then gives a non-zero $(1 + U + U^2)$ defect, proportional to the residues at the poles the two segments separate. The arc is therefore not a convenient choice among equivalent ones: it is the class in which $\hat{r}_f$ lies in W , and (79) says exactly what is given up in exchange for the simpler residue convention.

4.3 Period polynomials in the geodesic class

Definition 4.28. For $f \in F_k$, let r_f be given by Definition 3.7 with $L^*(f, s)$ evaluated in the geodesic reference class of Definition 4.4, and set

$$\hat{r}_f := r_f - \Phi(f) r_{E_k}. \quad (80)$$

$\mathrm{SL}_2(\mathbb{Z})$ is generated by S and T subject to $S^2 = (ST)^3 = -I$, so two closed paths control the relations. Their degeneration on the arc is a statement about paths alone, independent of f ; we record it separately from its consequence, since the case of a pole *on* the contour uses the geometry but not the consequence.

Lemma 4.29. Let τ_0 lie on γ^{arc} and let $\hat{\gamma}^S = \hat{\gamma}^T$. Then the closed path traced by S^2 retraces itself, and if moreover $\tau_0 = \rho + 1$ the closed path traced by $(ST)^3$ retraces itself. Both statements are independent of f .

Proof. S acts on the unit circle by $e^{i\theta} \mapsto e^{i(\pi-\theta)}$, preserving $\frac{\pi}{3} \leq \theta \leq \frac{2\pi}{3}$, reversing orientation and fixing i ; so $S\hat{\gamma}^S$ is $\hat{\gamma}^S$ reversed and the S^2 -path retraces, for any base point on the arc. For $(ST)^3$, the fixed points of $U = TS$ are the roots of $\tau^2 - \tau + 1$, namely $e^{\pm i\pi/3}$, so $U\tau_0 = \tau_0$ exactly at $\tau_0 = \rho + 1$ and the U -path is closed. With $\hat{\gamma}^S = \hat{\gamma}^T$ it is $S^{-1}\hat{\gamma}^T$ followed by $\hat{\gamma}^S$, the arc traversed and then retraced; its U - and U^2 -transports retrace $U\gamma^{\mathrm{arc}}$ and $U^2\gamma^{\mathrm{arc}}$ through the same fixed point. $\square$

Lemma 4.30. In the geodesic reference class, $\hat{C}_{\pm I} = 0$ for every $f \in F_k$ with no pole on $\mathrm{SL}_2(\mathbb{Z})\gamma^{\mathrm{arc}}$.

Proof. By Lemma 4.29 both relator paths retrace, so each bounds nothing; the winding number about every pole of f vanishes and Cauchy's theorem gives $\hat{C}_{\pm I} = C_{\pm I} = 0$. $\square$

Definition 4.28 covers those f whose poles avoid the contour, and since γ^{arc} is a curve that is the generic situation: the excluded locus is of codimension one in $\mathbb{H}$, so $\hat{r}_f$ is defined outright on a set of codimension zero. The remaining cases are reached by continuity from it. Three cases exhaust the possibilities:

1. f has no pole on $\mathrm{SL}_2(\mathbb{Z})\cdot\gamma^{\mathrm{arc}}$;
2. f has a pole on γ^{arc} , away from i and ρ ;
3. f has a pole at i or at ρ .

Lemma 4.31. Let $f \in F_k$ have no pole on $\mathrm{SL}_2(\mathbb{Z})\cdot\gamma^{\mathrm{arc}}$. Then $\hat{r}_f \in W$, with no condition on the location or the order of the poles of f .

Proof. Throughout, $\hat{C}$ is the cocycle of Lemma 3.9, so that $\hat{C}_S = r_f$ and $\hat{C}_{\gamma\delta} = \hat{C}_\gamma|_\delta + \hat{C}_\delta$.

For the first relation, taking $\gamma = \delta = S$ gives $\hat{C}_{S^2} = \hat{C}_S|_S + \hat{C}_S = r_f|_{(1+S)}$, and $S^2 = -I$, so Lemma 4.30 gives

$$r_f|_{(1+S)} = \hat{C}_{-I} = 0. \quad (81)$$

For the second, take $\gamma = T$, $\delta = S$, so that $U = TS$ and $\hat{C}_U = \hat{C}_T|_S + \hat{C}_S$. Applying the cocycle relation twice more,

$$\hat{C}_{U^3} = \hat{C}_U|_{U^2} + \hat{C}_{U^2} = \hat{C}_U|_{U^2} + \hat{C}_U|_U + \hat{C}_U = \hat{C}_U|_{(1+U+U^2)},$$

and $U^3 = -I$, so Lemma 4.30 makes the left side vanish. Substituting $\hat{C}_U = \hat{C}_T|_S + \hat{C}_S$ and using $\hat{C}_S = r_f$,

$$0 = \hat{C}_T|_{S(1+U+U^2)} + r_f|_{(1+U+U^2)}, \quad \text{so} \quad r_f|_{(1+U+U^2)} = -\hat{C}_T|_{S(1+U+U^2)}. \quad (82)$$

This is where the hypothesis $f \in S_k^!$ was used in Lemma 3.11, and it is the only place: there $\hat{C}_T$ vanished, here it need not.

The T -cochain is controlled by Φ alone, whatever path represents the T -segment. We isolate this, since it is used again below for segments that wind.

Lemma 4.32. *There is a constant A_0 , depending on k alone, such that for every $f \in F_k$ and every admissible $\hat{\gamma}_f^T$,*

$$\hat{C}_T = -(2\pi i)^{n+1} A_0 \int_{\hat{\gamma}_f^T} f(\tau) d\tau = -(2\pi i)^{n+1} A_0 \Phi(f). \quad (83)$$

In particular the right-hand side depends on the homotopy class of $\hat{\gamma}_f^T$ only through $\Phi(f)$, and Φ is by Definition 4.16 the integral over that same path.

Proof. The computation of $\hat{C}_T$ in the proof of Lemma 3.9 does not use $c_f(0) = 0$ until its final step. It shows that $A(\tau) := \chi(\tau; X + Y, Y) - \chi(\tau - 1; X, Y)$ is a 1-periodic polynomial, hence a constant A_0 depending on k alone, and leaves $\hat{C}_T$ as the stated multiple of $\int_{\hat{\gamma}_f^T} f d\tau$. No property of the path enters beyond its endpoints, which are fixed by Definition 2.1. $\square$

Since A_0 does not depend on f , neither does $\mathcal{D} := (2\pi i)^{n+1} A_0|_{S(1+U+U^2)}$, and (82) becomes

$$r_f|_{(1+U+U^2)} = \Phi(f) \mathcal{D}. \quad (84)$$

The defect is therefore proportional to $\Phi(f)$, with a constant of proportionality common to every $f \in F_k$ whose poles avoid $\text{SL}_2(\mathbb{Z}) \cdot \gamma^{\text{arc}}$.

It remains to identify $\mathcal{D}$. Apply (84) to $f = E_k$, which is holomorphic and so certainly admissible, and for which $\Phi(E_k) = 1$: this gives $\mathcal{D} = r_{E_k}|_{(1+U+U^2)}$. Hence

$$\hat{r}_f|_{(1+U+U^2)} = r_f|_{(1+U+U^2)} - \Phi(f) r_{E_k}|_{(1+U+U^2)} = \Phi(f) \mathcal{D} - \Phi(f) \mathcal{D} = 0,$$

while (81) applied to both f and E_k gives $\hat{r}_f|_{(1+S)} = 0 - \Phi(f) \cdot 0 = 0$. Both defining relations of W hold. Nothing in the argument constrains where the poles of f lie, or how deep they are, beyond their absence from $\text{SL}_2(\mathbb{Z}) \cdot \gamma^{\text{arc}}$, which was used only through Lemma 4.30. $\square$

Lemma 4.33. *Let $f \in F_k$ have poles on γ^{arc} , none at i and none on $\text{SL}_2(\mathbb{Z}) \cdot \rho$, and let $\hat{\gamma}^S$ be S -symmetric. Write $D_E := r_{E_k}|_{(1+U+U^2)}$. Then $\hat{r}_f|_{(1+S)} = 0$ for every admissible $\hat{\gamma}^T$. If $\hat{\gamma}^T$ passes on the same side of every pole as $\hat{\gamma}^S$ then $\hat{r}_f|_{(1+U+U^2)} = 0$; otherwise, with $p_1, \dots, p_m$ the poles at which the two segments part,*

$$\hat{r}_f|_{(1+U+U^2)} = -(2\pi i)^{n+2} \sum_{j=1}^m \nu_j a_{p_j} (X - p_j Y)^n - 2\pi i \left(\sum_{j=1}^m X_T(p_j) a_{p_j} \right) D_E, \quad (85)$$

where $a_p = \text{Res}_{\tau=p} f$ and $\nu_j \in \mathbb{Z} \setminus \{0\}$, and this is non-zero unless D_E lies in the span of $\{(X - p_j Y)^n\}_{j \leq m}$.

Proof. By Lemma 3.9, $\hat{C}_S = r_f$ with $\hat{C}_S = C_S - (E|_S - E)$ and $C_S = (2\pi i)^{n+1} \int_{\hat{\gamma}^S} \omega_f$; the T -segment enters r_f only through the coboundary $E - E|_S$. The cocycle relation gives $r_f|_{(1+S)} = \hat{C}_S|_{(1+S)} = \hat{C}_{S^2}$, and as n is even the coboundary drops, leaving the integral of ω_f over the closed path formed by $\hat{\gamma}^S$ followed by its S -image. In polar coordinates S acts by $(r, \theta) \mapsto (1/r, \pi - \theta)$, so an S -symmetric $\hat{\gamma}^S$ is carried onto itself with reversed orientation; that path retraces and the integral vanishes. No property of $\hat{\gamma}^T$ is used, and the subtracted term contributes nothing since $r_{E_k}|_{(1+S)} = 0$.

For the second relation, Φ is by construction the quantity $\hat{C}_T$ measures, so Definition 4.28 gives $\hat{C}_T = 0$; then $\hat{C}_U = \hat{C}_T|_S + \hat{C}_S = \hat{C}_S$ and $\hat{r}_f|_{(1+U+U^2)} = \hat{C}_{U^3} - \Phi(f)D_E$, with $\hat{C}_{U^3} = \hat{C}_{-I}$ the integral of ω_f over the closed path traced by $(TS)^3$. As $U = TS$, that path is assembled from both segments. If $\hat{\gamma}^S = \hat{\gamma}^T$ it retraces, as in Lemma 4.30, and the two terms cancel exactly as in Lemma 4.31.

If the segments pass on opposite sides at $p_1, \dots, p_m$, the two paths bound regions containing exactly those poles and no others, and the residue theorem gives the first term of (85); the second is the shift induced on Φ , which is the T -segment integral and therefore moves with $\hat{\gamma}^T$. For the last claim, the coefficient vector of $(X - pY)^n$ is $\left(\binom{n}{l}(-p)^l\right)_l$, so the matrix of $\{(X - p_j Y)^n\}_{j \leq m}$ is $\text{diag}\left(\binom{n}{l}\right)$ times a Vandermonde in $(-p_1, \dots, -p_m)$, of rank m whenever the p_j are distinct and $m \leq n+1$. The residue terms therefore cannot cancel among themselves, and (85) can vanish only if D_E lies in their span. $\square$

The shifted contour realises one of the two sides without a second prescription being needed.

Lemma 4.34. *Let f be as in Lemma 4.33. Then for every $\epsilon > 0$ the pair $(\hat{\gamma}_\epsilon^S, \hat{\gamma}_\epsilon^T)$ of Definition 4.5 passes on the same side of every pole of f on γ^{arc} , and $\hat{r}_f$ computed on it is independent of ϵ and equals the value of Lemma 4.33 on that pair.*

Proof. The two segments differ only inside $|\tau - \rho| = O(\epsilon)$, where by hypothesis f has no pole, so they pass on the same side of each pole on the arc and Lemma 4.33 applies with $m = 0$ in (85). Independence of ϵ is Lemma 4.3: varying ϵ slides τ_0 radially without crossing a pole, the poles on the arc lying at $|\tau| = 1$ and the sides taken by (51) being the same for every $\epsilon > 0$. $\square$

The two relations are therefore sensitive to the two segments in quite different ways. The S -relation constrains $\hat{\gamma}^S$ alone: S -symmetry makes the S^2 -loop degenerate whatever $\hat{\gamma}^T$ does. The $(1 + U + U^2)$ relation mixes them, since $U = TS$, and vanishes only when the two segments agree. Requiring $\hat{\gamma}^S = \hat{\gamma}^T$ in Definition 4.4 is thus not a stipulation of convenience: membership in W selects it.

What it does not do is select a single class. The two conditions just isolated are preserved by a whole family of deformations, and $\hat{r}_f$ is linear in the contour — both r_f , through the special values of Definition 3.7, and Φ are integrals over the segments — so the family may be combined.

Corollary 4.35. *Let $f \in F_k$ and let $\Gamma_1, \dots, \Gamma_N$ be reference pairs at $\tau_0 = \rho + 1$, each with $S\hat{\gamma}_j^S = -\hat{\gamma}_j^S$ as a chain and $\hat{\gamma}_j^T = \hat{\gamma}_j^S$. Then $\hat{r}_f(\Gamma_j) \in W$ for every j , and for any $w_1, \dots, w_N \in \mathbb{C}$ with $\sum_j w_j = 1$,*

$$\sum_j w_j \hat{r}_f(\Gamma_j) \in W. \quad (86)$$

The values so obtained form a coset of $\text{span}_{\mathbb{C}}\{v_p\} \subseteq W$, where v_p is the shift produced by one unit of winding about a pole p and, as the previous condition forces, -1 unit about Sp , taken on both segments at once:

$$v_p = \left[(2\pi i)^{n+1} (-1)^\ell \binom{n}{\ell} 2\pi i (r_S + r_T)(f, p, \ell + 1) \right]_{\ell=0}^n - \left[p \mapsto Sp \right] - 2\pi i (a_p - a_{Sp}) r_{E_k}, \quad (87)$$

with r_S, r_T as in Lemma 4.8, evaluated at $s = \ell + 1$, and $a_q = \text{Res}_{\tau=q} f$.

Proof. Each Γ_j satisfies the hypotheses of Lemma 4.29, so both relator paths retrace and $\hat{C}_{\pm I} = 0$; the argument of Lemma 4.31 then gives both defining relations, whatever the winding data, so $\hat{r}_f(\Gamma_j) \in W$. For (86), the segment integrals of Definition 2.3 are linear in the contour as a chain, hence so are L^* , r_f and Φ , hence so is $\hat{r}_f = r_f - \Phi(f)r_{E_k}$; and W is a linear subspace. The shift (87) is Lemma 4.8 at $X_S(p) = X_T(p) = 1$ and $X_S(Sp) = X_T(Sp) = -1$, assembled through Definition 3.7, whose $i^{\ell+1}$ cancels the prefactor $e^{-i\pi s/2}$ of Definition 2.3 at $s = \ell + 1$, and corrected by the induced change $\Delta\Phi = 2\pi i(a_p - a_{Sp})$ in the subtraction. The two winding numbers are opposite because $S\hat{\gamma}^S = -\hat{\gamma}^S$ is preserved only by windings that reverse under S ; a single unit about p alone leaves the family. $\square$

The generator v_p deserves more than this incidental role, and Section 5 gives it one. It is $\Xi_{f;p}$ of Definition 5.2, an integral of f against a single kernel over the closed contour $c_p - Sc_p$, and Proposition 5.5 proves $\Xi_{f;p} \in W$ outright — for a pole of any order, with no base point, no homotopy class and no period polynomial entering. The antisymmetry demanded here is exactly what that proof consumes: (96) is the relation $(1 - S)(1 + S) = 1 - S^2 = 0$, and (97) records what the symmetric choice costs instead.

Two consequences worth separating. First, W -membership is a property of the whole symmetric family and not a selector within it: the residual freedom is exactly $\dim \text{span}_{\mathbb{C}}\{v_p\}$, indexed by the S -orbits of poles the contour can wind around, and Proposition 4.9 is the case of a single such orbit. Second, (86) does not require the combination to be realised by any one contour, only that each constituent be. This is what Definition 4.7 uses at an elliptic point, where $W = 0$ is not attained by a single path but is an affine combination of two that are.

Lemma 4.36. *Let $f \in F_k$ have a pole on $\text{SL}_2(\mathbb{Z}) \cdot \rho$ and let $W = W(f, \epsilon)$ be as in Definition 4.6. Then $W \equiv -1 \pmod{3}$; every such value is realised by some connector; and there is a vector $c \in V_n$, independent of W , with*

$$\hat{r}_f|_{(1+S)} = 0, \quad \hat{r}_f|_{(1+U+U^2)} = -cW. \quad (88)$$

Proof. Write $\sigma := \hat{\gamma}_\epsilon^S$ for the spiral, from $S\tau_0$ to τ_0 , and β for the connector of Definition 4.5, from $T^{-1}\tau_0$ to $S\tau_0$, so that $\hat{\gamma}_\epsilon^T = \beta \cdot \sigma$. With γ_g a path from τ_0 to $g\tau_0$ and the cocycle rule $\gamma_{gh} = \gamma_g \cdot (g\gamma_h)$, we have $\gamma_S = \bar{\sigma}$ and $\gamma_T = T\hat{\gamma}_\epsilon^T = (T\beta)(T\sigma)$, whence

$$\gamma_U = \gamma_T \cdot (T\gamma_S) = (T\beta)(T\sigma)(T\bar{\sigma}) \simeq T\beta, \quad (89)$$

the spiral occurring once forwards and once reversed. The cancellation is exact, not an estimate: $\hat{\gamma}_\epsilon^S$ appears verbatim inside $\hat{\gamma}_\epsilon^T$, so $(T\sigma)(T\bar{\sigma})$ is a path followed by its own reverse and is null-homotopic rel endpoints. Hence γ_U is homotopic to the T -translated connector, which lies in the disc $|\tau - (\rho + 1)| = O(\epsilon)$.

Since U fixes $\rho + 1$, each $U^j\gamma_U$ lies in a disc of the same size about the same point, so the relator path $\gamma_{U^3} = \gamma_U \cdot (U\gamma_U) \cdot (U^2\gamma_U)$ — closed, as $U^3 = -I$ — is supported in one $O(\epsilon)$ disc about $\rho + 1$. This is the analogue for $\epsilon \neq 0$ of Lemma 4.29, which is the case β trivial.

In the coordinate w of Definition 4.6 the map U fixes $w = 0$ and is exactly the rotation $w \mapsto \omega^{-1}w$, $\omega = e^{2\pi i/3}$, since $U'(\rho + 1) = (\rho + 1)^{-2} = e^{-2\pi i/3}$; winding about $\rho + 1$ is winding about $w = 0$, and U preserves increments of $\arg w$ exactly. The three pieces are U -translates of one another, hence each changes $\arg w$ by the same amount Δ , and the endpoints $\tau_0(\epsilon), U\tau_0(\epsilon), U^2\tau_0(\epsilon)$ sit at arguments $\frac{\pi}{3}, -\frac{\pi}{3}, -\pi$, so $\Delta = -\frac{2\pi}{3} + 2\pi k$ with $k \in \mathbb{Z}$ the class of β . Then $2\pi W = 3\Delta$ gives $W = -1 + 3k$, and every k occurs.

For (88), the S -relation holds because $\hat{\gamma}_\epsilon^S$ is S -symmetric, so the S^2 -path retraces by Lemma 4.29 and $\hat{C}_{S^2} = 0$ whatever $\hat{\gamma}_\epsilon^T$ does; as $r_{E_k}|_{(1+S)} = 0$ the subtraction contributes nothing. For the U -relation, (82) gives $r_f|_{(1+U+U^2)} = -\hat{C}_T|_{S(1+U+U^2)} - \hat{C}_{U^3}$, and Lemma 4.32 holds at every winding, so the first term is $\Phi(f)\mathcal{D}$ with $\mathcal{D} = D_E$ exactly as in Lemma 4.31; Definition 4.28 cancels it for every W . There remains $\hat{C}_{U^3}$, which by Cauchy's theorem is $2\pi i \sum_q W_q \text{Res}_q \omega_f$ over the poles q of f . By (89) the cycle lies in a single disc of radius $O(\epsilon)$ about $\rho + 1$. The poles of f are discrete and $\text{SL}_2(\mathbb{Z})$ -stable, hence locally finite, so there is $\epsilon_0 > 0$ below which that disc meets $f^{-1}(\infty)$ only at $\rho + 1$ itself. Therefore $W_q = 0$ for every $q \neq \rho + 1$, and $\hat{C}_{U^3} = 2\pi i W \text{Res}_{\rho+1} \omega_f$; setting $c := -2\pi i \text{Res}_{\rho+1} \omega_f|_{(1+U+U^2)}$ gives (88), and $c \neq 0$ because the defect is non-zero at $W = -1$. $\square$

Corollary 4.37. *For f as in Lemma 4.36, $\hat{r}_f \in W$ in the sense of Definition 4.7, and that class is the unique one in which it does.*

Proof. By (88) the defect vanishes exactly at $W = 0$, and $c \neq 0$ since the defect is non-zero at $W = -1$; uniqueness follows because $W \mapsto -cW$ is injective. The same argument at i , where U is replaced by S and the relator path is the S^2 -path, gives W odd and the prescription the mean of the two indentations. $\square$

Lemmas 4.33 and 4.36 are complementary rather than overlapping: the first excludes i and $\text{SL}_2(\mathbb{Z}) \cdot \rho$ by hypothesis, the second treats only those points, where the stabiliser forces W into a class that misses 0 and the L-integral must be read off the affine family rather than from a single contour. Together with Lemma 4.31 they cover every pole configuration.

Theorem 4.38. *For $f \in F_k$, and $W \subset V_n$ from Definition 3.10, the period polynomial $\hat{r}_f(X, Y) \in W$.*

Proof. Let $E \subset f^{-1}(\infty)$ be the poles of f lying on $\text{SL}_2(\mathbb{Z}) \cdot \rho$ or on $\text{SL}_2(\mathbb{Z}) \cdot i$, and N those lying on γ^{arc} but not at an elliptic point; the remaining poles meet γ^{arc} nowhere. The three mechanisms are local to these three families and may be applied together.

If E and N are both empty, Lemma 4.31 gives $\hat{r}_f \in W$ directly, with no condition on where the other poles lie or how deep they are.

If E is empty and N is not, Definition 4.4 takes $\hat{\gamma}_f^S = \hat{\gamma}_f^T$, and both relations hold by Lemma 4.33; the shifted contour of Definition 4.5 gives the same value by Lemma 4.34, so the two descriptions do not compete.

If E is non-empty, read L^* and $\hat{r}_f$ off the shifted class of Definition 4.5 at $W = 0$, as in Definition 4.7, at each elliptic point of E in turn. Lemma 4.36 shows the two defining relations vanish there, and Corollary 4.37 that $W = 0$ is the only class in which they do. Poles of f elsewhere are untouched and fall under the two previous cases. Hence $\hat{r}_f \in W$. $\square$

The construction uses only the $k - 1$ special values $L^*(\cdot, \ell + 1)$, $0 \leq \ell \leq n$, of the finite-contour L -function of Section 4.1, together with the single subtraction $\Phi(f)r_{E_k}$. No reference family is chosen, and no projection onto W is performed.

Two choices enter, and neither moves $\hat{r}_f$ out of W . In case 2 the side of the indentation is free, and by Lemma 4.33 the two differ within W . In case 3 nothing is free: Corollary 4.37 singles out $W = 0$, and ϵ is inert by Lemma 4.3. In Definition 4.28 the reference form E_k may be replaced by any g with $\Phi(g) = 1$, and two such choices shift $\hat{r}_f$ by $\Phi(f)r_{g-g'}$, where $g - g'$ runs over the forms annihilated by Φ ; how large that shift space is depends on the class from which g is drawn, and we do not settle it here.

A pole of prescribed order at a prescribed point is cut out by $J - J(z)$, and for that family $\hat{r}_f$ is an explicit series in $J(z)^{-1}$ whose coefficients do not depend on z .

Proposition 4.39. *Let $h \in M_{k-2}$, let $z \in \mathbb{H}$ with $|J(z)| > 984$, and set $f_z := h J'/(J - J(z)) \in F_k$. Then*

$$\hat{r}_{f_z} = - \sum_{m \geq 1} \hat{r}_{h J^{m-1} J'} J(z)^{-m}, \quad (90)$$

absolutely convergent in V_n , with $h J^{m-1} J' \in M_k^!$ for every m .

Proof. On γ^{arc} we have $S\tau = -\bar{\tau}$, so $J(\tau) = J(S\tau) = \overline{J(\tau)}$ is real there; it runs from $J(\rho) = -744$ to $J(i) = 984$ and back, whence $|J| \leq 984 < |J(z)|$ on the arc and

$$\frac{J'}{J - J(z)} = - \sum_{m \geq 1} J^{m-1} J' J(z)^{-m} \quad (91)$$

converges uniformly on it. By Definition 4.16 and Definition 3.7, Φ and each $L^*(\cdot, \ell + 1)$ are integrals over γ^{arc} against kernels continuous on that compact set, so both are linear and continuous under uniform convergence there, and $\hat{r}$ is assembled from them by (80). Term by term, $J' = -2\pi i E_4^2 E_6 / \Delta$ is holomorphic on $\mathbb{H}$ because Δ does not vanish there, so $h J^{m-1} J'$ has no pole off the cusp. $\square$

The $m = 1$ term is where the choice left open in Definition 4.28 becomes visible. Since $J' = -2\pi i (q^{-1} + O(q))$ carries no constant term, $h J'$ has constant term $-2\pi i c_h(1)$; so when $c_h(1) \neq 0$ the form $h J' / (-2\pi i c_h(1))$ is itself admissible as the reference of Definition 4.28, and with that choice $\hat{r}_{h J'} = 0$ and (90) begins at $m = 2$. Then $\hat{r}_{f_z} = O(J(z)^{-2})$, which is $O(e^{-4\pi \text{Im } z})$ rather than $O(e^{-2\pi \text{Im } z})$. Any other admissible reference restores the $m = 1$ term, by the shift recorded above. The family thus selects a reference form from that shift space, whereas f alone does not.

5 Residue polynomials

As is profoundly clear in the sub-section 4.3, the route to assigning a period polynomial to a weight- k $\text{SL}_2(\mathbb{Z})$ meromorphic modular form $f \in F_k$, entirely via its underlying L-function, is fraught with ambiguities. To be explicit:

1. To define $L^*(f, s) := L^*(f, s; [\hat{\gamma}_{\tau_0}^T], [\hat{\gamma}_{\tau_0}^S])$ associated with a meromorphic $f \in F_k$, one must choose a reference homotopy class. There are countably infinite number of distinct choices, here.
2. For our preferred reference contour, γ^{arc} , to define a period polynomial $\hat{r}_f \in W \subset V_{k-2}$, one must subtract $\Phi(f) r_{E_k}$. That subtraction is $c_f(0) r_{E_k}$ unless a pole lies between γ^{arc} and the horizontal line from $\rho \rightarrow \rho + 1$, in which case the residues in the intervening strip enter as well. But it is well-defined only modulo $r(\ker \Phi)$.

We were unable to naturally side-step either of these obstructions, either of which would render an attempt to make a canonical map between the space of meromorphic modular forms, $f \in F_k$, to the space of period polynomials $r_f \in W \subset V_{k-2}$, $f \in F_k \mapsto r_f \in W \subset V_{k-2}$. Indeed, both of these ambiguities are integrally tied to our definition of an L-function of a meromorphic modular form $f \in F_k$ via a contour integral and to our requirement that any candidate period polynomial of a modular form must lie in the privileged subspace $W \subset V_{k-2}$. This seems to imply some inherent difficulty in extending the notion of periods and quasi-periods of meromorphic modular forms, in the sense of this paper (i.e. $f \in F^* := \oplus_k F_k$) which is fundamentally tied to the nature of poles at finite locations in the fundamental domain.

Both obstructions, however, are statements about *open* contours: the first is the choice of a homotopy class joining $S\tau_0$ to τ_0 , and the second is the subtraction that repairs the resulting T -defect. A *closed* contour about a pole carries neither. What we show below is that the winding of such a contour is by itself enough: it produces an element of W , directly, with no base point, no homotopy class, and no period polynomial anywhere in the construction. This opens the door to defining essentially complementary objects — complementary in the sense that the objects are trivial/fail to exist for the privileged subspace of weakly holomorphic forms, $M_k^! \subset F_k$. To wit,

Definition 5.1. For $n = k - 2$, with $\mathbf{K}_T$ as in Proposition 3.8 and r_{E_k} as in Definition 4.28, set

$$\mathcal{K}(\tau; X, Y) := (2\pi i)^{n+1} \left[(X - \tau Y)^n + \mathbf{K}_T(\tau; X, Y) \right] - r_{E_k}(X, Y). \quad (92)$$

The subtraction r_{E_k} is constant in τ , so it contributes $2\pi i r_{E_k}$ times the enclosed residue of f and no more; it is the counterterm in the single non-parabolic direction, normalised by $\Phi(E_k) = 1$. A loop about a pole of nonzero residue is a non-parabolic cycle, and by the third display in the proof of Lemma 3.11 it is parabolicity, $\hat{C}_T = 0$, on which the relation $w|_{(1+U+U^2)} = 0$ rests.

Definition 5.2 (Residue polynomial for $f \in F_k$ at z). Let $f \in F_k$, let $z \in \mathbb{H}$ be a pole of f , let c_z be a simple, positively oriented loop enclosing z and no other pole of f , and set $\gamma_o(z) := c_z - S c_z$. We define the residue polynomial of f at z , $\Xi_{f;z}$ to be,

$$\Xi_{f;z} := \oint_{\gamma_o(z)} f(\tau) \mathcal{K}(\tau; X, Y) d\tau. \quad (93)$$

The loop is S -antisymmetric, $S\gamma_o = -\gamma_o$, and not merely a loop about the pair $\{z, Sz\}$. This is forced: Lemma 4.29 makes the S^2 - and $(ST)^3$ -paths retrace by $S\hat{\gamma}^S = -\hat{\gamma}^S$, so a winding added to both segments must reverse under S . The symmetric alternative $c_z + S c_z$ fails, and fails in the S -relation; see (97) below. Since S is holomorphic, $S c_z$ is a positively oriented loop about Sz , so γ_o encircles z once positively and Sz once negatively. At $z = i$ it degenerates to the zero cycle, because $Si = i$; correspondingly $\Xi_{f;i} = 0$. At $z \in \mathrm{SL}_2(\mathbb{Z}) \cdot \rho$ the two points are the arc endpoints ρ and $S(\rho + 1) = \rho$.

Write $P_\tau := (X - \tau Y)^n$, and for a 1-chain c in $\mathbb{H}$ avoiding the poles of f set

$$A(c) := \int_c f(\tau) P_\tau d\tau, \quad B(c) := \int_c f(\tau) \mathbf{K}_T(\tau; X, Y) d\tau. \quad (94)$$

Two facts about these, one for each kernel, give the result outright.

Lemma 5.3. For every $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ and every such chain c , $A(\gamma c) = A(c)|_{\gamma^{-1}}$.

Proof. Substitute $\tau = \gamma w$ in $A(\gamma c)$. Then $d\tau = (cw + d)^{-2} dw$ and $f(\gamma w) = (cw + d)^k f(w)$, while

$$X - (\gamma w)Y = \frac{(cX - aY)w + (dX - bY)}{cw + d},$$

so the integrand becomes $f(w)((cX - aY)w + (dX - bY))^n dw$ because $k - n - 2 = 0$. Since $\gamma^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$, that bracket is $P_w|_{\gamma^{-1}}$. $\square$

Lemma 5.4. $\mathbf{K}_T(\tau) - \mathbf{K}_T(\tau - 1) = -P_\tau|_{(1-S)}$, and consequently, in the sense of Hurwitz continuation, $\mathbf{K}_T(\tau) = \sum_{m \geq 1} P_{\tau+m}|_{(1-S)}$.

Proof. At $s = \ell + 1$ Definition 2.2 reads $\tilde{k}_T(\tau, \ell + 1) = \zeta(-\ell, \tau + 1) - (-1)^\ell \zeta(\ell - n, \tau + 1)$, and $\zeta(w, x) - \zeta(w, x + 1) = x^{-w}$ gives $\tilde{k}_T(\tau, \ell + 1) - \tilde{k}_T(\tau - 1, \ell + 1) = -\tau^\ell + (-1)^\ell \tau^{n-\ell}$. Assembling over ℓ with the weights of Proposition 3.8 yields $-P_\tau + (\tau X + Y)^n$, and $P_\tau|_S = (\tau X + Y)^n$ for n even. For the second form, $\zeta(-m, x)$ continues $\sum_{j \geq 0} (j + x)^m$, so $\zeta(-\ell, \tau + 1)$ continues $\sum_{m \geq 1} (\tau + m)^\ell$; assembling gives $\sum_{m \geq 1} (P_{\tau+m} - P_{\tau+m}|_S)$. $\square$

Proposition 5.5. $\Xi_{f;z} \in W$, for a pole of any order, and depends on neither τ_0 nor the reference class of Definition 4.2.

Proof. Put $\mathfrak{a} := A(c_z)$ and let $\mathcal{T} := \sum_{m \geq 1} T^{-m}$ act by $A(T^m c_z) = \mathfrak{a}|_{T^{-m}}$, which is Lemma 5.3. That lemma gives $A(\gamma_\circ) = \mathfrak{a}|_{(1-S)}$, and, since $P_{\tau+m} = P_\tau|_{T^{-m}}$, Lemma 5.4 gives $B(c_z) = \mathfrak{a}|_{\mathcal{T}(1-S)}$ and hence $B(\gamma_\circ) = \mathfrak{a}|_{(1-S)\mathcal{T}(1-S)}$. So with $E := 1 + \mathcal{T}(1-S)$,

$$A(\gamma_\circ) + B(\gamma_\circ) = \mathfrak{a}|_{(1-S)E}. \quad (95)$$

Now $S^2 = -I$ acts trivially on V_n for n even, so $(1-S)(1+S) = 1 - S^2 = 0$ and therefore

$$(1-S)E(1+S) = (1-S)(1+S) + (1-S)\mathcal{T}(1-S^2) = 0, \quad (96)$$

every term of the sum carrying the factor $1 - S^2$. With $r_{E_k}|_{(1+S)} = 0$, which holds by Lemma 4.30 and the second display in the proof of Lemma 3.11, this gives $\Xi_{f;z}|_{(1+S)} = 0$.

For the remaining relation, $\gamma_\circ$ changes Φ of Definition 4.16 by $\oint_{\gamma_\circ} f d\tau$, so the bracket (95) alone is not parabolic; the subtraction of r_{E_k} in (92), normalised by $\Phi(E_k) = 1$, restores $\hat{C}_T = 0$, and the third display in the proof of Lemma 3.11 then gives $\Xi_{f;z}|_{(1+U+U^2)} = 0$. Nothing in (93) refers to τ_0 or to a homotopy class, and no step above used the order of the pole. $\square$

The symmetric alternative fails at (96), and fails there only: $E(1+S) = (1+S)$, so

$$\left[\oint_{c_z + S c_z} f \mathcal{K} d\tau \right] \Big|_{(1+S)} = (2\pi i)^{n+1} \mathfrak{a}|_{(1+S)^2} = 2(2\pi i)^{n+1} \mathfrak{a}|_{(1+S)}, \quad (97)$$

which vanishes only when $\mathfrak{a}|_{(1+S)}$ does. The same computation with c_z alone gives $\mathfrak{a}|_{(1+S)}$, half of (97).

Further, we suspect that the related objects,

Definition 5.6 (Full residue polynomial for $f \in F_k$). Let $f \in F_k$, let Z denote a set of representatives, modulo $\mathrm{SL}_2(\mathbb{Z})$, for the pole locations of f in $\mathbb{H}$, let $\Xi_{f;z}$ be as in Definition 5.2, and let n_z denote the order of the stabiliser of z in $\mathrm{PSL}_2(\mathbb{Z})$, so that $n_z = 2$ on $\mathrm{SL}_2(\mathbb{Z}) \cdot i$, $n_z = 3$ on $\mathrm{SL}_2(\mathbb{Z}) \cdot \rho$, and $n_z = 1$ otherwise. We define the full residue polynomial of f , Ξ_f , to be,

$$\Xi_f := \sum_{z \in Z} \frac{\Xi_{f;z}}{n_z}, \quad (98)$$

$$R(f) := \sum_{z \in Z} \frac{1}{n_z} \left(\mathrm{Res}_{\tau=z} f - \mathrm{Res}_{\tau=Sz} f \right), \quad (99)$$

$$\Sigma_f := \Xi_f + 2\pi i R(f) r_{E_k}. \quad (100)$$

to also be very interesting in their own right, perhaps even more interesting.

The weights $1/n_z$ are those of the valence formula [4], which counts orders at i and at ρ with $\frac{1}{2}$ and $\frac{1}{3}$; in that sense Ξ_f is a divisor sum. Away from the elliptic points $f|_k S = f$ gives $\mathrm{Res}_{\tau=Sz} f = z^n \mathrm{Res}_{\tau=z} f$ at a simple pole, so the bracket in (99) is $(1 - z^n) \mathrm{Res}_{\tau=z} f$; at $z = i$ it vanishes, consistently with $\Xi_{f;i} = 0$.

Proposition 5.7. *Let $f \in F_k$. Then*

1. Ξ_f, Σ_f and R are linear in f , and vanish identically on $M_k^!$;
2. Σ_f is independent of the reference form, while replacing E_k in Definition 4.28 by any g with $\Phi(g) = 1$ sends $\Xi_f \mapsto \Xi_f + 2\pi i R(f) r_{E_k-g}$;
3. $\Xi_f \in W$, whereas $\Sigma_f|_{(1+S)} = 0$ and $\Sigma_f|_{(1+U+U^2)} = 2\pi i R(f) r_{E_k}|_{(1+U+U^2)}$.

In particular $\Xi_f = \Sigma_f$ is both free of the reference form and an element of W precisely when $R(f) = 0$.

Proof. Linearity is immediate from (93) and (98), and $Z = \emptyset$ for $f \in M_k^!$. For (2), r_{E_k} enters (92) additively and is constant in τ , so it contributes $-r_{E_k} \oint_{\gamma_\circ(z)} f d\tau = -2\pi i r_{E_k} (\text{Res}_{\tau=z} f - \text{Res}_{\tau=Sz} f)$ to (93); weighting by $1/n_z$ and summing over Z gives the stated shift, which (100) cancels. For (3), $\Xi_f \in W$ by Proposition 5.5. The second display in the proof of Lemma 3.11 uses only $\hat{C}_{\pm I} = 0$, which holds for E_k by Lemma 4.30, so $r_{E_k}|_{(1+S)} = 0$; the third display uses $\hat{C}_T = 0$, which fails for E_k since $\Phi(E_k) = 1$. The two identities follow from (100). $\square$

$R(f)$ is a canonical linear functional on F_k : no contour, base point or reference form enters it. It is the whole of the obstruction, and it does not vanish in general — for the family of Proposition 4.39, $Z = \{z\}$ and $\text{Res}_{\tau=z} f_z = h(z)$, so $R(f_z) = h(z)(1 - z^n)$.

At a simple pole Definition 5.2 is wholly explicit. Writing $\tilde{k}_T(w, \ell + 1) = -B_{\ell+1}(w + 1)/(\ell + 1) + (-1)^\ell B_{n-\ell+1}(w + 1)/(n - \ell + 1)$ for the Bernoulli polynomials of Definition 3.12, the coefficient of $X^{n-\ell}Y^\ell$ in $\Xi_{f;z}$ is

$$2\pi i a_z \left\{ (2\pi i)^{n+1} (-1)^\ell \binom{n}{\ell} \left[z^\ell - z^{n-\ell} + \tilde{k}_T(z, \ell + 1) - z^n \tilde{k}_T(-1/z, \ell + 1) \right] - (1 - z^n) (r_{E_k})_\ell \right\}, \quad (101)$$

$a_z = \text{Res}_{\tau=z} f$. So $\Xi_{f;z}$ is $a_z (2\pi i)^{n+2}$ times a polynomial in z with rational coefficients, less $2\pi i a_z (1 - z^n) r_{E_k}$: its only transcendental content beyond a_z and powers of $2\pi i$ is r_{E_k} , entering with the coefficient $(1 - z^n)$. Write $\Xi_{f;z} = a_z \Xi_k^\circ(z)$ for the resulting map $\Xi_k^\circ : \mathbb{H} \rightarrow W$, which depends on k alone.

Proposition 5.8. $\Xi_k^\circ(z) = \sum_{m=-1}^{n+1} c_m z^m$ with every $c_m \in W$, and

$$\Xi_k^\circ(-1/z) = -z^{-n} \Xi_k^\circ(z), \quad \text{equivalently} \quad c_{n-m} = (-1)^{m+1} c_m. \quad (102)$$

Proof. In (101), $\tilde{k}_T(w, \ell + 1)$ is a polynomial in w of degree $\max(\ell + 1, n - \ell + 1) \leq n + 1$. Hence $\tilde{k}_T(z, \ell + 1)$ contributes powers $z^0, \dots, z^{n+1}$, while $z^n \tilde{k}_T(-1/z, \ell + 1)$ contributes $z^{n-\max(\ell+1, n-\ell+1)}, \dots, z^n$, the least exponent being z^{-1} , attained at $\ell = 0$ and $\ell = n$; the remaining terms lie in that range. So Ξ_k° is a Laurent polynomial supported in $-1 \leq m \leq n + 1$. Since $\Xi_k^\circ(z) \in W$ for every z by Proposition 5.5, the monomials z^m are linearly independent as functions of z , and W is a linear subspace, each c_m lies in W .

For (102), S is holomorphic and $S^2 = 1$ in $\text{PSL}_2(\mathbb{Z})$, so $\gamma_\circ(Sz) = c_{Sz} - S c_{Sz} = c_{Sz} - c_z = -\gamma_\circ(z)$ and therefore $\Xi_{f;Sz} = -\Xi_{f;z}$. Lemma 5.3 at $\gamma = S$ gives $a_{Sz} = z^n a_z$, so $z^n a_z \Xi_k^\circ(-1/z) = -a_z \Xi_k^\circ(z)$. Comparing coefficients of z^m on the two sides of $\Xi_k^\circ(-1/z) = -z^{-n} \Xi_k^\circ(z)$ gives $(-1)^m c_m = -c_{n-m}$. $\square$

So the dependence of $\Xi_{f;z}$ on the pole location is rational: $z \mapsto \Xi_k^\circ(z)$ is a curve in W of degree at most $n + 2$, and (102) is an involution on it. Its coordinate functions in any basis of W are

Laurent polynomials in z , so the curve is contained in no proper subspace of W precisely when the c_m span; at $k = 12$, where $\dim W = 3$, they are computed to do so.

Two consequences for the ambiguity of Proposition 5.7. If f has a single simple pole orbit, then $R(f) = n_z^{-1}(1 - z^n)a_z$, which vanishes exactly when $z^n = 1$, that is when the pole lies on $|\tau| = 1$ at an n -th root of unity; for such f the residue polynomial is free of the reference form. At $k = 12$ the roots of unity meeting $\frac{\pi}{3} \leq \arg \tau \leq \frac{2\pi}{3}$ are $e^{2\pi i/5}$ and $e^{3\pi i/5}$, a single S -pair. And $\Xi_{f;z}$ determines a_z whenever $\Xi_k^\circ(z) \neq 0$, so within one orbit the whole residue data is recovered, by Lemma 5.3; across distinct orbits it is not, since Ξ_f records only the weighted sum.

A semi-structured list of reasons why, below:

- Ξ_f and $\Xi_{f;z}$ are inherently independent of τ_0 , and of the reference homotopy class. They just depend on the contour which encircles their residue, just as period polynomials are for the standard case $f \in M_k^!$.
- Ξ_f and $\Xi_{f;z}$ are naturally in $W \subset V_{k-2}$ for every $f \in F_k$ and every pole z , elliptic or not
- Ξ_f carries global information about the divisor structure of the modular form: it is a divisor sum, weighted at i and at ρ exactly as the valence formula weights them, and it records the whole polar part of f at each pole and not merely the residue
- Ξ_f denotes a unique polynomial in $W \subset V_{k-2}$ exactly when $R(f) = 0$; otherwise the reference form of Definition 4.28 survives in it, and $R(f)$ measures how much

Poles at the cusp do not enter. The kernel $\mathcal{K}$ is integrated against f on loops in $\mathbb{H}$, so only the polar parts of f at finite locations contribute, and no $c_f(0)$ appears anywhere. This is why Ξ_f vanishes on $M_k^!$, and it means that admitting or excluding poles at the cusp changes nothing.

5.1 An example localised in W at weight $k = 18$

Take $g = \Delta E_4 \in M_{16}$ and

$$f_z := g \frac{J'}{J - J(z)} \in F_{18}, \quad J' = -2\pi i (J + 744) \frac{E_6}{E_4}, \quad (103)$$

for z in the interior of the fundamental domain. Three things make this the tractable case. $J'/(J - J(z))$ has weight 2, so f_z has weight $k = 18$ and $n = 16$; f_z has a single pole orbit, so $\Xi_{f_z} = \Xi_{f_z;z}$ and the local and global residue polynomials coincide; and $\dim S_{18} = 1$, so $\dim W = 3$ and W has a cuspidal part rather than reducing to the Eisenstein line. By Lemma 5.3 the residue is $\text{Res}_{\tau=z} f_z = g(z)$, so

$$\Xi_{f_z} = \Delta(z) E_4(z) \Xi_{18}^\circ(z), \quad (104)$$

a weight-16 cusp form evaluated at z times the universal map of Proposition 5.8, which depends on k alone. In particular $|\Xi_{f_z}| \asymp e^{-4\pi \text{Im } z}$.

The parity involution $\varepsilon : P(X, Y) \mapsto P(-X, Y)$ preserves W : it acts as $(-1)^\ell$ on the coefficient of $X^{n-\ell} Y^\ell$, and $\varepsilon S \varepsilon^{-1} = -S$ while $\varepsilon U \varepsilon^{-1} = S U^{-1} S^{-1}$, so on $\ker(1 + S)$ the two U -relations agree. Since $p_0 = X^n - Y^n$ is ε -even,

$$\dim W^{\text{even}} = \dim S_k + 1, \quad \dim W^{\text{odd}} = \dim S_k. \quad (105)$$

At $k = 18$ the odd part is therefore a line. Since $\dim S_{18} = 1$, Definition 3.12 applies, and the three polynomials $W_-^{(18)}$, $W_+^{(18)}$, p_0 have *disjoint* supports in ℓ — odd ℓ ; $\ell \in \{2, 4, 6, 10, 12, 14\}$;

and $\ell \in \{0, 16\}$ — so coordinates in that basis are read off rather than solved for. Note $\ell = n/2$ is absent from $W_+^{(18)}$, its defining bracket having $a_i = -a_{n-i}$ and hence $a_{n/2} = 0$; the same midpoint forces $c_{n/2} = 0$ in (102) whenever $k \equiv 2 \pmod{4}$.

Writing $\Xi_k^\circ(z) = \sum_m c_m z^m$ and taking the reference form to be E_k itself, the coordinates of c_m in this basis obey a selection rule: the W_- coordinate is supported on even m , and the W_+ and p_0 coordinates on odd m . All three are rational multiples of $(2\pi i)^{n+2}$ with one exception. The exception is p_0 at $m \in \{0, n\}$, where the rational part vanishes and the coordinate is instead the single constant by which r_{E_k} fails to be rational — for $r_{E_{18}}$ the W_+ coordinate vanishes identically and the W_- coordinate is rational, so $r_{E_{18}}$ is rational along W_- and irrational only along p_0 . Consequently the rational and the transcendental parts of Ξ_k° occupy disjoint powers of z .

Explicitly, writing $\Xi_{f_z} = \Delta(z)E_4(z)[\mu_-(z)W_-^{(18)} + \mu_+(z)W_+^{(18)} + \mu_0(z)p_0]$ and dividing out $(2\pi i)^{18}$, $\mu_- = \frac{2}{131601}P_-$ and $\mu_+ = \frac{2}{3}P_+$, where

$$P_-(z) = -18000 + 350936 z^2 - 1096675 z^4 + 1140542 z^6 - 1140542 z^{10} + 1096675 z^{12} - 350936 z^{14} + 18000 z^{16}, \quad (106)$$

$$P_+(z) = -24 z + 154 z^3 - 273 z^5 + 143 z^7 + 143 z^9 - 273 z^{11} + 154 z^{13} - 24 z^{15} \quad (107)$$

have coprime integer coefficients, and $131601 = 3 \cdot 43867$ carries the numerator of B_{18} . The scalar $\Delta(z)E_4(z)$ and the normalisations of $W_\pm^{(18)}$ both cancel in the ratio, so the position of Ξ_{f_z} inside the plane spanned by $W_-^{(18)}$ and $W_+^{(18)}$ is the single rational function

$$R(z) := \frac{\mu_+(z)}{\mu_-(z)} = 43867 \frac{P_+(z)}{P_-(z)} \in \mathbb{P}^1. \quad (108)$$

Lemma 5.9. $R(-1/z) = R(z)$ and $R(-z) = -R(z)$, and $R(z) \in i\mathbb{R} \cup \{\infty\}$ for z on the imaginary axis and for $|z| = 1$.

Proof. By (102) each coordinate of Ξ_k° satisfies $\mu(-1/z) = -z^{-n}\mu(z)$, so the common factor cancels in (108); and μ_- is even in z while μ_+ is odd, which gives the second identity. For the third, $\mu_\pm$ have real coefficients, so $R(\bar{z}) = \overline{R(z)}$. On the imaginary axis $\mu_-(iy)$ is real and $\mu_+(iy)$ is i times a real number. For $|z| = 1$ one has $\bar{z} = 1/z$, hence $\overline{R(z)} = R(1/z) = -R(-1/z) = -R(z)$. $\square$

So the $W^\pm$ -localisation depends only on the $\mathrm{SL}_2(\mathbb{Z})$ -orbit of the pole through its S -class, and is purely imaginary exactly on the two reflection loci. Since R has rational coefficients, $R(z) \in \mathbb{Q}(z)$; at a point of class number one this is an explicit element of an imaginary quadratic field. The two lying on the reflection loci of Lemma 5.9 are

$$R\left(\frac{1+\sqrt{-3}}{2}\right) = \frac{43867}{8129}\sqrt{-3}, \quad R(\sqrt{-2}) = \frac{43867 \cdot 18}{119513}\sqrt{-2}, \quad (109)$$

purely imaginary as they must be, while off those loci both coordinates appear, as in

$$R\left(\frac{1+\sqrt{-7}}{2}\right) = \frac{43867(-107 + 316769\sqrt{-7})}{3931333248}, \quad R\left(\frac{1+\sqrt{-11}}{2}\right) = \frac{43867(-30607 + 4873558\sqrt{-11})}{75795816699}. \quad (110)$$

The remaining class-number-one points give elements of $\mathbb{Q}(\sqrt{-d})$ of the same shape and larger height. At $z = i$ both $\mu_\pm$ vanish and R is undefined, consistently with $\Xi_{f,i} = 0$.

These values are algebraic because R is a rational function with rational coefficients, and not because the pole is a CM point. The arithmetic of Ξ_{f_z} sits entirely in the scalar $\Delta(z)E_4(z)$, a CM value of a weight-16 cusp form, which (108) divides out.

Both $\mu_{\pm}$ vanish at $z = \pm i$: in the variable $w = z^2$, μ_+/z is palindromic of odd degree and μ_- anti-palindromic of degree 8, so $w = -1$ is a root of each. This is the algebraic shadow of $\Xi_{f,i} = 0$. The anti-palindromy gives μ_- the further roots $z = \pm 1$.

The reference form matters here and not only in Proposition 5.7. Replacing E_k by another g with $\Phi(g) = 1$ changes r_{E_k} by the period polynomial of a cusp form, which has nonzero W_- and W_+ coordinates; the rationality above is a statement about the Eisenstein series and is destroyed by such a change.

References

- [1] K. Bringmann, K.-H. Fricke, and Z. Kent, *Special L-values and periods of weakly holomorphic modular forms*, Ramanujan J. **24** (2011), 119–139.
- [2] F. Brown, *A class of non-holomorphic modular forms III*, arXiv:1710.07912 (2017).
- [3] D. A. McGady, *L-functions for meromorphic modular forms and sum rules in conformal field theory*, arXiv:1806.09874 (2018).
- [4] T. M. Apostol, “Modular functions and Dirichlet series in number theory,” Graduate Texts in Mathematics, London, Springer (1997).
- [5] W. Duke and P. Jenkins, *On the zeros and coefficients of certain weakly holomorphic modular forms*, Pure Appl. Math. Q. **4** (2008), no. 4, 1327–1340.
- [6] W. Kohnen and D. Zagier, *Modular forms with rational periods*, in *Modular Forms* (R. A. Rankin, ed.), Ellis Horwood, 1984, pp. 197–249.
- [7] F. Brown and R. Hain, *Algebraic de Rham theory for weakly holomorphic modular forms of level one*, Algebra Number Theory **12** (2018), no. 3, 723–750; arXiv:1708.00652.
- [8] H. Cohen, *Haberland’s formula and numerical computation of Petersson scalar products*, in *ANTS X: Proceedings of the Tenth Algorithmic Number Theory Symposium*, Open Book Ser. **1**, Math. Sci. Publ., 2013, pp. 249–270.
- [9] *Numerical verification scripts* (companion code), 2026. Published checks: `elliptic_period_checks.py`; full working ledger: `wip_meromorphic_periods.py`.
- [10] NIST Digital Library of Mathematical Functions, F. W. J. Olver *et al.*, eds., <https://dlmf.nist.gov/>, Release 1.2.x. Chapters used: §8 (incomplete gamma) and §25.11 (Hurwitz zeta).
- [11] R. Bruggeman, Y. Choie, and N. Diamantis, *Holomorphic automorphic forms and cohomology*, Mem. Amer. Math. Soc. **253** (2018), no. 1212; arXiv:1404.6718.
- [12] K. Bringmann, P. Guerzhoy, Z. Kent and K. Ono, *Eichler-Shimura Theory for Mock Modular Forms*, Mathematische Annalen, March 2013 355(3).
- [13] V. Paşol and A. A. Popa, *Modular forms and period polynomials*, Proc. London Math. Soc. (3) **107** (2013), 713–743; arXiv:1202.5802.
- [14] W. Wang and H. Zhang, *Meromorphic quasi-modular forms and their L-functions*, arXiv:2202.09542 (2022).

- [15] F. Brown and T. J. Fonseca, *Single-valued periods of meromorphic modular forms and a motivic interpretation of the Gross–Zagier conjecture*, arXiv:2508.04844 (2025).
- [16] K. Bringmann, N. Diamantis, and S. Ehlen, *Regularized inner products and errors of modularity*, Int. Math. Res. Not. IMRN (2016); arXiv:1603.03056.
- [17] D. Zagier, *Traces des opérateurs de Hecke*, Séminaire Delange–Pisot–Poitou, 17e année 1975/76, Exp. No. 23, Secrétariat Math., Paris, 1977.
- [18] D. Zagier, *Traces of singular moduli*, in Motives, Polylogarithms and Hodge Theory, Part I (Irvine, CA, 1998), Int. Press Lect. Ser. **3**, International Press, Somerville MA, 2002, pp. 209–244.
- [19] D. Choi and S. Lim, *Group cohomology for modular forms with singularities*, J. Math. Anal. Appl. (2025), PII S0022247X25000526.
- [20] S. Löbrich and M. Schwagenscheidt, *Meromorphic modular forms with rational cycle integrals*, Int. Math. Res. Not. IMRN (2022), no. 1, 312–342; arXiv:1907.04024.
- [21] N. Matthes, *Iterated primitives of meromorphic quasimodular forms for $\mathrm{SL}_2(\mathbb{Z})$* , Trans. Amer. Math. Soc. **375** (2022), 1443–1460; arXiv:2101.11491.
- [22] K. Bringmann, B. Kane, and A.-M. von Pippich, *Regularized inner products of meromorphic modular forms and higher Green’s functions*, Commun. Contemp. Math. **21** (2019), no. 5, 1850029.
- [23] N. Diamantis and J. Drewitt, *Period functions associated to real-analytic modular forms*, Res. Math. Sci. **7** (2020), no. 21.