

L-functions for all meromorphic $\mathrm{SL}_2(\mathbb{Z})$ modular forms, via manifestly finite contours

David A. McGady

Abstract

We give a single manifestly finite, regulator-free functional realizing in one object L-functions quasi-periods and period polynomials of meromorphic modular forms on $\mathrm{SL}_2(\mathbb{Z})$, with generic poles throughout the upper half-plane $\mathbb{H}$ and at the cusp of $\mathrm{SL}_2(\mathbb{Z})$. Additionally, our construction preserves holomorphicity. It is a two-segment contour integral, $L^*(f, s) = L_T^*(f, s; \gamma_{\tau_0}^T) + L_S^*(f, s; \gamma_{\tau_0}^S)$, where L_T^* and L_S^* have distinct integration kernels, and $\gamma_{\tau_0}^S$ is a path in $\mathbb{H}$ from $S(\tau_0)$ to τ_0 , and the same for $\gamma_{\tau_0}^T$. For weakly holomorphic forms our L-function is independent of τ_0 and of the path, and reproduces classical L-functions for holomorphic modular forms, reproduces the L-functions of weak modular forms in [1], which previously needed non-holomorphic regularization to be defined, and reproduces Brown's quasi-periods [2]. For properly meromorphic modular forms with poles inside the interior of $\mathbb{H}$, our L-function depends on the homotopy classes, defined in terms of pole-crossings. There is a natural choice of trivial homotopy class, which renders our L-functions canonical when the modular form has a pole in the interior of the fundamental domain, generalizing [3].

1 Introduction and preliminary concepts

Our primary goal in this note is to put forth a unifying generalization of the L-function associated to a given modular form. Our L-function naturally encompasses the classical case of L-functions associated to holomorphic modular forms [4], and seamlessly unifies Ref. [1]'s extension to L-functions of modular forms with poles at the cusp with Ref. [3]'s extension to L-functions of modular forms with poles at finite locations, exclusively away from cusps. Secondly, we show that our L-functions reproduce the numerical results for quasi-periods associated to weak cusp forms [2].

Note: Throughout this document, we refer to L-integrals and L-functions. The classical L-function-as-Dirichlet-series which has been suitably extended to the complex s plane is simply not defined when the modular form that defines the L-function has poles either at the cusp or at finite locations within the upper half-plane. So, if there is ever a doubt, we state clearly: *any L-function in this note is defined by an L-integral*. The focus of this note is to define L-integrals for any integer weight $\mathrm{SL}_2(\mathbb{Z})$ modular form which may have poles at cusps or away from cusps.

1.1 Modular forms

Write $\mathrm{Re}(z)$ and $\mathrm{Im}(z)$ for the real and imaginary parts of a complex number $z \in \mathbb{C}$. Let $\mathbb{H} := \{\tau \in \mathbb{C} \mid \mathrm{Im}(\tau) > 0\}$ denote the upper half-plane. $\mathrm{SL}_2(\mathbb{Z})$ is the group generated by $T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $S := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

There is a natural $\mathrm{SL}_2(\mathbb{Z})$ action on functions $f : \mathbb{H} \rightarrow \mathbb{C}$, graded by integers k : $(f|_k \gamma)(\tau) := (c\tau + d)^{-k} f(\gamma \cdot \tau)$ for all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$. Similarly, $\gamma \cdot \tau := \frac{a\tau + b}{c\tau + d}$.

In this vein, a *holomorphic modular form* of weight k for $\mathrm{SL}_2(\mathbb{Z})$ is a holomorphic function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that $(f|_k \gamma)(\tau) = f(\tau)$ for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ (aka $f(\tau + 1) = f(-\frac{1}{\tau})\tau^{-k} = f(\tau)$) for all

$\tau \in \mathbb{H}$, and $f(\tau) = \mathcal{O}(1)$ as $\text{Im}(\tau) \rightarrow \infty$, and which has no poles at finite locations within $\mathbb{H}$. By virtue of the periodicity $f(\tau) = f(\tau + 1)$ and holomorphicity, each holomorphic modular form has a q -series decomposition,

$$f(\tau) = \sum_n c_f(n) e^{2\pi i n \tau} = \sum_n c_f(n) q^n \quad , \quad q := e^{2\pi i \tau} \quad . \quad (1)$$

In order for f to be bounded as $\tau \rightarrow i\infty$, we conclude that $c_f(n) = 0$ for all $n < 0$, i.e. that $f = \sum_{n \geq 0} c_f(n) q^n$; for such modular forms f we write $f \in M_k$. If a modular form f additionally has a vanishing constant term, i.e. if $c_f(0) = 0$, then f is a *cusp form*, and we write $f \in S_k$. Clearly, $S_k \subset M_k$. Further, M_k is a finite-dimensional vector-space, as is S_k . Holomorphic $\text{SL}_2(\mathbb{Z})$ modular forms must have even integer weights, and $k \geq 4$ [4].

A product of modular forms of weights k_1 and k_2 is a modular form of weight $k_1 + k_2$, so $M_* := \bigoplus_k M_k$ is naturally a (graded) ring. This graded ring is freely generated by the two modular forms $E_4 \in M_4$ and $E_6 \in M_6$, the $k = 4, 6$ specializations of the *weight k Eisenstein series* E_k , which may be defined as follows:

$$E_k := \frac{1}{2} \sum_{(m,n)=1} \frac{1}{(m+n\tau)^k} = 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \quad , \quad (2)$$

where $\sum_{(m,n)=1}$ denotes summing over all relatively prime $(m, n) \in \mathbb{Z}^2$, the Bernoulli number B_n is the coefficient of $t^n/n!$ of the Laurent expansion of $t/(e^t - 1)$ about $t = 0$, and $\sigma_m(n) := \sum_{d|n} d^m$ is the Divisor function where $d|n$ means $n/d \in \mathbb{N}$. In short, $M_* = \mathbb{C}[E_4, E_6]$. We now define the *modular discriminant*:

$$\Delta := \frac{E_4^3 - E_6^2}{1728} = q + \mathcal{O}(q^2) \in S_{12} \quad . \quad (3)$$

Holomorphicity is most canonically relaxed via allowing a modular form to have a pole at the cusp of $\text{SL}_2(\mathbb{Z})$, i.e. for a weight k modular form for $\text{SL}_2(\mathbb{Z})$ to have negative powers of q in its q -series expansion. In this case, f diverges as $\tau \rightarrow i\infty$ as $f \sim e^{-2\pi N \tau}$, where $-N$ is the largest negative number such that $c_f(n)$ is non-zero. Such a modular form f is called a *weak modular form*, and we write $f \in M_k^!$. If a weak modular form f has a vanishing coefficient of q^0 , i.e. if $c_f(0) = 0$, it is called a *weak cusp form* and we write $f \in S_k^!$. Clearly, $S_k^! \subset M_k^!$, $S_k \subset S_k^!$ and $M_k \subset M_k^!$. Two important weak cusp forms in this note are as follows:

$$J := \frac{E_4^3}{\Delta} - 744 = \frac{1}{q} + \mathcal{O}(q) \in S_0^! \quad , \quad (4)$$

$$\widehat{\Delta} = (J^2 + 24J - 393444)\Delta := \frac{1}{q} + \mathcal{O}(q^2) \in S_{12}^! \quad . \quad (5)$$

In this work, we further relax from even weakly holomorphic modular forms, and consider *meromorphic modular forms* of weight k which are quotients: $f = \frac{f_1}{f_2}$ where $f_i \in M_{k_i}$, the denominator f_2 is not identically zero, and $k = k_1 - k_2$. We write F_k for the vector space of meromorphic modular forms of weight k , and set $F_* := \bigoplus_k F_k$. Then F_k vanishes unless k is an even integer, and F_* may be regarded as a “graded field of fractions” of M_* .

1.2 L-functions for holomorphic modular forms and periods for cusp forms

For any $f \in S_k$, one may define the associated L-function:

$$L(f, s) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s) \stackrel{f \in S_k}{:=} \frac{(2\pi)^s}{\Gamma(s)} \int_0^\infty f(it) t^{s-1} dt = \sum_{n \geq 1} \frac{c_f(n)}{n^s} \quad . \quad (6)$$

A similar definition applies for $f \in M_k$. (The classical definition of $L^*(f, s)$ for $f \in M_k$ is the focus of Remark 3.6, below.) Similarly, one may define a period polynomial for $f \in S_k$ in terms of this L-function,

$$r_f(X, Y) := (2\pi i)^{k-1} \sum_{\ell=0}^{k-2} (-1)^\ell \binom{k-2}{\ell} i^{\ell+1} L^*(f, \ell+1) X^{k-2-\ell} Y^\ell. \quad (7)$$

This polynomial, $r_f(X, Y) \in V_{k-2} := \mathbb{C}[X, Y]_{k-2}$, has a natural action under $\mathrm{SL}_2(\mathbb{Z})$, $(P|_\gamma)(X, Y) = P(aX + bY, cX + dY)$ for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$. This period polynomial can be decomposed into a basis of highly constrained polynomials, which are annihilated by $|(1+T)$ and $|(1+U+U^2)[6]$, where $U := TS$ generates an order-3 cyclic subgroup of $\mathrm{SL}_2(\mathbb{Z})$. We define the *periods* of a cusp form to be coefficients of r_f within this highly constrained sub-space W of V_{k-2} , i.e. $W \subset V_{k-2}$.

Why this work? As classically computed, these L-functions and associated period polynomials (when defined in terms of the special values of the associated L-function) are not defined when f has poles either at the cusp of $\mathrm{SL}_2(\mathbb{Z})$ or within the interior of $\mathbb{H}$. As stated above, there are extant, but partial and non-overlapping, extensions of the definition of $L(f, s)$ to $f \in M_k^!$ [1] or $f \in F_k$ when f is regular at the cusp of $\mathrm{SL}_2(\mathbb{Z})$ [3]. In this note, we provide a construction that unifies these disparate extensions. We start with generic L-integrals in section 2, then specialize to $f \in M_k^!$ in section 3, and finally explore the more general context of $f \in F_k$ in section 4.

2 A finite L-integral

Definition 2.1. Let $\tau_0 \in \mathbb{H}$ and $f \in F_k$, let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ denote the generators of $\Gamma = \mathrm{SL}_2(\mathbb{Z})$, and let $f^{-1}(\infty)$ be the locus of all poles of $f \in \mathbb{H}$ (inclusive of the cusp of $\mathrm{SL}_2(\mathbb{Z})$). The S - and T -segments at τ_0 are smooth paths in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$ from $\sigma^{-1}\tau_0$ to τ_0 ,

$$\gamma_{\tau_0}^S : -1/\tau_0 \longrightarrow \tau_0, \quad \gamma_{\tau_0}^T : \tau_0 - 1 \longrightarrow \tau_0. \quad (8)$$

Avoiding poles is automatic for $f \in M_k^!$, as long as the base-point τ_0 , is not at a cusp of $\mathrm{SL}_2(\mathbb{Z})$, i.e. there is only the trivial homotopy class. When $f \in F_k$, and f has poles at finite and non-zero imaginary heights, integration contours fall into nontrivial homotopy classes.

Definition 2.2. Let $\zeta(s, x) = \sum_{m \geq 0} (m+x)^{-s}$ as a function $\zeta : \mathbb{C}_s \times (\mathbb{C} \setminus \mathbb{N})_x \rightarrow \mathbb{C}$ denote the Hurwitz zeta function (meromorphic in s , holomorphic in $x \in \mathbb{H}$). We define Hurwitz T -kernel as

$$\tilde{k}_T(\tau, s) := \zeta(1-s, \tau+1) - e^{i\pi(s-1)} \zeta(1-(k-s), \tau+1). \quad (9)$$

Definition 2.3 (Generic L-integral). Let $f \in F_k$, $\tau_0 \in \mathbb{H}$, and $s \in \mathbb{C}$. Further, let $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$ be the S - and T -segments at τ_0 , as in Definition 2.1, and let k_T be as defined in Definition 2.2. The L-integral of f is

$$L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) := L_S^*(f, s; \gamma_{\tau_0}^S) + L_T^*(f, s; \gamma_{\tau_0}^T) := e^{-\pi i s/2} \left(\int_{\gamma_{\tau_0}^S} f(\tau) \tau^{s-1} d\tau + \int_{\gamma_{\tau_0}^T} f(\tau) \tilde{k}_T(\tau, s) d\tau \right) \quad (10)$$

We use this L -integral to define a generalized L -function:

Definition 2.4 (Generic L-function). Let $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ be as defined in Definition 2.3. We define the generalized L-function of $f \in F_k$ to be,

$$L(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T). \quad (11)$$

Having defined the L-integral and L-function, we prove they are manifestly finite for $f \in F_k$:

Lemma 2.5 (Finiteness of L-integrals). *For $f \in F_k$, both $L_S^*(f, s; \gamma_{\tau_0}^S)$ and $L_T^*(f, s; \gamma_{\tau_0}^T)$ in Definition 2.3 are each, separately finite for every $s \in \mathbb{C}$, every $\tau_0 \in \mathbb{H}$, and for every path $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$.*

Proof. Both segments are compact and lie in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, so f is continuous on them; τ^{s-1} is continuous on $\mathbb{H}$, and $\tilde{k}_T(\tau, s)$ is continuous there since $\zeta(\nu, \tau + 1)$ is holomorphic in $\tau \in \mathbb{H}$ (Definition 2.2). A continuous integrand on a compact path is integrable, for every $s \in \mathbb{C}$. $\square$

We now turn to path-dependence and τ_0 -independence of the L-integral and the L-function of $f \in F_k$. This holds for all $f \in M_k^!$:

Proposition 2.6 (Basepoint & path independence for $f \in M_k^!$). *For $f \in M_k^!$ the value $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ is independent of $\tau_0 \in \mathbb{H}$ and of the paths: with no interior poles, $\mathbb{H}$ is simply connected, so all admissible contours are homotopic. We write $L^*(f, s)$ for the common value and $L(f, s) := \frac{(2\pi)^s}{\Gamma(s)} L^*(f, s)$.*

Proof. With no interior poles the integrands $f(\tau)\tau^{s-1}$ and $f(\tau)\tilde{k}_T(\tau, s)$ are holomorphic on $\mathbb{H}$, which is simply connected; by Cauchy's theorem each integral is unchanged under any deformation fixing its endpoints, hence depends on those endpoints alone. This is path-independence.

For independence of τ_0 show that variations in τ_0 for the S-term and T-term in $L^*(f, s)$ cancel, i.e. that $\partial_{\tau_0} L_S^*(f, s; \gamma_{\tau_0}^S) = -\partial_{\tau_0} L_T^*(f, s; \gamma_{\tau_0}^T)$. For L_T^* , note that the T -segment runs from $T^{-1}\tau_0 = \tau_0 - 1$ to τ_0 — two endpoints a unit apart, moving together with τ_0 — so by Leibniz's rule and the T -periodicity $f(\tau_0 - 1) = f(\tau_0)$,

$$\partial_{\tau_0} L_T^*(f, s; \gamma_{\tau_0}^T) = f(\tau_0) \tilde{k}_T(\tau_0, s) - f(\tau_0 - 1) \tilde{k}_T(\tau_0 - 1, s) = f(\tau_0) \delta_T \tilde{k}_T(\tau_0 - 1, s), \quad (12)$$

where $\delta_T \tilde{k}_T(\tau, s) := \tilde{k}_T(\tau + 1, s) - \tilde{k}_T(\tau, s)$. This is the whole role of δ_T : the unit gap between the endpoints turns the ∂_{τ_0} boundary term into a unit shift of the kernel. That shift is elementary — the kernel was built for it — since $\zeta(\nu, x + 1) - \zeta(\nu, x) = -x^{-\nu}$ gives, on both terms at $x = \tau + 1$,

$$\delta_T \tilde{k}_T(\tau, s) = -(\tau + 1)^{s-1} + e^{i\pi(s-1)}(\tau + 1)^{k-1-s} \quad (13)$$

$$\implies \partial_{\tau_0} L_T^*(f, s; \gamma_{\tau_0}^T) = f(\tau_0) [-\tau_0^{s-1} + e^{i\pi(s-1)}\tau_0^{k-1-s}]. \quad (14)$$

The S -segment runs $-1/\tau_0 \rightarrow \tau_0$; its lower endpoint moves with $\partial_{\tau_0}(-1/\tau_0) = \tau_0^{-2}$, so by Leibniz, the modularity $f(-1/\tau_0) = \tau_0^k f(\tau_0)$, and the phase $(-1/\tau_0)^{s-1} = e^{i\pi(s-1)}\tau_0^{1-s}$,

$$\partial_{\tau_0} L_S^*(f, s; \gamma_{\tau_0}^S) = f(\tau_0)\tau_0^{s-1} - f(-1/\tau_0)(-1/\tau_0)^{s-1}\tau_0^{-2} = f(\tau_0)[\tau_0^{s-1} - e^{i\pi(s-1)}\tau_0^{k-1-s}]. \quad (15)$$

Thus τ_0 -variations are equal and opposite, $\partial_{\tau_0} L_S^* = -\partial_{\tau_0} L_T^*$. Thus $\partial_{\tau_0} L^* = \partial_{\tau_0} L_S^* + \partial_{\tau_0} L_T^* = 0$. $\square$

Remark 2.7 (Recovering the classical L-integral for $f \in S_k$). *Let $f \in S_k$. As $f^{-1}(\infty)$ does not include the cusp $i\infty$, $\lim_{\tau_0 \rightarrow i\infty} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ exists. In this limit, the S -segment maps onto the standard L -function contour, $[0, i\infty]$. Simultaneously, as $f(it) \sim e^{-2\pi t}$ as $t \rightarrow \infty$, the T -segment vanishes exponentially. Thus, in this limit, $L^*(f, s)$ exactly maps onto the classical L -function integral [4]:*

$$\lim_{\tau_0 \rightarrow i\infty} L^*(f, s) = \int_0^\infty t^{s-1} f(it) dt. \quad (16)$$

3 L-functions for weak modular forms

Proposition 3.1. Let $\Gamma(s, x) := \int_x^\infty e^{-t} t^{s-1} dt$ denote the incomplete Gamma function. For $n \neq 0$, $\nu \in \mathbb{C}$, and $z \in \mathbb{H}$,

$$\int_z^{z+1} e^{2\pi i n u} \zeta(\nu, u) du = e^{i\pi(1-\nu)/2} (2\pi n)^{\nu-1} \Gamma(1-\nu, -2\pi i n z). \quad (17)$$

Proof. Write $\zeta(\nu, u) = \sum_{m \geq 0} (m+u)^{-\nu}$ and shift $u \mapsto u+m$ in the m -th term; as $e^{2\pi i n(u+m)} = e^{2\pi i n u}$ for integer m , the unit segments stack into one horizontal ray,

$$\int_z^{z+1} e^{2\pi i n u} \zeta(\nu, u) du = \sum_{m \geq 0} \int_{z+m}^{z+1+m} e^{2\pi i n v} v^{-\nu} dv = \int_z^{z+\infty} e^{2\pi i n v} v^{-\nu} dv. \quad (18)$$

Substitute $w = -2\pi i n v$, so that $v = iw/(2\pi n)$, $dv = i dw/(2\pi n)$, and $e^{2\pi i n v} = e^{-w}$: the lower endpoint becomes $w = -2\pi i n z$, and for $n > 0$ the ray rotates to $w \rightarrow +\infty$ along the positive reals — legal, since $e^{-w} w^{-\nu}$ decays in the right half-plane, so the closing arc vanishes. With $v^{-\nu} = i^{-\nu} (2\pi n)^{\nu} w^{-\nu}$,

$$\int_z^{z+\infty} e^{2\pi i n v} v^{-\nu} dv = i^{1-\nu} (2\pi n)^{\nu-1} \int_{-2\pi i n z}^{\infty} e^{-w} w^{-\nu} dw = e^{i\pi(1-\nu)/2} (2\pi n)^{\nu-1} \Gamma(1-\nu, -2\pi i n z). \quad (19)$$

For $n < 0$ the ray instead grows as $\text{Im} v \rightarrow \infty$ and the rotation is illegal; but both sides are analytic in n and agree for $n > 0$, so the identity continues to $n < 0$ with Γ on its principal branch. At $n = 0$ the prefactor $(2\pi n)^{\nu-1}$ is singular; that case is handled separately. $\square$

Theorem 3.2 (The L-function for $f \in M_k^!$). For $f \in M_k^!$, we may set $\tau_0 = i$, and prove

$$L^*(f, s) = -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right). \quad (20)$$

Here, trivially, $L^*(f, s) = i^k L^*(f, k-s)$. The L-function is $L(f, s) := (2\pi)^s L^*(f, s)/\Gamma(s)$.

Proof. By path independence and base-point independence from Proposition 2.6, we may freely set $\tau_0 = i$. Here, $L_S^*(f, s; \gamma_{\tau_0}^S) = 0$ (the S -contour vanishes in length and there are no nontrivial homotopy classes as $f \in M_k^!$ has no poles which a finite contour can wrap), and the L-integral simplifies dramatically:

$$L^*(f, s) = L_T^*(f, s; \gamma_i^T) = e^{-\pi i s/2} \int_{i-1}^i f(\tau) \tilde{k}_T(\tau, s) d\tau. \quad (21)$$

We then expand f in terms of its q-series, $\sum_n c_f(n) e^{2\pi i n \tau}$, and substitute the u -variable of Proposition 3.1 as $u = \tau + 1$, and note that $e^{2\pi i n(u-1)} = e^{2\pi i n u}$. When $n \neq 0$, straightforward substitutions of ν, z into Proposition 3.1 suffice: $1-\nu = s$ gives the contribution of the first term in $\tilde{k}_T(\tau, s)$, and $1-\nu = k-s$ gives the second:

$$e^{-\pi i s/2} \left(e^{\pi i s/2} \frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} - e^{i\pi(s-1)} e^{\pi i(k-s)/2} \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right) = \frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}}. \quad (22)$$

We may then evaluate the $n = 0$ mode's contribution follows from noting the closed form solution for the definite integral,

$$\int_{i-1}^i \zeta(\sigma, \tau + 1) d\tau = \int_i^{i+1} \zeta(\sigma, u) du = \frac{i^{1-\sigma}}{\sigma-1}. \quad (23)$$

This result follows from noting $\partial_a(\zeta(\sigma-1, a)/(1-\sigma)) = \zeta(\sigma, a)$ and $\zeta(\sigma-1, a+1) - \zeta(\sigma-1, a) = -a^{1-\sigma}$. Applying this to the two distinct terms of $\tilde{k}_T(\tau, s)$ at $\sigma = 1-s$ and $\sigma = 1-(k-s)$ gives

$$e^{-\pi is/2} \int_{i-1}^i c_f(0) \tilde{k}_T(\tau, s) d\tau = -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right). \quad (24)$$

Combining the $n=0$ contributions with the $n \neq 0$ contributions reproduces Eq. (20). Finally, the functional equation, $L^*(f, s) = i^k L^*(f, k-s)$, follows by direct inspection. $\square$

3.1 Explicit S- and T-contours for general τ_0

Lemma 3.3 (L_T^* for $M_k^!$). For $f \in M_k^!$, and $\tau_0 \in \mathbb{H}$ away from cusps, $L_T^*(f, s; \gamma_{\tau_0}^T)$ evaluates to:

$$L_T^*(f, s; \gamma_{\tau_0}^T) = -c_f(0) \left(\frac{(\tau_0/i)^s}{s} + i^k \frac{(\tau_0/i)^{k-s}}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, -2\pi in\tau_0)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, -2\pi in\tau_0)}{(2\pi n)^{k-s}} \right). \quad (25)$$

Proof. Expand $f = \sum_n c_f(n) q^n$ on the compact segment $[\tau_0-1, \tau_0]$, and restrict attention to the sub-sum, $\sum_{n \neq 0}$. For each such $n \neq 0$, apply the Proposition 3.1 in $\tilde{k}_T(\tau, s) = \zeta(1-s, \tau+1) - e^{i\pi(s-1)} \zeta(s-(k-1), \tau+1)$ at $\nu = 1-s$ and $\nu = s-(k-1)$. The phases for the second term work-out identically as in the proof of Theorem 3.2.

For $n=0$, the constant $c_f(0)$ multiplies $e^{-\pi is/2} \int_{\tau_0-1}^{\tau_0} \tilde{k}_T(\tau, s) d\tau$; the Hurwitz antiderivative $\int_{\tau_0-1}^{\tau_0} \zeta(\sigma, \tau+1) d\tau = \tau_0^{1-\sigma}/(\sigma-1) -$ from $\partial_\tau \zeta(\sigma-1, \tau+1) = -(\sigma-1)\zeta(\sigma, \tau+1)$ and the shift $\zeta(\sigma-1, \tau_0+1) - \zeta(\sigma-1, \tau_0) = -\tau_0^{1-\sigma}$ — applied to the two halves of $\tilde{k}_T$ at $\sigma = 1-s$ and $\sigma = s-k+1$ yields $-e^{-\pi is/2} \tau_0^s/s = -(\tau_0/i)^s/s$ and $-e^{\pi is/2} \tau_0^{k-s}/(k-s) = -i^k (\tau_0/i)^{k-s}/(k-s)$ respectively. $\square$

Corollary 3.4 (L_T^* as $\tau_0 \rightarrow i\infty$). Let $f \in S_k^!$, where $f = \sum_{n \geq -N} c_f(n) q^n$ for some $-N \in \mathbb{Z}$. The T -contour's contribution to $L^*(f, s)$ is given by Lemma 3.3. As $\tau_0 = it$ goes to the cusp $i\infty$, the T -contour scales exactly as $f(it)$ scales: $L_T^*(f, s; \gamma_{\tau_0}^T) \sim f(it) \sim e^{2\pi Nt}$.

Proof. For the asymptotics, $\Gamma(\sigma, z) \sim z^{\sigma-1} e^{-z}$ as $|z| \rightarrow \infty$ for $|\arg z| < \frac{3\pi}{2}$ [10, §8.11] gives $|c_f(n)|$ -weighted go as $e^{-2\pi n \text{Im}(\tau_0)}$. The smallest mode n in the q -series, i.e. $n = -N$, dominates the scaling of $f(it)$ and of $L_T^*(f, s; \gamma_{it}^T)$. Combining these results yields the claimed asymptotics. $\square$

Now, exploiting τ_0 -independence of $L^*(f, s)$, its explicit value as given in Theorem 3.2, and the explicit T -contribution $L_T^*(f, s; \gamma_{\tau_0}^T)$ trivially implies clean, closed-form expressions for $L_S^*(f, s; \gamma_{\tau_0}^S)$:

Corollary 3.5 (L_S^* for $M_k^!$). For $f \in M_k^!$ and $\tau_0 \in \mathbb{H}$, $L_S^*(f, s; \gamma_{\tau_0}^S)$ evaluates to

$$L_S^*(f, s; \gamma_{\tau_0}^S) = c_f(0) \left(\frac{(\tau_0/i)^s - 1}{s} + i^k \frac{(\tau_0/i)^{k-s} - 1}{k-s} \right) + \sum_{n \neq 0} c_f(n) \left(\frac{\Gamma(s, 2\pi n) - \Gamma(s, -2\pi in\tau_0)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n) - \Gamma(k-s, -2\pi in\tau_0)}{(2\pi n)^{k-s}} \right). \quad (26)$$

Proof. Recall $L^*(f, s) - L_T^*(f, s; \gamma_{\tau_0}^T) = L_S^*(f, s; \gamma_{\tau_0}^S)$. Now, $L^*(f, s)$ for $f \in M_k^!$ is given by Theorem 3.2, and $L_T^*(f, s; \gamma_{\tau_0}^T)$ is given by Lemma 3.3. Their difference yields Eq. (26). $\square$

3.2 Classical Mellin recovery and canonical uniqueness.

As discussed in Remark 2.7, for $f \in S_k$, taking the cusp-limit $\tau_0 \rightarrow i\infty$ in Theorem 3.2 collapses the S -segment onto the classical Mellin contour $[0, i\infty]$ while via Corollary 3.4 the T -segment vanishes exponentially as $L_T^*(f, s; \gamma_{\tau_0}^T) \sim f(\tau_0) \sim e^{-2\pi\text{Im}(\tau_0)N}$ (as $f \in S_k \Leftrightarrow f = \sum_{n \geq N} c_f(n)q^n$ with $N \geq 1$). I.e., here, one recovers the classical L-integral of f without obstruction:

$$\lim_{\tau_0 \rightarrow i\infty} L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) = \int_0^\infty t^{s-1} f(it) dt. \quad (27)$$

This story changes when $f = 1/q^N(1 + O(q)) \in S_k^!$ for some $N \geq 1$.

In short, when f has a polar q -series, it has a pole at the cusp of $\text{SL}_2(\mathbb{Z})$. Thus, the limit $\tau_0 \rightarrow i\infty$, which was crucial to recovering the classical Mellin contour in Remark 2.7 simply does not exist.

However, it can still be instructive to examine $L_T^*(f, s; \gamma_{\tau_0}^T)$, and $L_S^*(f, s; \gamma_{\tau_0}^S)$ as τ_0 goes towards $i\infty$. In this context, while of course the sum $L_T^*(f, s; \gamma_{\tau_0}^T) + L_S^*(f, s; \gamma_{\tau_0}^S) = L^*(f, s)$ is manifestly invariant under changes of the base-point, each of these two terms can indeed change values. Indeed, Corollary 3.4 clearly shows that as $\tau_0 \rightarrow i\infty$, when $f \in S_k^!$ has negative powers of q in its q -series, then $L_T^*(f, s; \gamma_{\tau_0}^T) \sim e^{2\pi N \text{Im}(\tau_0)}$. Discarding $\gamma_{\tau_0}^T$ therefore does violence to the result: When the S -contour is isolated from its T -partner, and $\tau_0 \rightarrow i\infty$, it diverges. This is the precise reason that [1] needed to regularize their divergent L-integral representation for properly weak cusp forms.

That said, there is one further aspect of our L-function even in the classical case of $f \in M_k$ that is worthy of remark:

Remark 3.6 (Recovering the classical L-integral for $f \in M_k$). *In textbook treatments [4], when $f \in M_k$, its L-integral is defined as*

$$\begin{aligned} L^*(f, s)|_{\text{cl}} &:= \int_0^\infty (f(it) - c_f(0))t^{s-1}dt = \frac{\Gamma(s)}{(2\pi)^s} \sum_{n=1}^\infty \frac{c_f(n)}{n^s} \\ &= -c_f(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n=1}^\infty c_f(n) \left(\frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}} \right) \end{aligned} \quad (28)$$

Classically, this constant-term subtraction is needed because $\int_0^\infty t^{s-1}dt$ is a pure divergence, so the q -series cannot commute with the integration contour.

However, as is clear from our general Theorem 3.2 for L-integrals for $f \in M_k^!$, and from the specific results of the S - and T -contours in Lemma 3.3 and Corollary 3.5, our L-integral requires no such explicit subtraction of the constant term $-c_f(0)$. Thus, not only does our prescription naturally handle divergences at poles when $f \in M_k^!$ has polar terms in its q -series, but it also obviates the need to by-hand remove the constant mode from the kernel of the L-integral that is needed when defining L-functions in terms of the classical Mellin contour, $\int_0^\infty$.

3.3 Period polynomials, periods and quasi-periods

Because $L^*(f, s)$ is canonical for every $f \in S_k^!$, and because period polynomials for a given modular form can be written in terms of their L-functions evaluated at integer- s in the critical strip, they need no separate construction.

Definition 3.7 (Period polynomial for $S_k^!$). *For $f \in S_k^!$ and $n = k - 2$, the period polynomial is*

$$r_f(X, Y) := (2\pi i)^{n+1} \sum_{\ell=0}^n (-1)^\ell \binom{n}{\ell} i^{\ell+1} L^*(f, \ell+1) X^{n-\ell} Y^\ell. \quad (29)$$

Proposition 3.8 (Period polynomial as a contour integral). *Let $n := k - 2$. With $(X - \tau Y)^n$ in place of the L -weight τ^{s-1} , and $\mathbf{K}_T(\tau; X, Y) := \sum_{\ell=0}^n \binom{n}{\ell} (-1)^\ell \tilde{k}_T(\tau, \ell + 1) X^{n-\ell} Y^\ell$ in place of $\tilde{k}_T$, the same two segments assemble r_f directly:*

$$r_f(X, Y) = (2\pi i)^{n+1} \left[\int_{\gamma_{\tau_0}^S} f(\tau) (X - \tau Y)^n d\tau + \int_{\gamma_{\tau_0}^T} f(\tau) \mathbf{K}_T(\tau; X, Y) d\tau \right]. \quad (30)$$

Proof. By Definition 2.3, the $X^{n-\ell} Y^\ell$ coefficient of the bracket is

$$r_f|_{X^{n-\ell} Y^\ell} = (-1)^\ell \binom{n}{\ell} i^{\ell+1} [L_S^*(f, \ell + 1) + L_T^*(f, \ell + 1)] = (-1)^\ell \binom{n}{\ell} i^{\ell+1} L^*(f, \ell + 1). \quad (31)$$

This matches Definition 3.7 term by term, and proves the claim. $\square$

Having now defined the period polynomial associated to a weak cusp form $f \in S_k^!$, we now seek to find its (*quasi*-)periods. First, a number of helpful definitions. For starters, let $f \in S_k^!$, $\tau_0 \in \mathbb{H}$ and $\gamma \in \mathrm{SL}_2(\mathbb{Z})$, and define

$$C_\gamma(f; \tau_0) := (2\pi i)^{n+1} \int_{\gamma^{-1}\tau_0}^{\tau_0} \omega_f, \quad \text{where } \omega_f := f(\tau)(X - \tau Y)^n d\tau. \quad (32)$$

Finally, it is helpful to define two slight abstractions related to $\mathbf{K}_T(\tau; X, Y)$ and to $L_T^*(f, s; \gamma_{\tau_0}^T)$:

$$\chi(\tau; X, Y) := \sum_{\ell=0}^n \binom{n}{\ell} (-1)^\ell \zeta(-\ell, \tau + 1) X^{n-\ell} Y^\ell, \quad (33)$$

$$E(f; \tau_0) := (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) \chi(\tau; X, Y) d\tau. \quad (34)$$

For each τ , $\chi(\tau; \cdot, \cdot) \in V_{k-2} = \mathbb{C}[X, Y]_{k-2}$; the $\mathrm{SL}_2(\mathbb{Z})$ -slash $(P|_\gamma)(X, Y) := P(aX + bY, cX + dY)$ acts on the polynomial variables alone, holding τ fixed:

$$\chi|_\gamma(\tau; X, Y) := \chi(\tau; aX + bY, cX + dY), \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}). \quad (35)$$

Thus $|_T$ sends $(X, Y) \mapsto (X + Y, Y)$ with τ untouched. Being linear and τ -independent, $|_\gamma$ commutes with $\int d\tau$, so slashing the polynomial E agrees with slashing χ under the integral — the two occurrences of $|_\gamma$ in Lemma 3.9 denote the same operation.

Lemma 3.9. *Let $f \in S_k^!$, $\tau_0 \in \mathbb{H}$, and let ω_f , $C_\gamma(f; \tau_0)$, $\chi(\tau; X, Y)$, and $E(f; \tau_0)$ be as defined in Eqs. (32), (33) and (34). Then $\hat{C}_\gamma := C_\gamma - (E|_\gamma - E)$ is a 1-cocycle for $\mathrm{SL}_2(\mathbb{Z})$ with $\hat{C}_S = r_f$ and $\hat{C}_T = 0$.*

Proof. First, we prove $\hat{C}_\gamma$ is a cocycle. Path-additivity and covariance of ω_f give the Eichler–Shimura cocycle relation $C_{\gamma\delta} = C_\gamma|_\delta + C_\delta$ [11, 6]; as $\gamma \mapsto E|_\gamma - E$ is a coboundary, $\hat{C}$ is a cocycle.

Having established it is a cocycle, we now show that $\hat{C}_S = r_f$. At $s = \ell + 1$, Definition 2.2 reads $\tilde{k}_T(\tau, \ell + 1) = \zeta(-\ell, \tau + 1) - (-1)^\ell \zeta(-(n - \ell), \tau + 1)$; assembling $\mathbf{K}_T$ and re-indexing $\ell \mapsto n - \ell$ in the second term (with n even) gives $\mathbf{K}_T = \chi - \chi|_S$. So the T -term of r_f is $E - E|_S$, and by Proposition 3.8 $r_f = C_S + (E - E|_S) = C_S - (E|_S - E) = \hat{C}_S$.

Finally, we show that $\hat{C}_T = 0$. The Hurwitz shift $\zeta(-\ell, \tau) = \zeta(-\ell, \tau + 1) + \tau^\ell$ gives the point-wise identity $\chi(\tau - 1; X, Y) - \chi(\tau; X, Y) = (X - \tau Y)^n$; applied twice it shows $A(\tau) :=$

$\chi(\tau; X+Y, Y) - \chi(\tau-1; X, Y)$ satisfies $A(\tau-1) = A(\tau)$. Crucially, because $A(\tau)$ is a polynomial, and because $A(\tau) = A(\tau+1)$ for generic τ , then $A(\tau)$ is constant: $\partial_\tau A(\tau) = 0 \Leftrightarrow A(\tau) = A_0$. Thus,

$$\begin{aligned}
\hat{C}_T &= C_T - (E|_T - E) \\
&= (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau)(X - \tau Y)^n d\tau - (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) [\chi|_T(\tau; X, Y) - \chi(\tau; X, Y)] d\tau \\
&= (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) \left([\chi(\tau-1; X, Y) - \chi(\tau; X, Y)] - [\chi(\tau; X+Y, Y) - \chi(\tau; X, Y)] \right) d\tau \\
&= (2\pi i)^{n+1} \int_{\gamma_{\tau_0}^T} f(\tau) [\chi(\tau-1; X, Y) - \chi(\tau; X+Y, Y)] d\tau = -(2\pi i)^{n+1} \int_{\tau_0-1}^{\tau_0} f(\tau) A(\tau) d\tau \\
&= -(2\pi i)^{n+1} A_0 \int_{\tau_0-1}^{\tau_0} f(\tau) d\tau = -(2\pi i)^{n+1} A_0 c_f(0) = 0,
\end{aligned} \tag{36}$$

as all elements $f \in S_k^!$ must have $c_f(0) = 0$. $\square$

Definition 3.10. $W := \{w \in \mathbb{C}[X, Y]_{k-2} : w|_{(1+S)} = w|_{(1+U+U^2)} = 0\} \subset V_n$

Lemma 3.11. For $f \in S_k^!$, and $W \subset V_n$ from Definition 3.10, the period polynomial $r_f(X, Y) \in W$.

Proof. Let $\hat{C}_\gamma$ be the parabolic cocycle of Lemma 3.9. Thus, $r_f = \hat{C}_S$ and $\hat{C}_T = 0$. Now, note that $\hat{C}_\gamma(f; \tau_0) = 0$ when $\gamma = \pm I$ for the simple reason that when $\gamma = \pm I$, $\hat{C}_\gamma$ is a closed-contour integral of a function which has no poles in $\mathbb{H}$. Finally, direct computation, starting from the identities $-I = U^3 = S^2$ and using the cocycle law $\hat{C}_{\gamma\delta} = \hat{C}_\gamma|_\delta + \hat{C}_\delta$ and clever use of $\hat{C}_T = 0$, shows $r_f|_{(1+S)} = r_f|_{(1+U+U^2)} = 0$:

$$\hat{C}_U = \hat{C}_{TS} = \hat{C}_T|_S + \hat{C}_S = \hat{C}_S = r_f, \tag{37}$$

$$0 = \hat{C}_{S^2} = \hat{C}_S|_S + \hat{C}_S = \hat{C}_S|_{(1+S)} = r_f|_{(1+S)}, \tag{38}$$

$$0 = \hat{C}_{U^3} = \hat{C}_U|_{UU} + \hat{C}_{UU} = \hat{C}_U|_{U^2} + (\hat{C}_U|_U + \hat{C}_U) = \hat{C}_U|_{(1+U+U^2)} = r_f|_{(1+U+U^2)}. \tag{39}$$

Thus, $r_f \in W \subset V_n$, and finishes the proof. $\square$

To define periods and quasi-periods, we recall the standard fact that $\dim(W) = 2 \dim(S_k) + 1$ (see, e.g. [6]), and define the inner-product:

$$\langle P, Q \rangle := \sum_a (-1)^a \binom{n}{a}^{-1} p_a q_{n-a}. \tag{40}$$

Now for the basis elements for k where $\dim(S_k) = 1$:

Definition 3.12. Let $\dim S_k = 1$, $n := k-2$, $p_0(X) := X^n - 1$, $B_\ell := \frac{\ell!}{2\pi i} \oint \frac{t^{-\ell}}{e^t - 1} dt$, and $B_m^0(X) := \sum_{i \neq 1} \binom{m}{i} B_i X^{m-i}$. Motivated by [6, Thm. 1'], we define:

$$W_-^{(k)}(X) := \text{pp} \left(\frac{1}{n-1} (B_{n-1}^0(X) + X^n B_{n-1}^0(\frac{1}{X})) - \frac{1}{3} (B_3^0(X) + X^n B_3^0(\frac{1}{X})) \right), \tag{41}$$

$$W_+^{(k)}(X) := \text{pp} \left(\frac{1}{2} (B_2^0(X) - X^n B_2^0(\frac{1}{X})) + \frac{1}{n} (B_n^0(X) - X^n B_n^0(\frac{1}{X})) - c p_0(X) \right), \tag{42}$$

$$v_0^{(k)}(X) := p_0(X) - W_+^{(k)}(X) \left(\langle W_+^{(k)}, p_0 \rangle / \langle W_+^{(k)}, W_+^{(k)} \rangle \right), \tag{43}$$

where pp acts on any polynomial with rational coefficients by multiplying by all coefficient-denominators and dividing by the gcd, c eliminates the $p_0(X)$ component of $W_+^{(k)}$, and $\langle P, Q \rangle$ as in Eq. (40).

Lemma 3.13. *For k with $\dim S_k = 1$, the space W of Lemma 3.11 has $\dim W = 3$, and $\{W_+^{(k)}, W_-^{(k)}, v_0^{(k)}\}$ is an orthogonal basis of W under $\langle \cdot, \cdot \rangle$ from Eq. (40).*

Proof. First, let $W_\pm^{(k)}(X, Y) := Y^{k-2}W_\pm^{(k)}(X/Y)$, and let $v_0^{(k)}(X, Y) := Y^{k-2}v_0^{(k)}(X/Y)$.

Now, by Eichler–Shimura, $\dim W = 2 \dim S_k + 1 = 3$. Theorem 1' of [6] gives the odd and even period polynomials of the cusp form $R_n \propto \Delta$ as modified-Bernoulli combinations; its $n = 2$ case is $r^-(R_2)$, odd under $X \mapsto -X$, and its $n = 1$ case is $r^+(R_1)$, even. Our $W_-^{(k)}$ is the primitive part of $r^-(R_2)$, and $W_+^{(k)}$ the primitive part of $r^+(R_1)$ with its Eisenstein p_0 -component removed by c ; both lie in W , as does $p_0 = X^{k-2} - Y^{k-2}$ (even). The three are independent, hence a basis of W . Since $k - 2$ is even, $\langle \cdot, \cdot \rangle$ is symmetric and pairs opposite $X \mapsto -X$ parities to zero; the odd $W_-^{(k)}$ is thus orthogonal to the even $W_+^{(k)}$ and $v_0^{(k)}$, while $v_0^{(k)} \perp W_+^{(k)}$ by construction. Hence $\{W_+^{(k)}, W_-^{(k)}, v_0^{(k)}\}$ is orthogonal. $\square$

Definition 3.14 (Periods and Quasi-Periods). *Let k be such that $\dim(S_k) = 1$, $n := k - 2$, $W_\pm^{(k)}(X, Y) := Y^n W_\pm^{(k)}(X/Y)$ be as in Definition 3.12, and $r_f(X, Y)$ be as in Definition 3.7. Now, define the two numbers $\omega^\pm(f)$ as follows:*

$$\omega^\pm(f) := \frac{\langle r_f, W_\pm^{(k)} \rangle}{\langle W_\pm^{(k)}, W_\pm^{(k)} \rangle}, \quad (44)$$

where $\langle \cdot, \cdot \rangle$ is as in Eq. (40). When f is a cusp form, $\omega^\pm(f)$ are its periods; when f is a weak cusp form, $\omega^\pm(f)$ are its quasi-periods, sometimes denoted $\eta^\pm(f)$.

Now, having defined our periods and quasi-periods, we use them together with our L-functions to extract periods and quasi-periods for $f \in S_k^!$ (when $\dim(S_k) = 1$). To this effect, we recall the notation of [5]. Decomposing the modular weight $k = 12\lambda + k'$ where $k' \in \{0, 4, 6, 8, 10, 14\}$, we see $\lambda = \dim(S_k)$ when $\lambda \geq 0$. For any λ , we may denote $f_{k,m} := 1/q^m + O(q^{\lambda+1}) \in M_k^!$. When $\dim(S_k) = 1$, we have $f_{k,m} = 1/q^m + O(q^2)$. Here, define $\Delta_k := f_{k,-1} = q + O(q^2) \in S_k$ and $\hat{\Delta}_k := f_{k,1} = 1/q + O(q^2) \in S_k^!$. Further, define

$$\text{pd}_k := \begin{pmatrix} \omega^+(\Delta_k) & \omega^-(\Delta_k) \\ \eta^+(\hat{\Delta}_k) & \eta^-(\hat{\Delta}_k) \end{pmatrix}, \quad \alpha_k := \frac{(-1)^{1+k/2} \det(\text{pd}_k)}{(2\pi i)^{k-1} (k-2)!} \quad (45)$$

k	$\omega^+(\Delta_k)$	$\omega^-(\Delta_k)/i$	$\eta^+(\hat{\Delta}_k)$	$\eta^-(\hat{\Delta}_k)/i$	α_k
12	-6.89168e7	-5.58502e6	1.27202e11	1.02767e10	1
16	-8.81481e11	-1.54063e10	4.14120e13	6.83541e11	13/30
18	-5.07833e13	-5.43305e12	-2.19200e14	-3.36339e13	2/3
20	-4.22882e16	-3.73170e14	6.38640e16	-1.89823e14	17/5
22	-1.49974e18	-4.24489e16	1.43795e18	-4.41050e16	19/21
26	-1.18673e22	-3.60961e21	1.18653e22	-3.61016e21	23/15

Table 1: Periods $\omega^\pm(\Delta_k)$ and quasi-periods $\eta^\pm(\hat{\Delta}_k)$, and the rational number α_k . While $\alpha_{12} = 1$ matches Brown [2], α_k for $k \geq 16$ appear to be new, interesting, rational numbers, albeit with implicit dependence on the period-polynomial basis normalizations—in contrast to the normalization-free Petersson check above. We have not found a clear pattern for the α_k . See [9] for the scripts that extracted these figures (to 50-digit precision).

Brown–Hain [7, Prop. 5.6] show that $\det(\text{pd}_k)$ satisfies a Legendre-type relation, $\det(\text{pd}_k) \in \mathbb{Q}^\times \cdot (2\pi i)^{k-1}$. Brown [2, §8.3] numerically finds $\alpha_k = 1$; we reproduce this result at $k = 12$ and

give the analogous rational multiples for $k \in \{16, 18, 20, 22, 26\}$ in Table 1, below. Note: The value of α_k is tied to the normalization of Definition 3.12. Other normalizations, e.g. not including the pp-operator in defining $W_{\pm}^{(k)}$ would lead to different α_k -values.

4 Meromorphic modular forms

For $f \in F_k$ with poles in $\mathbb{H}$, the two segment integrals of Definition 2.3 depend on their contours; we show their sum depends only on the homotopy class of the pair.

Proposition 4.1. *For $f \in F_k$ and $\tau_0 \in \mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, the L-integral $L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T)$ depends only on the homotopy class of its two segments in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, and not on the basepoint τ_0 . At fixed τ_0 each of L_S^*, L_T^* is separately path-independent.*

Proof. At fixed τ_0 the integrands $f(\tau)\tau^{s-1}$ and $f(\tau)\tilde{k}_T(\tau, s)$ are holomorphic on $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, so Cauchy's theorem makes each of L_S^*, L_T^* constant on the homotopy class of its own segment. Under a motion of the basepoint, $\partial_{\tau_0} L^*$ is evaluated at the moving endpoints alone; the computation of Proposition 2.6 — which uses only $f(\tau_0 - 1) = f(\tau_0)$ and $f(-1/\tau_0) = \tau_0^k f(\tau_0)$ — cancels the boundary terms of L_S^* against those of L_T^* , the interior poles lying off the contours. Hence $\partial_{\tau_0} L^* = 0$ along any motion in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, while each summand may separately vary. $\square$

Definition 4.2. *For $f \in F_k$ and $\tau_0 \in \mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$, let $[\gamma_{\tau_0}^S], [\gamma_{\tau_0}^T]$ denote the homotopy classes of the two segments in $\mathbb{H} \setminus \Gamma \cdot f^{-1}(\infty)$.*

Thus, with $\gamma_{\tau_0}^S \in [\gamma_{\tau_0}^S]$ and $\gamma_{\tau_0}^T \in [\gamma_{\tau_0}^T]$, by Proposition 4.1 the value

$$L^*(f, s; [\gamma_{\tau_0}^S], [\gamma_{\tau_0}^T]) := L^*(f, s; \gamma_{\tau_0}^S, \gamma_{\tau_0}^T) = L_S^*(f, s; \gamma_{\tau_0}^S) + L_T^*(f, s; \gamma_{\tau_0}^T) \quad (46)$$

is independent of the chosen representative paths and independent of τ_0 (subject to natural constraints of τ_0 monodromies around singular points, which move between homotopy classes).

We would like to speak of “the” L-integral. Homotopy classes of course split this into L-integrals for contours in different homotopy classes. And, so, in what follows, we define the reference homotopy classes:

Definition 4.3 (Reference homotopy class for $f \in F_k$). *Let $f \in F_k$, let τ_0 be the base-point of $L^*(f, s)$, and let $f^{-1}(\infty)$ be the poles of f . If f has no poles on the line $\text{Im}(\tau) = 1$, choose $\delta = 0$. Else, choose $\delta > 0$ such that no pole lies in the strip $1 < \text{Im}(\tau) \leq 1 + \delta$. Set $\hat{\tau}_0^f := i(1 + \delta)$. The reference homotopy class $[\hat{\gamma}_f] = [\hat{\gamma}_f^T] + [\hat{\gamma}_f^S]$ is:*

- $[\hat{\gamma}_f^T]$: contains the horizontal segment from $T^{-1}\hat{\tau}_0^f$ to $\hat{\tau}_0^f$;
- $[\hat{\gamma}_f^S]$: contains the near-vertical path from $S^{-1}\hat{\tau}_0^f$ to $\hat{\tau}_0^f$ along the imaginary axis, taking a pole at i by principal value.

For $\delta = 0$, the reference paths match the paths $\gamma_i^S : i \rightarrow i$ and $\gamma_i^T : (i-1) \rightarrow i$ for $L^*(f, s)$ when $f \in M_k^1$.

With this definition, we may thus unambiguously write

$$L^*(f, s) := L^*(f, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]) . \quad (47)$$

Now, having fixed the reference value $L^*(f, s)$ in the reference homotopy class of Definition 4.3, consider two paths $\gamma_{\tau_0}^S$ and $\gamma_{\tau_0}^T$ which may be in any homotopy class. Further, let $X_S(p), X_T(p) \in \mathbb{Z}$ count the windings $\gamma_{\tau_0}^S, \gamma_{\tau_0}^T$ has about each pole $\tau_p \in f^{-1}(\infty)$ relative to the reference-class.

Lemma 4.4. *Let $f \in F_k$, $r_S(f, \tau_p) := \text{Res}_{\tau=\tau_p}(f(\tau)\tau^{s-1})$ and $r_T(f, \tau_p) := \text{Res}_{\tau=\tau_p}(f(\tau)\tilde{k}_T(\tau, s))$. Then,*

$$L^*(f, s; [\gamma_{\tau_0}^S], [\gamma_{\tau_0}^T]) = L^*(f, s) + 2\pi i e^{-\pi i s/2} \sum_{\tau_p \in f^{-1}(\infty)} \left[X_S(\tau_p) r_S(f, \tau_p) + X_T(\tau_p) r_T(f, \tau_p) \right]. \quad (48)$$

Proof. Two pairs whose winding data differ by one unit at a single pole p , on one segment, bound a loop encircling p once; the residue theorem adds $2\pi i$ times the residue of that segment's integrand, $f(\tau)\tau^{s-1}$ for γ^S or $f(\tau)\tilde{k}_T(\tau, s)$ for γ^T , and the prefactor $e^{-\pi i s/2}$ of Definition 2.3 multiplies through. Summing over crossings gives (48). $\square$

This content provides the basics to discuss our finite L-functions for $f \in F_k$. However, we wish to present these L-functions in terms of their q-series coefficients, just as we have done for L-functions for $f \in M_k^!$ in the previous section. To do this, we must introduce polylogarithmic projections of Ref. [3]:

This polylog projection isolates the q-series contribution of a given pole of an $f \in F_k$ into a known special function, an integral of a polylogarithm of negative index, and leaves behind an object $(I - \hat{P})f$ with that one pole removed. For our purposes, we find it essential to project-out all poles at, and above, the line $\text{Im } \tau = 1$. We must do this because the reference contours which describe our reference homotopy classes are, at a minimum, contour integrals of f across the horizontal segment from $i - 1$ to i . Were we to allow poles at/above this line, q-series diverges on the contour ($|c_f(n)| \gtrsim n^{m-1} e^{2\pi n \text{Im } \tau_p}$ for a pole of order m [3]).

Within this general context, we lay-out the starting definitions and facts:

Definition 4.5. *Let $f \in F_k$ have, at a pole τ_p of order m_p , the Laurent principal part $\sum_{m=1}^{m_p} r_{f, \tau_p}^*(m) (\tau - \tau_p)^{-m}$; the pole weights $r_{f, \tau_p}^*(m)$ are thus its Laurent coefficients. With the polylogarithm $\text{Li}_s(x) := \sum_{n \geq 1} x^n n^{-s}$, the polylog projection of f at τ_p is*

$$\hat{P}_{\tau_p} f(\tau) := \sum_{m=1}^{m_p} r_{f, \tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)}), \quad (49)$$

and $\hat{P} := \sum_{\text{Im } \tau_p \geq 1} \hat{P}_{\tau_p}$.

Lemma 4.6. *The normalized block $\frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)})$ has principal part exactly $(\tau - \tau_p)^{-m}$, so $\hat{P}_{\tau_p} f$ matches the principal part of f and $(I - \hat{P}_{\tau_p})f$ is holomorphic at τ_p .*

Proof. By the small- t expansion $\text{Li}_{1-m}(e^{-t}) = (m-1)! t^{-m} + O(1)$ with $t = -2\pi i(\tau - \tau_p)$, the block $\frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)})$ has a pure pole $(\tau - \tau_p)^{-m} + O(1)$ at τ_p , with no lower-order poles. Summing against the Laurent coefficients $r_{f, \tau_p}^*(m)$ therefore reproduces the principal part of f term by term, leaving $(I - \hat{P}_{\tau_p})f$ regular at τ_p . $\square$

Definition 4.7. *With $\hat{P}$ as in Definition 4.5, let $c_{Rf}(n)$ be the Fourier coefficients of the pole-projected form $(I - \hat{P})f$ in its expansion valid for large $\text{Im } \tau$:*

$$(I - \hat{P})f(\tau) = \sum_n c_{Rf}(n) q^n, \quad q = e^{2\pi i \tau}. \quad (50)$$

Definition 4.8. *$F(\infty)$ is the space of meromorphic functions $g : \mathbb{H} \rightarrow \mathbb{C}$ with $g(\tau + 1) = g(\tau)$, finitely many poles in each compact subset of the strip $|\text{Re } \tau| \leq \frac{1}{2}$, and bounded pole order at $q = 0$.*

Lemma 4.9. $F_k \subset F(\infty)$ and $(I - \hat{P})f \in F(\infty)$. Every pole of $(I - \hat{P})f$ has $\text{Im } \tau < 1$, with a maximal height $y_b < 1$ (set $y_b = 0$ if $(I - \hat{P})f$ is holomorphic).

Proof. A weakly holomorphic form is T -periodic, meromorphic, and $O(e^{C\text{Im}\tau})$ at the cusp, so $F_k \subset F(\infty)$; the projected polylogs $\text{Li}_{1-m}(e^{2\pi i(\tau-\tau_p)})$ share these properties, so $(I - \hat{P})f \in F(\infty)$ (and leaves F_k , being non-modular). By Lemma 4.6, $\hat{P}$ cancels the principal part of every pole with $\text{Im}\tau \geq 1$, so the poles of $(I - \hat{P})f$ lie strictly below $\text{Im}\tau = 1$; finite in each compact set and accumulating only at $\text{Im}\tau = 0$, they have a maximal height $y_b < 1$. $\square$

The Fourier coefficients of a pole-projected form are controlled by its highest pole, an estimate we import from [3]:

Lemma 4.10. With y_b as in Lemma 4.9 and m_b the maximal order among poles at height y_b , there is a $C > 0$ such that, for every $\varepsilon > 0$,

$$C n^{m_b-1} e^{2\pi n y_b} \leq |c_{Rf}(n)| \leq e^{2\pi n(y_b+\varepsilon)}, \quad (51)$$

the upper bound for all large n , the lower for infinitely many n . In particular $|c_{Rf}(n)| e^{-2\pi n} \rightarrow 0$ geometrically, at base $e^{-2\pi(1-y_b-\varepsilon)}$.

Proof. This is the pole-height/order Fourier-growth estimate of [3] for a T -periodic meromorphic function, applied to $(I - \hat{P})f$ (Lemma 4.9). $\square$

For $n \neq 0$, define

$$G_{s,k}(n) := \frac{\Gamma(s, 2\pi n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, 2\pi n)}{(2\pi n)^{k-s}}. \quad (52)$$

Each pole with $\text{Im } \tau_p \geq 1$ contributes one projected polylog to $L^*(f, s)$, whose L-integral we now record.

Definition 4.11. For a pole projected at z with $\text{Im } z > 1$, let $I_N(s, z)$ be the L-integral of the projected polylog $\text{Li}_{-N}(e^{2\pi i(-z)})$ in the minimal class $[\hat{\gamma}_f^S], [\hat{\gamma}_f^T]$ of Definition 4.3:

$$I_N(s, z) := L^*(\text{Li}_{-N}(e^{2\pi i(-z)}), s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]). \quad (53)$$

Lemma 4.12. Let $f \in F_k$ have a pole of order p at z , $\text{Im } z > 1$, with projection weights $r_{f,z}^*(m)$ (Definition 4.5). Polylog inversion $\text{Li}_{-N}(w) = (-1)^{N+1} \text{Li}_{-N}(1/w) - \delta_{N,0}$ ($N \in \mathbb{N}_0$) resums (53) into (each exponentially convergent, base $e^{2\pi(1-\text{Im}(z))}$)

$$L^*(\hat{P}_z f, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]) = \sum_{m=1}^p r_{f,z}^*(m) \frac{(-2\pi i)^m}{(m-1)!} I_{m-1}(s, z), \quad (54)$$

$$I_N(s, z) = \delta_{N,0} \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + (-1)^{N+1} \sum_{n=1}^{\infty} n^N e^{2\pi i n z} G_{s,k}(-n). \quad (55)$$

Proof. By Proposition 4.1 we may evaluate $I_N(s, z)$ in the $\delta \rightarrow 0$ representative of the minimal class; there the projected polylog — regular at i since $\text{Im } z > 1$ — has vanishing γ^S -integral and $\hat{\gamma}_f^T \rightarrow [i-1, i]$. On $[i-1, i]$ one has $|e^{2\pi i(\tau-z)}| > 1$, so the defining series of Li_{-N} diverges; inversion rewrites it as $(-1)^{N+1} \sum_{n \geq 1} n^N e^{2\pi i n(z-\tau)}$, convergent since $|e^{2\pi i(z-\tau)}| < 1$, and term-by-term integration against $\tilde{k}_T$ gives $\sum_n n^N e^{2\pi i n z} G_{s,k}(-n)$, the mode- $(-n)$ block of Theorem 3.2; for $N = 0$ the leftover -1 in $\text{Li}_0(w) = -1 - \text{Li}_0(1/w)$ contributes $\frac{1}{s} + \frac{i^k}{k-s}$. The rate is $e^{-2\pi(\text{Im } z - 1)}$, from $\Gamma(s, -2\pi n) \sim (2\pi n)^{s-1} e^{2\pi n}$ against $|e^{2\pi i n z}| = e^{-2\pi n \text{Im } z}$. $\square$

Extracting similar results for L^* evaluated on the polar terms for poles on the line $\text{Im}(\tau) = 1$ requires more care. To do this away from the special point $z = i$, we find it helpful to define the following two functions:

$$\Pi_N(s, z) := \sum_{\lambda=0}^{N+1} \frac{\text{Li}_{\lambda+1-N}(e^{2\pi i(z-i)})}{(-2\pi)^{\lambda+1}} \{(s-1)_{(\lambda)} + i^k(k-s-1)_{(\lambda)}\}, \quad (56)$$

$$R_N(s, z) := \sum_{n=1}^{\infty} n^N e^{2\pi i n z} \left[(s-1)_{(N+2)} \frac{\Gamma(s-N-2, -2\pi n)}{(-2\pi n)^s} + i^k(k-s-1)_{(N+2)} \frac{\Gamma(k-s-N-2, -2\pi n)}{(-2\pi n)^{k-s}} \right], \quad (57)$$

where $(a)_{(\lambda)} := a(a-1)\cdots(a-\lambda+1)$ is the falling factorial ($(a)_{(0)} := 1$), Li_s the polylogarithm of Definition 4.5, and $\Gamma, \Gamma(a, x)$ the gamma and upper incomplete gamma, on principal branches [10].

Lemma 4.13. *Let $f \in F_k$ have a pole of order p at z with projection weights $r_{f,z}^*(m)$ (Definition 4.5). With Π_N, R_N as in (56)–(57), for $\text{Im } z > 1$ and every $s \in \mathbb{C}$,*

$$L^*(\hat{P}_z f, s; [\hat{\gamma}_f^S], [\hat{\gamma}_f^T]) = \sum_{m=1}^p r_{f,z}^*(m) \frac{(-2\pi i)^m}{(m-1)!} I_{m-1}(s, z), \quad (58)$$

$$I_N(s, z) = \delta_{N,0} \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + (-1)^{N+1} [\Pi_N(s, z) + R_N(s, z)], \quad (59)$$

and the second line's right side extends $I_N(s, \cdot)$ continuously to $\text{Im } z = 1$ for $z \notin i + \mathbb{Z}$.

The crux of this Lemma is to show the existence of the smooth continuation of $I_N(s, z)$ from Lemma 4.12, derived for $\text{Im}(z) > 1$, to the $\text{Im}(z) = 1$ -line away from the elliptic point $z = i$. This elliptic point requires special care, which must focus on, below.

Proof. We manipulate the main result of Lemma 4.12,

$$I_N(s, z) = \delta_{N,0} \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + (-1)^{N+1} \sum_{n=1}^{\infty} n^N e^{2\pi i n z} \left(\frac{\Gamma(s, -2\pi n)}{(-2\pi n)^s} + i^k \frac{\Gamma(k-s, -2\pi n)}{(-2\pi n)^{k-s}} \right), \quad (60)$$

Our proof proceeds in two steps. First, we isolate the possible sources of divergence on the line $\text{Im}(z) = 1$ into a finite sum of polylogarithms $\text{Li}_{\lambda+1-N}(e^{2\pi i(z-i)})$ for $\lambda = 0, \dots, N+1$. To do this, we use the recurrence relation [10] for each fixed- n term,

$$\Gamma(a, x) = e^{-x} \sum_{\lambda=0}^{N+1} (a-1)_{(\lambda)} x^{a-1-\lambda} + (a-1)_{(N+2)} \Gamma(a-N-2, x) \quad (61)$$

to each gamma in $G_{s,k}(-n)$ at $x = -2\pi n$. Dividing each incomplete-Gamma term by $(-2\pi n)^s$ and $(-2\pi n)^{k-s}$ yields, then, a common factor of $1/(-2\pi n)^{\lambda+1}$ for each term in the finite sum. Further, this finite-sum is proportional to $e^{2\pi n}$; we leave the trailing incomplete gammas as-is:

$$G_{s,k}(-n) = \left\{ e^{2\pi n} \sum_{\lambda=0}^{N+1} \frac{(s-1)_{(\lambda)}}{(-2\pi n)^{\lambda+1}} + \frac{(s-1)_{(N+2)} \Gamma(s-N-2, -2\pi n)}{(-2\pi n)^s} \right\} + i^k(s \rightarrow k-s). \quad (62)$$

Now, we consider the n -sum $G_{s,k}(-n)$, starting with the falling factorial terms:

$$\sum_{n=1}^{\infty} n^N e^{2\pi i n z} e^{2\pi n} \left(\sum_{\lambda=0}^{N+1} \frac{(s-1)_{(\lambda)}}{(-2\pi n)^{\lambda+1}} + i^k(s \rightarrow k-s) \right) = \sum_{\lambda=0}^{N+1} \frac{\text{Li}_{\lambda+1-N}(e^{2\pi i(z-i)})}{(-2\pi)^{\lambda+1}} \{(s-1)_{(\lambda)} + i^k(k-s-1)_{(\lambda)}\}, \quad (63)$$

i.e., $\Pi_N(s, z)$ of (56), while the reduced-gamma terms remaining are $R_N(s, z)$ of Eq. (57), which together establish the lemma's identity (59).

The second main step of our proof is to show that $\Pi_N(s, z)$ is regular as $\text{Im}(z) \rightarrow 1$ for $z \neq i$, while $R_N(s, z)$ is regular for all z with $\text{Im}(z) = 1$. We treat each piece in turn, beginning with R_N .

Each term of R_N is internal to the $G_{s,k}(-n)$ in the mode $\sum e^{2\pi i n z} G_{s,k}(-n)$ from Lemma 4.12. Clearly, $|e^{2\pi i n z}| = e^{-2\pi n \text{Im} z}$. Thus, the incomplete gamma's exponential growth cancels against the exponentially falling $e^{2\pi i n z}$ factors. As $n \rightarrow \infty$, we have:

$$\left| n^N e^{2\pi i n z} \frac{\Gamma(s-N-2, -2\pi n)}{(-2\pi n)^s} \right| \sim \left| \frac{n^N e^{2\pi i n z} (-2\pi n)^{s-N-3} e^{2\pi n}}{(-2\pi n)^s} \right| \sim \left| \frac{e^{2\pi i n z} e^{2\pi n}}{(-2\pi)^{3+N} n^3} \right| \sim \frac{e^{2\pi n(1-\text{Im}(z))}}{(2\pi)^{N+3} n^3} \quad (64)$$

where the detailed asymptotics of $\Gamma(s, x)$ in the first scaling relation follow from [10, §8.11]. (The $i^k(s \rightarrow k-s)$ -partner follows identically.) Hence the n -th term of R_N is $O(n^{-3} e^{-2\pi n(\text{Im} z - 1)})$, uniformly for $\text{Im} z \geq 1$. There $e^{-2\pi(\text{Im} z - 1)} \leq 1$, so $\sum_n n^{-3} e^{-2\pi n(\text{Im} z - 1)} \leq \zeta(3)$: the series converges absolutely and uniformly, up to and including the edge $\text{Im} z = 1$, so R_N is continuous there.

For Π_N , recall that it is a sum over $N+2$ polylogarithms of integer index, $\Pi_N(s, z) = c_2(s) \text{Li}_2(w) + c_1(s) \text{Li}_1(w) + \dots + c_{1-N}(s) \text{Li}_{1-N}(w)$, where $w := e^{2\pi i(z-i)}$. Each Li_a is single-valued and analytic on the cut plane $\mathbb{C} \setminus [1, \infty)$: for $a \leq 0$ it is rational with a pole of order $1-a$ at $w = 1$; $\text{Li}_1(w) = -\log(1-w)$ has a logarithmic branch point at $w = 1$; and $\text{Li}_2(w)$ is finite at its branch point $w = 1$. All three singularities sit at the endpoint $w = 1$, and the cuts of Li_1, Li_2 run along the ray $[1, \infty)$. The edge is the unit circle $|w| = 1$, which meets that radial cut only at $w = 1$; approaching it from inside the disc ($\text{Im} z > 1$) gives the analytic boundary value of each Li_a at every $w \neq 1$. Hence Π_N is analytic along the whole edge except at $z \in i + \mathbb{Z}$, where each index ≤ 1 term diverges. With the continuity of R_N , (59) extends $I_N(s, \cdot)$ continuously to $\text{Im} z = 1$ for $z \notin i + \mathbb{Z}$. $\square$

The one edge point the formula misses is $z = i$, where Π_N is singular ($w = 1$); we compute $I_N(s, i)$ directly. Its finiteness rests on a relation among the Laurent coefficients of f at the S -fixed point.

Lemma 4.14. *Let $f \in F_k$ have a pole of order P at i , with pole weights $r_{f,i}^*(m)$ (Definition 4.5), i.e. principal part $\sum_{m=1}^P r_{f,i}^*(m)(\tau - i)^{-m}$. Modularity $f(-1/\tau) = \tau^k f(\tau)$ is equivalent to*

$$\sum_{m=r}^P r_{f,i}^*(m) \binom{m}{r} i^m = i^k \sum_{m=r}^P r_{f,i}^*(m) \binom{k}{m-r} i^{-m}, \quad r = 1, \dots, P. \quad (65)$$

The leading relation ($r = P$) is $((-1)^P - i^k) r_{f,i}^*(P) = 0$, so the pole order matches $k \bmod 4$ (P odd $\Leftrightarrow k \equiv 2$, P even $\Leftrightarrow 4 \mid k$); at $P = 1$ this is $(1 + i^k) r_{f,i}^*(1) = 0$, and for a double pole the $r = 1$ relation is $r_{f,i}^*(1) = \frac{i(k-2)}{2} r_{f,i}^*(2)$.

Proof. Write $\tau = i + \varepsilon$. Then $-1/\tau - i = -\varepsilon/(1 - i\varepsilon)$, so the principal part of $f(-1/\tau)$ is $\sum_m r_{f,i}^*(m)(i - \varepsilon^{-1})^m$ and that of $\tau^k f(\tau)$ is $i^k(1 - i\varepsilon)^k \sum_m r_{f,i}^*(m)\varepsilon^{-m}$. Matching the coefficient of ε^{-r} through $(i - \varepsilon^{-1})^m = \sum_j \binom{m}{j} i^{m-j} (-1)^j \varepsilon^{-j}$ and $(1 - i\varepsilon)^k = \sum_l \binom{k}{l} (-i)^l \varepsilon^l$ gives $(-1)^r \sum_{m=r}^P r_{f,i}^*(m) \binom{m}{r} i^{m-r} = i^k \sum_{m=r}^P r_{f,i}^*(m) \binom{k}{m-r} (-i)^{m-r}$; dividing through by i^r (with $-1 = i^2$ and $-i = i^{-1}$) yields (65). $\square$

Definition 4.15. For $N \geq 0$ and any cutoff $J \geq N + 2$, with $H_N := \sum_{r=1}^N \frac{1}{r}$ and $c_j^+(s) := (s-1)_{(j)} + i^k(k-s-1)_{(j)}$, set

$$\mathcal{E}_N^+(s) := \sum_{n \geq 1} n^N e^{2\pi n} \left[(s-1)_{(J)} \frac{\Gamma(s-J, 2\pi n)}{(2\pi n)^s} + i^k(k-s-1)_{(J)} \frac{\Gamma(k-s-J, 2\pi n)}{(2\pi n)^{k-s}} \right], \quad (66)$$

and then

$$\text{FP}_N(s, k) := \frac{c_N^+(s) H_N}{(2\pi)^{N+1}} + \sum_{\substack{0 \leq j \leq J \\ j \neq N}} \frac{c_j^+(s) \zeta(j+1-N)}{(2\pi)^{j+1}} + \frac{(-1)^N N!}{(2\pi)^{N+1}} \sum_{t=0}^{N-1} \frac{(-1)^t (s-1)_{(t)}}{t! (N-t)} + \mathcal{E}_N^+(s). \quad (67)$$

Lemma 4.16. Let $f \in F_k$ have a pole of order P at i with pole weights $r_{f,i}^*(m)$ (Definition 4.5), and $\hat{P}_i f$ its projection. Evaluated in the minimal class at basepoint $i(1+\delta)$, the pole's contribution to $L^*(f, s)$ has the finite $\delta \rightarrow 0$ limit

$$L^*(\hat{P}_i f, s) := \lim_{\delta \rightarrow 0} L^*(\hat{P}_i f, s; i(1+\delta)) = \sum_{m=1}^P \frac{(-2\pi i)^m r_{f,i}^*(m)}{(m-1)!} \text{FP}_{m-1}(s, k), \quad (68)$$

entire in s . The limit exists precisely because the pole weights obey the relations of Lemma 4.14 at the S -fixed point i : they cancel the individually divergent contour pieces.

Proof. Fix s ; evaluate in the minimal class at basepoint $i(1+\delta)$, $\delta > 0$, and set $u := 2\pi i(\tau - i)$.

Block expansion. At its branch point $w = 1$ the polylogarithm expands as $\text{Li}_{1-m}(e^u) = (m-1)!(-u)^{-m} + \sum_{p \geq 0} \frac{\zeta(1-m-p)}{p!} u^p$ ([10, §25.12]); multiplying by $\frac{(-2\pi i)^m}{(m-1)!}$ and using $-u = -2\pi i(\tau - i)$,

$$\frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^u) = (\tau - i)^{-m} + \frac{(-2\pi i)^m}{(m-1)!} \sum_{p \geq 0} \frac{\zeta(1-m-p)}{p!} u^p, \quad (69)$$

a pure order- m pole (matching Lemma 4.6) plus an entire, T -periodic tail. Hence $\hat{P}_i f = \sum_{m=1}^P r_{f,i}^*(m)(\tau - i)^{-m} + (\text{regular at } i)$. Because the pole lies below the T -segment (height $1+\delta$), $|e^u| < 1$ there and the block is used *without* inversion: no constant ($n=0$) mode appears, so no $\frac{1}{s} + \frac{i^k}{k-s}$ term arises and $L^*(\hat{P}_i f, s)$ is entire in s . (A pole *above* the contour would invert $\text{Li}_0 = -1 - \text{Li}_0(1/w)$, and the leftover -1 is the source of that pole in Lemma 4.12; here it is absent.)

Finite part is FP. Write $N := m-1$; the finite part collects a T -segment piece (the ζ -sum, harmonic term, and $\mathcal{E}^+$) and an S -segment piece (the polynomial term).

On γ^T the block is the convergent series $\text{Li}_{1-m}(e^u) = \sum_{n \geq 1} n^N e^{2\pi n} e^{2\pi i n \tau}$; integrating against $\tilde{k}_T$ from the basepoint $\tau_0 = i(1+\delta)$ gives, by Theorem 3.2, $\sum_{n \geq 1} n^N e^{2\pi n} \left[\frac{\Gamma(s, X_n)}{(2\pi n)^s} + i^k \frac{\Gamma(k-s, X_n)}{(2\pi n)^{k-s}} \right]$ with $X_n := -2\pi i n \tau_0 = 2\pi n(1+\delta)$. Expanding by the incomplete- Γ recursion of Lemma 4.13, $\Gamma(s, X) = e^{-X} \sum_{j=0}^{J-1} (s-1)_{(j)} X^{s-1-j} + (s-1)_{(J)} \Gamma(s-J, X)$, and using $e^{2\pi n} e^{-X_n} = e^{-2\pi n \delta}$ and $X_n^{s-1-j} = (2\pi n)^{s-1-j} (1+\delta)^{s-1-j}$, the j -sum becomes (with $c_j^+(s)$ collecting the s - and $i^k(k-s)$ -parts)

$$\sum_{j=0}^{J-1} \frac{c_j^+(s) (1+\delta)^{s-1-j}}{(2\pi)^{j+1}} \text{Li}_{j+1-N}(e^{-2\pi \delta}) + \mathcal{E}_N^+(s), \quad (70)$$

$\mathcal{E}_N^+$ being the reduced- Γ remainder of (66). With $\text{Li}_{j+1-N}(e^{-2\pi \delta}) = \Gamma(N-j)(2\pi \delta)^{j-N} + \sum_{p \geq 0} \frac{\zeta(j+1-N-p)}{p!} (-2\pi \delta)^p$, two finite (δ^0) parts survive:

- (i) the regular part at δ^0 gives, for $j \neq N$, the ζ -value $\frac{c_j^+(s)}{(2\pi)^{j+1}} \zeta(j+1-N)$ — the ζ -sum of (67);
- (ii) for $j < N$ the singular $\Gamma(N-j)(2\pi\delta)^{j-N}$ crosses the δ^{N-j} coefficient $\binom{s-1-j}{N-j}$ of $(1+\delta)^{s-1-j}$; by the falling-factorial identity $(s-1)_{(j)}(N-j-1)! \binom{s-1-j}{N-j} = (s-1)_{(N)}/(N-j)$ (and its $k-s$ partner) these sum to

$$\frac{c_N^+(s)}{(2\pi)^{N+1}} \sum_{j=0}^{N-1} \frac{1}{N-j} = \frac{c_N^+(s) H_N}{(2\pi)^{N+1}}, \quad (71)$$

the harmonic term of (67).

(The $j = N$ term is $-\frac{c_N^+(s)}{(2\pi)^{N+1}} \log(2\pi\delta) + o(1)$, a divergence with no finite part.) On γ^S the pure pole $\sum_m r_{f,i}^*(m)(\tau-i)^{-m}$, integrated by principal value against τ^{s-1} along the axis ($\tau = i(1+v)$), has the δ^0 finite part

$$i^{-m} \text{PV} \int \frac{(1+v)^{s-1}}{v^m} dv \Big|_{\delta^0} = i^{-m} (-1)^N \sum_{t=0}^{N-1} \frac{(-1)^t (s-1)_{(t)}}{t! (N-t)}, \quad (72)$$

the polynomial term of (67). Collecting, the finite part of block m is $\frac{(-2\pi i)^m}{(m-1)!} r_{f,i}^*(m) \text{FP}_{m-1}(s, k)$.

Divergences cancel by (65). The residual pieces are the $\delta \rightarrow 0$ divergences $\delta^{-(m-1)}, \dots, \log \delta$ from the pole part $\sum_m r_{f,i}^*(m)(\tau-i)^{-m}$ at the S -segment endpoints straddling i . At order r their coefficient is

$$\sum_{m=r}^P r_{f,i}^*(m) \binom{m}{r} i^m - i^k \sum_{m=r}^P r_{f,i}^*(m) \binom{k}{m-r} i^{-m}, \quad (73)$$

precisely the difference of the two sides of the modularity relation (65), hence 0 for $f \in F_k$. Every δ^{-r} and $\log \delta$ term therefore cancels and the $\delta \rightarrow 0$ limit exists; break (65) at order r and that divergence survives (at $r = 1$, a $\log \delta$).

The finite part and the (vanishing) divergences give (68). The cutoff J is immaterial, shifting terms between the ζ -sum and $\mathcal{E}_N^+$ as in Lemma 4.13. $\square$

Theorem 4.17. *Let $f \in F_k$ have poles τ_p of order P_p with $\text{Im } \tau_p \geq 1$ in the strip $|\text{Re } \tau| \leq \frac{1}{2}$, pole-projected Fourier coefficients $c_{Rf}(n)$ (Definition 4.7), and projection weights $r_{f,\tau_p}^*(m)$ (Definition 4.5). In the minimal class,*

$$L^*(f, s) = -c_{Rf}(0) \left(\frac{1}{s} + \frac{i^k}{k-s} \right) + \sum_{n \neq 0} c_{Rf}(n) G_{s,k}(n) + \sum_{\tau_p \neq i} \sum_{m=1}^{P_p} r_{f,\tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} I_{m-1}(s, \tau_p) + L^*(\hat{P}_i f, s), \quad (74)$$

with I_{m-1} from Lemmas 4.12–4.13 and the last term (present only for a pole at i) from Lemma 4.16. The right side is entire in s apart from at most simple poles at $s = 0$ and $s = k$, and reduces to the weakly-holomorphic L^* of Theorem 3.2 when f is holomorphic on $\mathbb{H}$.

Proof. Split $f = (I - \hat{P})f + \hat{P}f$. By Lemma 4.10 the Fourier series of the pole-projected part converges on $\hat{\gamma}_f^T$; Theorem 3.2 applied term by term gives its L^* as the first two sums. The projection $\hat{P}f = \sum_{\tau_p} \sum_m r_{f,\tau_p}^*(m) \frac{(-2\pi i)^m}{(m-1)!} \text{Li}_{1-m}(e^{2\pi i(\tau - \tau_p)})$ contributes, for each $\tau_p \neq i$, the L -integrals $I_{m-1}(s, \tau_p)$ of Definition 4.11, evaluated by Lemmas 4.12–4.13; the pole at i contributes $L^*(\hat{P}_i f, s)$ of Lemma 4.16. Entirety and the pole locations follow from those of $G_{s,k}$ and the $\delta_{N,0}$ terms; when f is holomorphic on $\mathbb{H}$, $\hat{P}f = 0$ and $c_{Rf} = c_f$, recovering Theorem 3.2. $\square$

References

- [1] K. Bringmann, K.-H. Fricke, and Z. Kent, *Special L-values and periods of weakly holomorphic modular forms*, Ramanujan J. **24** (2011), 119–139.
- [2] F. Brown, *A class of non-holomorphic modular forms III*, arXiv:1710.07912 (2017).
- [3] D. A. McGady, *L-functions for meromorphic modular forms and sum rules in conformal field theory*, arXiv:1806.09874 (2018).
- [4] T. M. Apostol, “Modular functions and Dirichlet series in number theory,” Graduate Texts in Mathematics, London, Springer (1997).
- [5] W. Duke and P. Jenkins, *On the zeros and coefficients of certain weakly holomorphic modular forms*, Pure Appl. Math. Q. **4** (2008), no. 4, 1327–1340.
- [6] W. Kohnen and D. Zagier, *Modular forms with rational periods*, in *Modular Forms* (R. A. Rankin, ed.), Ellis Horwood, 1984, pp. 197–249.
- [7] F. Brown and R. Hain, *Algebraic de Rham theory for weakly holomorphic modular forms of level one*, Algebra Number Theory **12** (2018), no. 3, 723–750; arXiv:1708.00652.
- [8] H. Cohen, *Haberland’s formula and numerical computation of Petersson scalar products*, in *ANTS X: Proceedings of the Tenth Algorithmic Number Theory Symposium*, Open Book Ser. **1**, Math. Sci. Publ., 2013, pp. 249–270.
- [9] *Numerical verification scripts* (companion code), 2026. Published checks: `elliptic_period_checks.py`; full working ledger: `wip_meromorphic_periods.py`.
- [10] NIST Digital Library of Mathematical Functions, F. W. J. Olver *et al.*, eds., <https://dlmf.nist.gov/>, Release 1.2.x. Chapters used: §8 (incomplete gamma) and §25.11 (Hurwitz zeta).
- [11] R. Bruggeman, Y. Choie, and N. Diamantis, *Holomorphic automorphic forms and cohomology*, Mem. Amer. Math. Soc. **253** (2018), no. 1212; arXiv:1404.6718.
- [12] K. Bringmann, P. Guerzhoy, Z. Kent and K. Ono, *Eichler-Shimura Theory for Mock Modular Forms*, Mathematische Annalen, March 2013 355(3).
- [13] V. Paşol and A. A. Popa, *Modular forms and period polynomials*, Proc. London Math. Soc. (3) **107** (2013), 713–743; arXiv:1202.5802.
- [14] W. Wang and H. Zhang, *Meromorphic quasi-modular forms and their L-functions*, arXiv:2202.09542 (2022).
- [15] F. Brown and T. J. Fonseca, *Single-valued periods of meromorphic modular forms and a motivic interpretation of the Gross–Zagier conjecture*, arXiv:2508.04844 (2025).
- [16] K. Bringmann, N. Diamantis, and S. Ehlen, *Regularized inner products and errors of modularity*, Int. Math. Res. Not. IMRN (2016); arXiv:1603.03056.
- [17] N. Diamantis and J. Drewitt, *Period functions associated to real-analytic modular forms*, Res. Math. Sci. **7** (2020), no. 21.