

Application: Flow Matching

Flow Straight and Fast:
Learning to Generate and Transfer Data with Rectified Flow

Xingchao Liu*
University of Texas at Austin
xcliu@utexas.edu

Chengyue Gong*
University of Texas at Austin
cygong@cs.utexas.edu

Qiang Liu
University of Texas at Austin
lqiang@cs.utexas.edu




Figure 2: (a) Linear interpolation of data input $(X_0, X_1) \sim \pi_0 \times \pi_1$. (b) The rectified flow Z_t induced by (X_0, X_1) ; the trajectories are “rewired” at the intersection points to avoid the crossing. (c) The linear interpolation of the end points (Z_0, Z_1) of flow Z_t . (d) The rectified flow induced from (Z_0, Z_1) , which follows straight paths.





Figure 1: The trajectories of rectified flows for image generation (π_0 : standard Gaussian noise, π_1 : cat faces, top two rows), and image transfer between human and cat faces (π_0 : human faces, π_1 : cat faces, bottom two rows), when simulated using Euler method with step size $1/N$ for N steps. The first rectified flow induced from the training data (*1-rectified flow*) yields good results with a very small number (e.g., ≥ 2) of steps; the straightened reflow induced from *1-rectified flow* (denoted as *2-rectified flow*) has nearly straight line trajectories and yield good results even with one discretization step.

Application: Modeling Cell Trajectories

Cell

Optimal-Transport Analysis of Single-Cell Gene Expression Identifies Developmental Trajectories in Reprogramming

Graphical Abstract



Authors

Geoffrey Schiebinger, Jian Shu, Marcin Tabaka, ..., Rudolf Jaenisch, Aviv Regev, Eric S. Lander

Correspondence

jianshu@broadinstitute.org (J.S.), aregev@broadinstitute.org (A.R.), lander@broadinstitute.org (E.S.L.)

In Brief

Application of a new analytical approach to examine developmental trajectories of single cells offers insight into how paracrine interactions shape reprogramming.

